Research

BMJ Open Diabetes Research & Care

Coronary artery disease in patients with atypical chest pain with and without diabetes mellitus assessed with coronary CT angiography

Marije M G Krul,¹ Kjell Bogaard,² Remco J J Knol,³ Albert C van Rossum,⁴ Paul Knaapen,⁴ Jan H Cornel,¹ Friso M van der Zant³

To cite: Krul MMG,

Bogaard K, Knol RJJ, et al. Coronary artery disease in patients with atypical chest pain with and without diabetes mellitus assessed with coronary CT angiography. *BMJ Open Diabetes Research and Care* 2014;**2**:000004. doi:10.1136/ bmjdrc-2013-000004

MMGK and KB contributed equally.

Received 18 November 2013 Revised 26 February 2014 Accepted 26 March 2014



 ¹Department of Cardiology, Medical Center Alkmaar, Alkmaar, The Netherlands
²Department of Cardiology, Rijnland Hospital, Leiderdorp, The Netherlands
³Department of Nuclear Medicine, Medical Center Alkmaar, Alkmaar, The Netherlands
⁴Department of Cardiology, VU Medical Center, Amsterdam, The Netherlands

Correspondence to

Marije MG Krul; marijekrul@hotmail.com

ABSTRACT

Introduction: Coronary artery disease (CAD) in diabetes mellitus (DM) is often widespread when diagnosed. Non-invasive coronary calcium scoring and coronary CT angiography (CAC-score/CCTA) are accurate in the detection of CAD. This study compared CAD characteristics as identified by CCTA between patients with and without DM with atypical chest pain. **Methods:** CAD was defined as CAC-score >0 and/or presence of coronary plaque. Several CAD characteristics (number of affected segments, obstructive (>50% stenosis) CAD and CAD distribution) were compared on a per patient and segment basis. Subanalysis of duration of DM (<5 or >5 years) and gender was performed.

Results: A total of 1148 patients (63.3% men, mean age 57.7±10.7), of whom 99 (8.6%) suffered from DM, were referred for CCTA. There was no difference in the prevalence of CAD between patients with and without DM (53.5% vs 50.9%, p=0.674). However patients with DM showed more affected coronary segments compared with patients without DM (2.5±3.4 vs 1.7 ±2.4, p=0.003). Multivariate analysis indicated that DM was an independent predictor of obstructive CAD (OR 2.16. 95% CI 1.23 to 3.78), as were age, women, and Diamond-Forrester score. In our study, obstructive CAD was more prevalent in women than in men (DM 40.0% vs 14.1%, p=0.003; non-DM 16.8% vs 8.4%, p<0.001). Patients suffering from DM >5 years showed more distal plaques (11.2% vs 7.7%, p=0.030). **Conclusions:** Patients with atypical chest pain and DM showed more extensive CAD, as well as more obstructive CAD, particularly in women. Diabetes duration (>5 years) was not associated with more obstructive coronary disease or different plaque morphology, although more distal disease was present.

INTRODUCTION

The number of people with diabetes mellitus (DM) is increasing due to population growth, aging, urbanization, increasing prevalence of obesity, and physical inactivity, with an estimated number of 200 million patients worldwide.¹ Patients with DM have a

Key messages

- Approximately 50% of middle-aged individuals with cardiovascular risk factors presenting with atypical chest pain will have previously undiagnosed coronary artery disease (CAD); presence of diabetes history does not affect this relationship.
- Patients with diabetes mellitus (DM) showed more affected coronary segments and more obstructive CAD than patients without DM.
- Obstructive CAD was detected more frequently in women with DM, as compared with men with DM.
- Duration of DM (>5 years) is associated with the occurrence of distal plaques.

twofold to fourfold likelihood of developing coronary artery disease (CAD) with marked morbidity and mortality.² CAD in patients with DM is often more advanced at the time of diagnosis compared with patients without DM.^{3 4}

Non-invasive cardiac imaging for the detection of CAD has evolved rapidly in the past decades. Coronary CT angiography (CCTA) offers detailed information on the extent and morphology of CAD. CCTA combined with coronary calcium scoring (CAC-score) is sensitive and specific in the detection of CAD and has been validated extensively.⁵ The real strength of CCTA is the exclusion of CAD in patients with a low-to-intermediate cardiovascular risk profile, and for this particular indication, there is a broad endorsement from international societies.⁶ Owing to technological improvements of CCTA, the radiation dose per patient has been reduced to acceptable levels. The latest generation scanners produce submillisievert (mSv) scans without compromising image quality.

The aim of the present study was to compare characteristics of CAD in patients

with atypical chest pain with and without DM using a CCTA scanner of the latest generation, and to investigate the relationship between CAD and the duration of DM.

METHODS

Study design

Consecutive patients with atypical chest pain, referred for CCTA to exclude CAD at the Medical Center Alkmaar (MCA) were entered in a database prospectively between December 13, 2011 and March 08, 2013. Patients were categorized in two groups, either with or without known DM. Type 1 and 2 DM were included. DM was diagnosed according to the American Diabetes Association criteria: fasting glucose level of \geq 7 mmol/L, symptoms of DM and casual plasma glucose of \geq 11.1 mmol/L, or the need for oral hypoglycemic agents or insulin.⁸ Only patients in sinus rhythm and without contraindications to CCTA were included. Other exclusion criteria were a history of CAD, myocardial infarction (MI), and/or revascularization procedures. All patients gave written informed consent.

The following patient characteristics were recorded: general data such as age and gender, a full cardiovascular risk profile, active medication use, Duke risk score calculation,⁹ and the Diamond-Forrester risk score calculation.¹⁰ Positive family history of premature CAD was defined as the presence of CAD in first-degree relatives younger than 55 (men) or 65 (women) years of age.¹¹ Smoking was defined as nicotine misuse within the previous 5 years. Hypertension was defined as a systolic blood pressure above 140 mm Hg and/or a diastolic blood pressure above 90 mm Hg or the use of antihypertensive drugs.¹² High cholesterol was defined as a total cholesterol >5.0 mmol/L or the use of lipid-lowering therapy.¹³ The duration of DM was categorized in two groups (<5; >5 years). Furthermore, differences between women and men were analyzed.

Patient preparation

CCTA was performed under fasting conditions. Medication could be used normally, with the exception of metformin (in case estimated glomerular filtration rate (eGFR) <60 mL/min) and sildenafil. To prevent contrast-induced nephropathy, patients with an eGFR between 30 and 60 mL/min received volume expansion therapy, using 0.9% NaCl intravenously, before and after CCTA according to the local hospital safety protocol. In case of a heart rate above 60 bpm, 100 mg atenolol was given orally 1 h prior to the CCTA. Additionally, up to 30 mg metoprolol was administered intravenously to decrease the heart rate if the heart rate still exceeded 60 bpm. Each patient also received two doses of 0.4 mg nitroglycerin sublingually.

CAC-score and CCTA data acquisition

All scans were performed with a dual source 128-slice (Somatom Definition Flash; Siemens Medical Systems, Erlangen, Germany). A 10–15 mL test bolus containing non-ionic low-osmolar iodinated radiocontrast (iopromide) was injected, followed by a flush of 40 mL saline, both at a flow rate of 6 mL/s. The time point of maximal contrast enhancement in the ascending aorta at the level of the pulmonary trunk was recorded, and an additional delay of 5 s was added to define the optimal time point for acquisition of coronary artery data. A dual-head injector then injected 48–75 mL contrast depending on kV used for the high-pitch flash scan and 75 mL in case of prospective or retrospectively triggered scan, followed by 45 mL 30/70% contrast/saline solution at a flow rate of 6 mL/s.

The tube voltage (80, 100, or 120 kV) and tube current were determined automatically by the scanning system based on body geometry.¹⁴ In this study, the total radiation dose of test bolus, topogram, CAC-score, and CCTA was used to estimate the effective radiation dose for each patient.

CAC-score and CCTA image analysis

All scans were read by two physicians, who are level 2 accredited by the Society of Computed Cardiovascular Tomography (SCCT).

For CAC-score, coronary calcifications were defined as dense lesions in coronary arteries with densities >130 Hounsfield units. Calcifications were manually assigned to coronary arteries and added to the Agatston score for each patient.¹⁵ Agatston scores were divided into three groups: a CAC-score between 0–100, 101–400, and more than 400.

Plaque characteristics were compared on a per patient and segment basis. Coronary arteries were divided into 18 segments according to the SCCT guidelines.¹⁶ Each segment was scored for the presence of coronary plaques. Structures $>1 \text{ mm}^2$ within and/or adjacent to the coronary artery lumen, which could be clearly distinguished from the vessel lumen, were scored as a coronary plaque.¹⁷ One coronary plaque was scored per coronary segment. Each coronary plaque was quantified for stenosis by visual estimation. The severity of stenosis was categorized (<25%; 25-49%; 50-69%; 70-99%; 100%). A stenosis of >50% was considered to be obstructive. The number of coronary segments with nonobstructive as well as obstructive plaques was determined for each patient. The morphology of a coronary plaque was categorized as calcified, non-calcified, or mixed lesion.

CAD was determined as CAC-score >0 and/or any coronary plaque. CAC-score=0 and no coronary plaques was defined as no CAD.

To compare the distribution of coronary plaques between patients with and without DM, patients were stratified by the location/distribution of atherosclerosis in either proximal or distal segments. The left main artery and proximal segments of the left anterior descending (LAD), right coronary artery (RCA), and ramus circumflexus (RCX) were assigned to the proximal

Table 1 Baseline, CAC-score and CCTA characteristics of study population with and without DM				
Variable	DM (n=99)	Non-DM (n=1049)	p Value	
Demographic				
Age (years)	58.6±9.9	57.7±10.8	0.435	
Sex (% male)	64 (64.6)	663 (63.2)	0.776	
Duration of DM (years)	6.6±6.9	0±0		
Risk factors				
Hypertension	73 (73.7)	676 (64.4)	0.063	
Dyslipidemia	86 (86.9)	823 (78.5)	0.049	
Obesity (BMI ≥30)	43 (43.4)	188 (17.9)	0.001	
Smoking	19 (19.2)	213 (20.3)	0.792	
Family history of CAD	53 (53.5)	518 (49.4)	0.429	
SBP (mm Hg)	138.8±17.4	133.1±17.2	0.678	
DBP(mm Hg)	81.2±12.4	80.7±11.5	0.833	
LDL cholesterol (mmol/L)	2.9±1.1	3.6±1.0	0.770	
HDL cholesterol (mmol/L)	1.2±0.3	1.4±0.4	0.003	
Total cholesterol(mmol/L)	4.9±1.4	5.7±1.1	0.060	
HbA1c (mmol/L)	7.2±1.1	5.7±0.4	0.001	
Duke score				
Low risk	36 (36.4)	599 (57.1)	0.001	
Intermediate risk	56 (56.5)	425 (40.5)	0.002	
High risk	7 (7.1)	26 (2.4)	0.007	
Diamond-Forrester score	39.5±21.6	38.2±22.2	0.562	
Medication				
ACE-I/ARB	43 (43.4)	238 (22.7)	0.001	
Aspirin	41 (41.4)	357 (34.0)	0.140	
β-blocker	52 (52.5)	437 (41.7)	0.037	
Statin	64 (64.6)	345 (32.9)	0.001	
Technical data				
Heart rate (bpm)	61.0±7.4	57.1±7.0	0.001	
Radiation dose (mSv)	3.8±2.7	2.2±1.8	0.001	
CCTA				
CAC-score	118±270.7	73.5±209.3	0.049	
0–100	74 (74.7)	866 (82.6)	0.057	
101–400	19 (19.2)	125 (11.9)	0.037	
>400	6 (6.1)	58 (5.5)	0.826	
Normal/no plaque	46 (46.5)	515 (49.1)	0.617	
CAD	53 (53.5)	534 (50.9)	0.674	
Obstructive CAD	23 (23.2)	121 (11.5)	0.001	
Non-obstructive CAD	30 (30.3)	413 (39.4)	0.076	
Segments				
Number of diseased segments	2.5±3.4	1.7±2.4	0.003	
Number of segments with obstructive plaques	0.4±1.0	0.2±0.6	0.001	
Number of segments with non-obstructive plaques	2.0±2.8	1.5±2.1	0.022	
Number of diseased segments LM	0.2±0.4	0.1±0.3	0.114	
Number of diseased segments RCA	0.7±1.1	0.4±0.8	0.008	
Number of diseased segments LAD	1.1±1.4	0.8±1.1	0.053	
Number of diseased segments RCX	0.6±1.0	0.3±0.7	0.001	

Categorical variables are expressed as count (%). Continuous variables are expressed as mean±SD.

ARB, angiotensin receptor blocker; BMI, body mass index; CAC-score, coronary calcium score; CAD, coronary artery disease; CCTA, cardiac CT coronary angiography; DBP, diastolic blood pressure; DM, diabetes mellitus; HbA1c, glycated hemoglobin; HDL, high-density lipoprotein; LAD, left anterior descending; LDL, low-density lipoprotein; LM, left main; mSv, millisievert; RCA, right coronary artery; RCX, ramus circumflexus; SBP, systolic blood pressure.

group. Mid and distal segments of LAD, RCA, and RCX and all evaluable other coronary branches were assigned to the distal group.

Statistical analysis

Continuous variables are presented as mean±SD and categorical variables as frequencies with percentages. One-sample Kolmogorov-Smirnov tests were used to test continuous data for normal distribution. Between groups, differences in mean values were tested for statistical differences with two-sample t tests and Mann-Whitney tests, when appropriate. Pearson χ^2 tests were used to test dichotomous parameters. To determine the relationship between plaque characteristics

Table 2	Baseline	characteristics	of patients	with DM with
and witho	ut CAD			

	DM with	DM without	
Variable	(n=53)	(n=46)	p Value
Demographic			
Age (years)	62.1±9.2	54.5±9.2	0.718
Sex (% male)	29 (54.7)	35 (76.1)	0.027
Duration of DM	7.1±7.4	6.0±6.3	0.468
(years)			
Risk factors			
Hypertension	29 (54.7)	17 (37.0)	0.077
Dyslipidemia	50 (94.3)	36 (78.3)	0.018
Obesity (BMI ≥30)	20 (37.7)	20 (43.5)	0.220
Smoking	8 (15.1)	11 (23.9)	0.266
Family history of CAD	28 (52.8)	25 (54.3)	0.880
Medication	()	· · · ·	
ACE-I/ARB	28 (52.8)	15 (32.6)	0.043
Aspirin	26 (49.1)	15 (32.6)	0.098
β-blocker	30 (56.6)	22 (47.8)	0.383
Statin	39 (73.6)	25 (54.3)	0.046
Technical data	()	· · · ·	
Heart rate (bpm)	59.3±6.6	62.7±8.0	0.849
Radiation dose (mSv)	3.6±2.9	4.0±2.6	0.640
Categorical variables are expressed as count (%). Continuous			

variables are expressed as mean±SD. ARB, angiotensin receptor blocker; BMI, body mass index; CAD,

coronary artery disease; DM, diabetes mellitus; mSv, millisievert.

and DM, logistic regression analyses were performed when the dependent variable was dichotomous. When the dependent variable was continuous, linear regression analyses were performed. Univariate analyses were performed using demographic variables, classical risk factors, risk scoring systems, and medication used (variables specified in table 1). Multivariable analyses were performed with variables that were significant in univariate analyses. p Values <0.05 were considered significant. The Duke risk score was not analyzed in the univariate and multivariate analyses because DM is one of the parameters in this particular score. The Diamond-Forrester score, not using DM as a parameter, was used instead. For the detailed per segment analysis, the unit of measure was each coronary segment and there were no adjustments or corrections made for the serial correlation between segments. Statistical Package for Social Sciences V.20 (SPSS Inc, Chicago, Illinois, USA) was used.

RESULTS

Study population

A total of 1176 patients with atypical chest pain were enrolled in the study, of which 28 patients were excluded. Twenty-three patients were excluded due to a history of CAD, MI, and/or revascularization procedures. Five patients suffered from arrhythmias during scan acquisition which resulted in unacceptable image quality, and thus, these patients were excluded. Table 1

summarizes the characteristics of patients with and without DM at the time of the CCTA. In total, 1148 patients (727 (63.3%) men; age (mean 57.7 years±10.7)) were included, of whom 99 (8.6%) were patients with known DM (mean 6.6 years ±6.9). Of the study population, 95.4% had a low-to-intermediate probability for CAD according to their Duke risk score (92.9% DM vs 97.9% non-DM, p=0.007). Patients with DM were more frequently obese and used more ACE-I/angiotensin receptor blocker, β-blocker, and statin therapy and had lower high-density lipoprotein levels (p values 0.001, 0.001, 0.037, 0.001, and 0.003, respectively). Table 2 compares the clinical and radiographic characteristics of patients with DM who did or did not have CAD. Patients with DM without CAD were more often of the male gender, showed less frequent dyslipidemia, and used less statin therapy (p values 0.020, 0.018, and 0.046, respectively). None of the other measured clinical variables differed between patients with DM with or without CAD.

CAD on CCTA

Coronary plaque characteristics on CCTA in patients with and without DM are shown in table 1. A normal CCTA was found in 561 (48.9%) patients, with no significant difference in prevalence between patients with and without DM (p=0.617). Patients with DM had a significantly higher CAC-score than patients without DM (118 ± 270 vs 73 ± 209 , p=0.049). Patients with DM had a significantly higher percentage of CAC-scores between 101 and 400 (19% vs 11.9%, p=0.037). Obstructive CAD was observed in 144 (12.5%) patients and more frequently in patients with DM (23.2% vs 11.5%, p<0.001). Higher CAC-scores were detected in patients with obstructive CAD (287.3±332.9 vs 47.2±173.4, p<0.001). The average number of diseased segments was higher in patients with DM compared with patients without DM (p=0.003). The number of segments with obstructive CAD plaques was higher in patients with DM compared with patients without DM. In patients with DM, the average number of diseased segments in the RCA (p=0.008) and RCX (p<0.001) was higher than in patients without DM. The difference in the average amount of diseased segments in the LAD did not reach statistical significance (p=0.053).

Figure 1 shows the prevalence of obstructive CAD across the subgroup gender. Women had an overall higher prevalence of obstructive CAD than men. In patients with DM, 40% of the women demonstrated obstructive CAD compared with 14.1% of the men (p=0.003). In patients without DM, 16.8% of the women compared with 8.4% of the men showed obstructive CAD (p<0.001).

Relationship of CAD characteristics on CCTA and DM

The results of the univariate and mulitivariate analyses of the correlation between plaque characteristics on CCTA and the presence of DM are listed in table 3. After correction for baseline characteristics, DM proved



Figure 1 Prevalence of obstructive coronary artery disease (CAD) across the different diabetes mellitus and gender subgroups.

to be an independent predictor of the total number of segments with plaque (p=0.016). The other independent predictors of the total number of segments with plaque were age (p<0.001), women (p<0.001), Diamond-Forrester score (p<0.001), and hypertension (p=0.023). Furthermore, DM was an independent predictor of the presence of obstructive plaque (OR 2.16, 95% CI 1.23 to 3.78), as were age (OR 1.04; 95% CI 1.02 to 1.07), women (OR 4.95; 95% CI 2.96 to 8.28), and Diamond-Forrester score (OR 1.02; 95% CI 1.00 to 1.03). DM was also an independent predictor of the number of segments with plaque in the distal segments (p=0.010), as were age (p<0.001), women (p<0.001), and Diamond-Forrester score (p=0.008).

Per segment analysis

A total of 17 240 segments were identified in the study population. After exclusion of 105 (0.61%) segments due to non-diagnostic image quality, a total of 17 135 coronary segments were included in the analysis. In the remaining 17 135 segments, 1998 (11.7%) segments contained plaque. Of the 17 135 segments, there were 243 (1.4%) segments with obstructive plaque, (44 of 1471 (3.0%) segments in patients with DM, 199 of 15 664 (1.3%) in patients without DM, p<0.001).

Accordingly, the distribution of plaque types differed between DM and non-DM: mixed plaques (13.3% vs 9.2%, p<0.001), calcified plaques (2.0% vs 1.3%, p=0.015), and non-calcified plaques (1.4% vs 0.8%, p=0.023). Patients with DM showed significantly more affected segments than patients without DM in the RCA (4.4% vs 2.7%, p<0.001), LAD (7.3% vs 5.6%, p=0.012), and RCX (3.8% vs 2.1%, p<0.001).

Subgroup analysis for duration of DM

Table 4 shows the subgroup analysis for duration of DM in the per segment analysis. Duration of DM was categorized into two groups (<5; >5 years). Although patients suffering from DM for more than 5 years had a higher percentage of involvement of the distal segments (11.2% vs 7.8%, p=0.030), duration of diabetes had minimal impact on total number of segments with plaque, number of obstructive lesions, and morphology of plaques.

DISCUSSION

In the present study, differences in characteristics of CAD between patients with and without DM were detected by means of CCTA. A significant, positive correlation between the presence of DM and total number of segments with plaque was demonstrated. Patients with DM had a significantly higher percentage of obstructive plaques, and this was particularly true for women. Patients suffering from DM for more than 5 years did not demonstrate more obstructive coronary disease or different plaque morphology, although more distal disease was present.

Table 3 Estimates of correlation of CCTA plaque characteristics with the presence of DM				
Variable	Univariate, parameter estimate	p Value	Multivariate, parameter estimate	p Value
Total calcium score	22.25	0.049	16.88	0.127
Total segments with plaque	0.394	0.003	0.292	0.016
CAD	1.11 (0.74 to 1.70)	0.617		
Obstructive plaque	2.32 (1.40 to 3.84)	0.001	2.16 (1.23 to 3.78)	0.007
Non-obstructive plaque	0.67 (0.43 to 1.05)	0.078		
LM (yes or no)	1.54 (0.90 to 2.70)	0.107		
RCA (yes or no)	1.45 (0.93 to 2.60)	0.101		
LAD (yes or no)	1.22 (0.81 to 1.85)	0.337		
RCX (yes or no)	1.59 (1.01 to 2.49)	0.044	1.36 (0.82 to 2.26)	0.229
Proximal plaques	0.142	0.019	0.089	0.110
Distal plaques	0.253	0.002	0.203	0.010

Data are ORs (95% CI) or estimates of correlation.

CAD, coronary artery disease; CCTA, cardiac CT coronary angiography; DM, diabetes mellitus; LAD, left anterior descending; LM, left main; RCA, right coronary artery; RCX, ramus circumflexus.

plaques—per segment analysis in subgroup duration of DM				
	DM <5 years (n=53; 797 segments)	DM >5 years (n=46; 683 segments)	p Value	
Total segments	790 (99.1)	681 (99.7)	0.149	
Total segments with plaque	121 (15.3)	125 (18.4)	0.108	
Obstructive plaque	23 (2.9)	21 (3.1)	0.847	
Non-obstructive plaque	98 (12.4)	104 (15.3)	0.111	
Plaque in LM	12 (1.5)	6 (0.9)	0.267	
Plaque in RCA	32 (4.1)	32 (4.7)	0.544	
Plaque in LAD	53 (6.7)	55 (8.1)	0.316	
Plaque in RCX	24 (3.0)	32 (4.7)	0.097	
Proximal plaque	59 (7.5)	49 (7.2)	0.841	
Distal plaque	62 (7.8)	76 (11.2)	0.030	
Mixed plaque	96 (12.2)	99 (14.5)	0.179	
Calcified plaque	15 (1.9)	15 (2.2)	0.681	
Non-calcified	10 (1.3)	11 (1.6)	0.573	

Prevalence, extent, localization, and morphology of

plaque

Categorical variables are expressed as count (%).

DM, diabetes mellitus; LAD, left anterior descending; LM, left

main; RCA, right coronary artery; RCX, ramus circumflexus.

Extent of CAD

Table 4

The present study showed a positive correlation between the presence of DM and the total number of segments with plaque. Patients with DM showed more affected segments than patients without DM. Higher prevalence of plagues in patients with DM has been reported in previous invasive as well as postmortem studies. Pundziute et al^{18} reported observations in 60 patients (including 19) with DM) using CCTA followed by conventional coronary angiography with intravascular ultrasound (IVUS); the authors demonstrated that patients with DM had significantly more affected segments on CCTA and with IVUS. In a postmortem study, Burke *et al*¹⁹ showed that patients with DM have a larger amount of total and distal plaques. In our study, in patients with atypical chest pain, DM was also associated with more obstructive CAD. This is in line with an earlier study performed by Chu *et al*²⁰ who demonstrated more obstructive plaques in symptomatic patients with DM as compared with patients without DM (2.5±2.5 vs 1.1±1.4). The CAC-score in our study was higher in patients with DM, compared with patients without DM (118±270.7 vs 73.5±209.3, p=0.049). This is in line with an earlier study by de Araujo *et al*²¹, who showed a significant difference in (median) CAC-score between patients with atypical chest pain with and without DM (68(0-311) vs 0 (0-67)), p<0.001). While our study looked at CAD in patients with atypical chest pain and DM, several studies have demonstrated CAD in asymptomatic patients with DM. Scholte *et al*²² revealed a high prevalence of CAD in asymptomatic patients with type 2 DM, with almost 80% showing CAD. Zeina et al also showed a higher

already developed serious CAD at the time of examination. One explanation could be that DM had existed long before it was diagnosed, due to lack of typical symptoms of the disease, and cardiovascular damage could have developed meanwhile.⁴ The previously mentioned studies demonstrated this in asymptomatic patients with DM.

Furthermore, in the present study women had a higher percentage of obstructive CAD in the DM and non-DM group. This is in contrast with several previous studies with CCTA that showed more obstructive CAD in men.^{21 25} These apparently conflicting results might be explained by the increasing notion of, and attention to, the atypical and late presentation of cardiac symptoms in women.^{26 27} On average, clinical symptoms of heart disease appear 10 years later in women than in men.²⁶ Studies in the Women's Ischemia Syndrome Evaluation (WISE) study support the concept of a multifactorial model where sex hormones interact with traditional and novel risk factors. Especially inflammatory process biomarkers (such as C reactive protein), leading to an increase in the functional expression of atherosclerotic plaque and vascular or metabolic alterations result in worse outcomes for women.²⁸ Also, medical diagnosis and treatment for stable angina is different in men and women.²⁹ Women were less likely to be referred for testing, in particular for exercise testing and invasive angiography. Antiplatelet and statin therapies were used significantly less in women than in men at the initial assessment and at 1 year, even in those in whom CAD had been confirmed, which unfortunately influences outcome.

Duration of DM

Previous studies have shown that the duration of type 2 DM correlates with all-cause mortality and significantly increases cardiovascular mortality.^{30 31} An autopsy study demonstrated an association between DM duration and the extent of atherosclerosis and myocardial lesions.³² Clarkson *et al^{33}* showed that vascular reactivity, a marker of impaired endothelial dysfunction, is impaired in patients with long-term type 1 DM. Also the degree of impairment is related to the duration of DM. Thus, one may speculate that longer exposure to hyperglycemia may increase the risk of endothelial dysfunction and thus increase the risk of developing CAD. To the best of our knowledge, no studies researched the correlation between the duration of DM and distal plaques. In our study, we found that patients suffering from DM for more than 5 years have more distal plaques, although

they did not demonstrate more obstructive coronary disease or different plaque morphology. Patients suffering from DM for a prolonged period will have more extensive and widespread disease.

Radiation dose CCTA

Bell *et al*^{β 4} showed in a prospective study that iodinated contrast, β -blockers, and nitroglycerin as part of a predefined CCTA protocol are safe and associated with a low rate of adverse events. In the past decade, studies have focused on radiation dose-reduction techniques with CCTA. The latest generation CCTA scanners can provide an effective dose below 1.0 mSv.⁷ In comparison, patient dose is typically 2.4 mSv for a $^{13}NH_3$ positron emission tomography scan, 2.5-5 mSv for diagnostic invasive selective coronary angiography, and 5-16 mSv for nuclear myocardial perfusion imaging by means of single photon emission CT.^{35–37} Safety and low radiation exposure with a high negative predictive value makes CCTA a very appropriate imaging technique for the exclusion of CAD in symptomatic patients and perhaps even as a presymptomatic screening tool. The radiation dose of CCTA in this study is higher in patients with DM compared with patients without DM. This difference could be explained by the fact that patients with DM are more obese compared with patients without DM. Obese patients require a higher tube voltage (kV) and current (mAs) to produce a diagnostic CCTA, which leads to a higher radiation dose.

Clinical implication

The combination of CAD and DM strongly increases cardiovascular mortality.^{30 31} The present study showed that a clinically significant number of patients with atypical chest pain and DM already had developed (serious) CAD at the time of first examination. Previous studies with asymptomatic patients also demonstrated a higher prevalence (80–93%) of CAD in patients with DM.² There is extensive evidence of prognostic information using CAC-score in symptomatic and asymptomatic patients.³⁸ However, the additional value of performing CCTA in these patients has been a matter of debate. In the CONFIRM registry, CCTA was unable to predict death of non-fatal MI in 7500 patients.³⁹ Nonetheless, in symptomatic patients the incremental prognostic value of CCTA has been established.⁴⁰ The best CCTA parameter for prediction of mortality was the number of proximal segments with mixed or calcified plaques.⁴¹ Thus, CCTA is useful in detection of CAD in patients with DM, either with or without symptoms of atypical chest pain. The high diagnostic yield in symptomatic patients with DM may justify the use of CCTA in daily clinical practice and can be considered as a criterion for appropriate use. Since prognostic value of performing CCTA in asymptomatic patients is unclear, further prospective studies are needed.

Limitations

Several limitations should be mentioned. First, this is a single-center study with a medium-size cohort in which the percentage of patients with DM was relatively small. Also, the examinations were performed at a single time point. There were some differences in the baseline characteristics of the two subgroups of patients, like obesity, and medication use, that could have contributed to the higher disease extent observed in patients with DM. Nevertheless, after adjusting for those differences, DM remained an independent predictor of plaque burden and obstructive CAD. Since patients referred for CCTA were suffering from atypical chest pain, but had no history of clinically diagnosed CAD, a potential referral bias has to be taken into account. In addition, limitations of CCTA in general should be mentioned; the presence of ischemia cannot be determined on CCTA, and abnormal CCTA findings should ideally be combined with functional data.

CONCLUSIONS

Although the prevalence of CAD in patients with atypical chest pain and DM did not differ from that in non-DM, patients with DM with CAD had more advanced CAD than patients without DM. They showed more affected coronary segments and more obstructive CAD than patients without DM; the latter especially in women with DM. Duration of DM (>5 years), in our study population, was not associated with more obstructive coronary disease or a different plaque morphology, although more distal disease was present.

Contributors MMGK and KB researched data and wrote the manuscript. RJJK researched data and reviewed/edited the manuscript. ACvR reviewed/edited the manuscript. PK and JHC contributed to the discussion and reviewed/ edited the manuscript. FMvdZ researched data, contributed to the discussion, and reviewed/edited the manuscript

Funding This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None.

Ethics approval The patients were prospectively included in the study and all patients signed informed consent. All patients did sign an informed consent for usage of their data.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 3.0) license, which permits others to distribute, remix, adapt, build upon this work noncommercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: http:// creativecommons.org/licenses/by-nc/3.0/

REFERENCES

- 1. International Diabetes Federation. International Diabetes Federation. 2013.
- Rana JS, Dunning A, Achenbach S, *et al.* Differences in prevalence, extent, severity, and prognosis of coronary artery disease among patients with and without diabetes undergoing coronary computed tomography angiography: results from 10,110 individuals from the CONFIRM (COronary CT Angiography EvaluatioN For Clinical

Cardiovascular and metabolic risk

Outcomes): an InteRnational Multicenter Registry. *Diabetes Care* 2012;35:1787–94.

- Goraya TY, Leibson CL, Palumbo PJ, et al. Coronary atherosclerosis in diabetes mellitus: a population-based autopsy study. J Am Coll Cardiol 2002;40:946–53.
- Hu FB, Stampfer MJ, Haffner SM, *et al.* Elevated risk of cardiovascular disease prior to clinical diagnosis of type 2 diabetes. *Diabetes Care* 2002;25:1129–34.
- Petcherski O, Gaspar T, Halon DA, et al. Diagnostic accuracy of 256-row computed tomographic angiography for detection of obstructive coronary artery disease using invasive quantitative coronary angiography as reference standard. Am J Cardiol 2013;111:510–15.
- 6. Taylor AJ, Cerqueira M, Hodgson JM, et al. ACCF/SCCT/ACR/AHA/ ASE/ASNC/NASCI/SCAI/SCMR 2010 appropriate use criteria for cardiac computed tomography. A report of the American College of Cardiology Foundation Appropriate Use Criteria Task Force, the Society of Cardiovascular Computed Tomography, the American College of Radiology, the American Heart Association, the American Society of Echocardiography, the American Society of Nuclear Cardiology, the North American Society of Cardiovascular Imaging, the Society for Cardiovascular Angiography and Interventions, and the Society for Cardiovascular Magnetic Resonance. J Am Coll Cardiol 2010;56:1864–94.
- Achenbach S, Marwan M, Ropers D, et al. Coronary computed tomography angiography with a consistent dose below 1 mSv using prospectively electrocardiogram-triggered high-pitch spiral acquisition. Eur Heart J 2010;31:340–6.
- American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care* 2009;32(Suppl 1):S62–7.
- 9. Wasfy MM, Brady TJ, Abbara S, *et al.* Comparison of the Diamond-Forrester method and Duke Clinical Score to predict obstructive coronary artery disease by computed tomographic angiography. *Am J Cardiol* 2012;109:998–1004.
- Diamond GA, Forrester JS. Probability of CAD. *Circulation* 1982;65:641–2.
- Taylor AJ, Bindeman J, Feuerstein I, et al. Coronary calcium independently predicts incident premature coronary heart disease over measured cardiovascular risk factors: mean three-year outcomes in the Prospective Army Coronary Calcium (PACC) project. J Am Coll Cardiol 2005;46:807–14.
- European Society of Hypertension-European Society of Cardiology Guidelines Committee. 2003 European Society of Hypertension-European Society of Cardiology guidelines for the management of arterial hypertension. J Hypertens 2003;21:1011–53.
- Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults. Executive Summary of The Third Report of The National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). JAMA 2001;285:2486–97.
- 14. Lee KH, Lee JM, Moon SK, *et al.* Attenuation-based automatic tube voltage selection and tube current modulation for dose reduction at contrast-enhanced liver CT. *Radiology* 2012;265:437–47.
- Agatston AS, Janowitz WR, Hildner FJ, et al. Quantification of coronary artery calcium using ultrafast computed tomography. J Am Coll Cardiol 1990;15:827–32.
- 16. Raff GL, Abidov A, Achenbach S, *et al.* SCCT guidelines for the interpretation and reporting of coronary computed tomographic angiography. *J Cardiovasc Comput Tomogr* 2009;3:122–36.
- Leber AW, Knez A, Becker A, *et al.* Accuracy of multidetector spiral computed tomography in identifying and differentiating the composition of coronary atherosclerotic plaques: a comparative study with intracoronary ultrasound. *J Am Coll Cardiol* 2004;43:1241–7.
- Pundziute G, Schuijf JD, Jukema JW, et al. Type 2 diabetes is associated with more advanced coronary atherosclerosis on multislice computed tomography and virtual histology intravascular ultrasound. J Nucl Cardiol 2009;16:376–83.
- Burke AP, Kolodgie FD, Zieske A, et al. Morphologic findings of coronary atherosclerotic plaques in diabetics: a postmortem study. Arterioscler Thromb Vasc Biol 2004;24:1266–71.
- Chu ZG, Yang ZG, Dong ZH, *et al.* Comparative analysis of coronary artery disease assessed by coronary CT angiography between patients with type 2 diabetes mellitus and non-diabetic patients. *Int J Cardiol* 2011;147:178–81.
- de Araujo GP, Garcia-Garcia HM, Carvalho MS, et al. Diabetes as an independent predictor of high atherosclerotic burden assessed by coronary computed tomography angiography: the coronary artery

disease equivalent revisited. Int J Cardiovasc Imaging 2012;29:1105–14.

- Scholte AJ, Schuijf JD, Kharagjitsingh AV, et al. Prevalence of coronary artery disease and plaque morphology assessed by multi-slice computed tomography coronary angiography and calcium scoring in asymptomatic patients with type 2 diabetes. *Heart* 2008;94:290–5.
- Zeina AR, Odeh M, Rosenschein U, et al. Coronary artery disease among asymptomatic diabetic and nondiabetic patients undergoing coronary computed tomography angiography. Coron Artery Dis 2008;19:37–41.
- Rivera JJ, Nasir K, Choi EK, *et al.* Detection of occult coronary artery disease in asymptomatic individuals with diabetes mellitus using non-invasive cardiac angiography. *Atherosclerosis* 2009;203:442–8.
- Qureshi W, Blaha MJ, Nasir K, *et al.* Gender differences in coronary plaque composition and burden detected in symptomatic patients referred for coronary computed tomographic angiography. *Int J Cardiovasc Imaging* 2013;29:463–9.
- McSweeney JC, Cody M, O'Sullivan P, et al. Women's early warning symptoms of acute myocardial infarction. *Circulation* 2003;108:2619–23.
- Goldberg RJ, O'Donnell C, Yarzebski J, et al. Sex differences in symptom presentation associated with acute myocardial infarction: a population-based perspective. Am Heart J 1998;136:189–95.
- Shaw LJ, Bairey Merz CN, Pepine CJ, et al. Insights from the NHLBI-Sponsored Women's Ischemia Syndrome Evaluation (WISE) Study: part I: gender differences in traditional and novel risk factors, symptom evaluation, and gender-optimized diagnostic strategies. J Am Coll Cardiol 2006;47:S4–20.
- Daly C, Clemens F, Lopez Sendon JL, et al. Gender differences in the management and clinical outcome of stable angina. *Circulation* 2006;113:490–8.
- Silbernagel G, Rosinger S, Grammer TB, et al. Duration of type 2 diabetes strongly predicts all-cause and cardiovascular mortality in people referred for coronary angiography. Atherosclerosis 2012;221:551–7.
- Fox CS, Sullivan L, D'Agostino RB Sr, et al. The significant effect of diabetes duration on coronary heart disease mortality: the Framingham Heart Study. Diabetes Care 2004;27:704–8.
- Burchfiel CM, Reed DM, Marcus EB, et al. Association of diabetes mellitus with coronary atherosclerosis and myocardial lesions. An autopsy study from the Honolulu Heart Program. Am J Epidemiol 1993;137:1328–40.
- Clarkson P, Celermajer DS, Donald AE, et al. Impaired vascular reactivity in insulin-dependent diabetes mellitus is related to disease duration and low density lipoprotein cholesterol levels. J Am Coll Cardiol 1996;28:573–9.
- Bell GW, Edwardes M, Dunning AM, *et al.* Periprocedural safety of 64-detector row coronary computed tomographic angiography: results from the prospective multicenter ACCURACY trial. *J Cardiovasc Comput Tomogr* 2010;4:375–80.
- Hoffmann U, Ferencik M, Cury RC, et al. Coronary CT angiography. J Nucl Med 2006;47:797–806.
- Kajander S, Ukkonen H, Sipila H, *et al.* Low radiation dose imaging of myocardial perfusion and coronary angiography with a hybrid PET/CT scanner. *Clin Physiol Funct Imaging* 2009;29:81–8.
- Stabin MG. Radiopharmaceuticals for nuclear cardiology: radiation dosimetry, uncertainties, and risk. J Nucl Med 2008;49:1555–63.
- Hecht HS, Narula J. Coronary artery calcium scanning in asymptomatic patients with diabetes mellitus: a paradigm shift. *J Diabetes* 2012;4:342–50.
- Cho I, Chang HJ, Sung JM, et al.; CONFIRM Investigators. Coronary computed tomographic angiography and risk of all-cause mortality and nonfatal myocardial infarction in subjects without chest pain syndrome from the CONFIRM Registry (coronary CT angiography evaluation for clinical outcomes: an international multicenter registry). *Circulation* 2012;126:304–13.
- Al-Mallah MH, Qureshi W, Lin FY, *et al.* Does coronary CT angiography improve risk stratification over coronary calcium scoring in symptomatic patients with suspected coronary artery disease? Results from the prospective multicenter international CONFIRM registry. *Eur Heart J Cardiovasc Imaging* 2014; 15:267–74.
- Nance JW Jr, Schlett CL, Schoepf UJ, et al. Incremental prognostic value of different components of coronary atherosclerotic plaque at cardiac CT angiography beyond coronary calcification in patients with acute chest pain. *Radiology* 2012;264:679–90.