BMJ Open Diabetes Research & Care

Inclusion of walnut in the diets of adults at risk for type 2 diabetes and their dietary pattern changes: a randomized, controlled, cross-over trial

Valentine Yanchou Njike,^{1,2} Niloufarsadat Yarandi,² Paul Petraro,² Rockiy G Ayettey,^{1,2} Judith A Treu,^{1,2} David L Katz^{1,2}

ABSTRACT

To cite: Njike VY, Yarandi N, Petraro P, *et al.* Inclusion of walnut in the diets of adults at risk for type 2 diabetes and their dietary pattern changes: a randomized, controlled, cross-over trial. *BMJ Open Diabetes Research and Care* 2016;**4**:e000293. doi:10.1136/bmjdrc-2016-000293

Received 5 July 2016 Revised 26 August 2016 Accepted 12 September 2016



¹Yale University Prevention Research Center, Derby, Connecticut, USA ²Griffin Hospital—Derby, Derby, Connecticut, USA

Correspondence to Dr David L Katz; davkatz7@gmail.com **Background:** In our recently published study, including walnuts in the diets of adults with prediabetes led to overall improvement in diet quality. This report adds to those study findings by examining the food groups displaced during walnut inclusion in the diets of those adults with prediabetes. **Methods:** Randomized, controlled, modified Latin

square parallel design with 2 treatment arms. The 112 participants (31 men, 81 women) were randomly assigned to a diet with or without dietary counseling to regulate calorie intake in a 1:1 ratio. Within each treatment arm, participants were further randomized to 1 of 2 sequence permutations to receive a walnutincluded diet with 56 g (366 kcal) of walnuts per day and a walnut-excluded diet. Participants in the calorieregulated arm received advice from a dietitian to preserve an isocaloric condition while including walnuts. We analyzed the 12 components of the 2010 Healthy Eating Index to examine dietary pattern changes of study participants.

Results: Seafood and plant protein foods intake significantly increased with walnut inclusion, compared with their exclusion (2.14 \pm 2.06 vs $-0.49\pm$ 2.33; p=0.003). The ingestion of healthful fatty acids also significantly increased with walnut inclusion, compared with their exclusion (1.43 \pm 4.53 vs $-1.76\pm$ 4.80; p=0.02). Dairy ingestion increased with walnut inclusion in the calorie-regulated phase, compared with walnut inclusion without calorie regulation (1.06 \pm 4.42 vs $-2.15\pm$ 3.64; p=0.02).

Conclusions: Our data suggest that walnut inclusion in the diets of adults at risk for diabetes led to an increase in intake of other healthful foods. **Trial registration number:** NCT02330848.

BACKGROUND

According to the WHO, the number of cases of adults with diabetes has quadrupled since 1980.¹ This increase in the number of cases of diabetes could be attributed to changes in dietary patterns which include an increase in the consumption of obesigenic foods such as sugary foods and beverages.² Currently,

Key messages

 The inclusion of walnuts in the diets of adults at risk for type 2diabetes improved their overall diet quality.

Research

- The inclusion of walnuts in the diets of adults at risk for type 2 diabetes facilitated the introduction of healthy food choices in their diets.
- The inclusion of walnuts in the diets of adults at risk for type 2 diabetes facilitated the replacement of unhealthy foods in their diets.

about 29.1 million Americans are known to have diabetes.³ An estimated 86 million Americans aged 20 and older have prediabetes.³ Among adults at risk of diabetes, 15–30% develop diabetes within 5 years.³

Potential complications of diabetes include heart attack, stroke, vision loss, kidney disease, nerve damage, and lower limb gangrene that could lead to amputation. The risk of death of individuals with diabetes is 50% more than that of those without diabetes. It is estimated that the total medical cost, work lost, and wages lost for individuals with diabetes is about \$245 billion. The medical cost for individuals with diabetes is twice as high as that for those without diabetes.³

Although there is no known cure for diabetes, the progression from prediabetes to type 2 diabetes could be prevented.4-7 A study by Knowler *et al*⁴ demonstrated that modest lifestyle changes (ie, exercise and nutrition) led to a reduction of 5-7% of body weight and a 58% reduction in the risk of developing type 2 diabetes among individuals at high risk of diabetes. In addition, lifestyle interventions have been shown to be cost-effective for the primary prevention of type 2 diabetes.⁸ Foods that are low in fat and calories, and high in fiber, with a focus on fruit, vegetables and whole grains are typically recommended to prevent the

progression from prediabetes to diabetes.⁹ Improved diet quality has been associated with lower risk of type 2 diabetes.¹⁰

In our recently published study,¹² the inclusion of walnuts in the diets of adults at risk for type 2 diabetes led to improvement in their overall diet quality. The foods displaced from the diets of these individuals with the inclusion of walnuts that led to the improvement in diet quality were unknown. We therefore examined the foods that were displaced with the inclusion of walnuts in the diets of these adults with prediabetes.

METHODS

Study population

One hundred and twelve adults at risk for type 2 diabetes (31 men, 81 women) were recruited from the Lower Naugatuck Valley of Connecticut through flyers and newspaper advertisements. Those responding to the advertisements (n=678) were prescreened using a semi-structured telephone interview. Details of the study methods, recruitment process, and participant flow have been previously published.¹²

Inclusion criteria: Adults between 25 and 75 years of age; non-smokers; high risk for diabetes, defined as meeting at least one of the following criteria: (1) overweight with increased waist circumference; (2) prediabetes based on fasting blood glucose between 100 and 126 mg/dL or glycated hemoglobin 5.7–6.4%; or (3) metabolic syndrome.

Exclusion criteria: Allergy to walnuts or any other nuts; anticipated inability to complete the study protocol for any reason; current eating disorder; restricted diets by choice (ie, vegetarian, vegan); regular use of high doses of vitamin E (>400 IU/day) or vitamin C (>500 mg/ day); use of fish oil, flaxseed oil, omega-3 fatty acid, and/or fiber supplements unless willing to discontinue supplementation for the study duration.

Study design

This was a randomized, controlled, modified Latin square parallel design study with two treatment arms to examine the effects of the inclusion of walnuts in the diets of adults at risk for type 2 diabetes on their dietary patterns over a 6-month period. After a 4-week run-in period of an ad libitum diet, participants were randomized to one of four possible sequence permutations, and then crossed over following inclusion of 56 g walnuts per day, or walnut exclusion, for 6 months in their diets in random sequence, with a 3-month washout between treatments.

Intervention

Walnut intake without calorie regulation: Participants were provided 392 g of walnuts (unshelled) per week (56 g or 2 oz/day) to include in their diets. Their calorie intake was not monitored or regulated, and thus allowed to float ad libitum.

Walnut intake with caloric regulation: The intervention group participants met with a registered dietitian and received instructions and recipes for inclusion of 392 g of walnuts per week (56 g or 2 oz/day) in their otherwise ad libitum meal plan. Participants received instruction to preserve an isocaloric condition after the addition of walnuts. The study dietitian customized dietary advice to make room calorically for walnuts in the diet, while accommodating the priorities of each study participant. The general approach emphasized a general reduction in portion sizes; participants also received advice, based on baseline dietary intake analysis, of food eliminations that they might want to consider. While the isocaloric condition was encouraged and monitored by the dietitian, participants were provided latitude in determining how to make room for the calories from the walnuts, to better approximate realworld conditions.

Control diet: During the control phase, participants also met with the dietitian and received instructions to follow their usual ad libitum diets, with the exception of avoiding intake of walnuts and specific walnut-containing products. Dietary monitoring and participant weighing was conducted to ensure that an isocaloric condition was maintained.

Outcome measure

To help the study team track any variation in dietary patterns over the course of the study, all participants were asked to provide information on the foods and beverages that they consumed by completing 24-hour recalls using a web-based Automated Self-Administered 24-Hour Recall (ASA24; available from the National Cancer Institute at http://riskfactor.cancer.gov/tools/ instruments/asa24/), which guided them through the process of completing the recall data. Diet quality and the food groups were assessed using the Healthy Eating Index (HEI) 2010. The HEI measures the diet quality, independent of quantity, which is used to assess compliance with the US Dietary Guidelines for Americans and monitor changes in dietary patterns. The HEI is a scoring metric that is used to determine the quality of a given dietary pattern, set of foods, or menu. There are 12 components in the HEI-2010. All of the key Dietary Guidelines food choice recommendations that relate to diet quality are reflected in the HEI-2010's 12 components. Nine of the components focus on adequacy (dietary components to increase) and three focus on moderation (dietary components to decrease). The performance of the HEI-2010 has been evaluated through an assessment of its content validity, construct validity, and reliability.¹³ We analyzed the 12 components of the HEI-2010 to assess the dietary pattern changes in the diets of our study participants.

Statistical analysis

Generalized linear models were used to analyze these data. Regression models were used to adjust for

BMJ Open Diab Res Care: first published as 10.1136/bmjdrc-2016-000293 on 19 October 2016. Downloaded from http://drc.bmj.com/ on April 27, 2024 by guest. Protected by copyright.

potential confounding factors (ie, age, gender, race, compliance, and treatment sequence). All analyses of end points were based on the intention-to-treat principle. Results are expressed as means±SD in text and tables except where otherwise stated. SAS software for Windows V.9.3 (SAS Institute, Cary, North Carolina, USA) was used to carry out all statistical analyses.

The sample size was estimated to provide $\geq 80\%$ power to detect a minimal difference of 7.5 in HEI score between treatment groups, with maximum allowable type I error of 5% (adjusted for three pair-wise comparisons). An SD of 10 in HEI and an attrition rate of 25% were used to compute the sample size.

RESULTS

Study participants

A total of 112 adults at risk for type 2 diabetes enrolled in this study. Most of the participants were female (ie, 72.3%). The proportion of female participants randomized to the calorie-regulated group was comparable to those randomized to the no calorie-regulated group. At baseline, the consumption of vegetables; greens and beans; total fruits (ie, 100% fruit juice); whole fruits (ie, all forms except juice); whole grains; protein foods; seafood and plant protein foods; fatty acids (ie, ratio of polyunsaturated and monounsaturated fatty acids to saturated fatty acids); sodium; refined grain and empty calories (ie, calories from solid fats, alcohol, and added sugars) was comparable in the participants randomized to the calorie-regulated group as compared with those randomized to the no calorie-regulated group. The consumption of dairy products (ie, milk products, such as fluid milk, yogurt, and cheese, and fortified soy beverages) was significantly higher in the calorie-regulated group as compared with the no calorie-regulated group (p=0.04). Baseline data of our study participants are presented in table 1.

Efficacy end point

Calorie-regulated arm: Participants' consumption of seafood and plant protein foods significantly increased from baseline during the walnut phase, as compared with the control $(2.14\pm2.06 \text{ vs } -0.49\pm2.33; \text{ p}=0.003)$. Their intake of sodium non-significantly decreased from baseline during the walnut-included phase, as compared with the walnut-excluded phase $(-3.20\pm4.46 \text{ vs} -0.52)$ ± 5.03 ; p=0.07). Their intake of sodium, empty calories, and dairy products significantly decreased from baseline (ie, -3.20±4.46, p<0.01; -3.15±6.03, p<0.05; and -2.15 ± 3.64 , p<0.05, respectively) with the inclusion of walnuts in the diet. The consumption of refined grains significantly decreased from baseline during the control phase $(-2.04\pm4.39; p<0.05)$. The consumption of fruits and proteins significantly increased from baseline during the walnut-excluded phase (ie, 1.52±2.26, p<0.05 and 0.90 ± 2.07 , p<0.05, respectively). The consumption of vegetables; greens and beans; whole fruit; whole grains; and fatty

	regulated/	regulated/						
Variable	walnuts (N=53)	walnuts (N=49)	p Value					
Gender	69.6%	75.0%	0.53					
Age (years)	56.5±11.7	53.3±11.1	0.15					
2010 Healthy	57.1±17.4	54.0±15.7	0.36					
Eating Index								
score								
Total	3.7±1.5	3.4±1.6	0.23					
vegetables*			0.50					
Greens and	2.3±2.4	2.0±2.3	0.52					
Total fruit+	27+20	27+20	0 02					
Whole fruitt	2.7±2.0 3.0+2.2	2.7 ± 2.0 3.0+2.3	0.92					
Whole grains	4 1+3 9	3.0+3.6	0.33					
Dairy&	6 4+3 1	5 1+3 7	0.17					
Total protein	3.8±1.6	4.3±1.3	0.10					
foods¶								
Seafood and	2.5±2.2	2.4±2.4	0.91					
plant protein								
foods¶,**								
Fatty acids++	4.2±3.8	4.6±3.8	0.60					
Sodium	4.4±3.6	4.0±3.9	0.65					
Refined grains	6.8±3.4	7.3±3.3	0.41					
Empty	13.9±5.5	13.8±6.1	0.95					
calories‡‡	00547 700 0	1000 0 007 5	0.00					
Total calories	2054.7±788.6	1892.8±867.5	0.33					
*Includes any bea	ns and peas not co	unted as total protei	n foods.					
‡Includes all forms	s except juice.							
§Includes all milk	products, such as fl	uid milk, yogurt, and	d cheese,					
and fortified soy be	everages.	and not with vocate						
when the total pro	tein foods standard	is otherwise not me	et.					
**Includes seafood	d, nuts, seeds, soy p	products (other than						
beverages) as wel	I as beans and pea	s counted as total p	rotein					
ttRatio of polyung	toods.							
saturated fatty acid	ds.	aneatoratou raity do						
‡‡Calories from so	olid fats, alcohol, an	d added sugars; thr	reshold					
for counting alcoho	ol is >13 g/1000 kca	u.						

cids (ie. ratio of polyupsaturated and monoupsaturated

acids (ie, ratio of polyunsaturated and monounsaturated fatty acids to saturated fatty acids) did not change significantly (p>0.05) from baseline during the walnut-included phase, compared with the walnut-excluded phase.

No calorie-regulated arm: The consumption of seafood and plant protein foods and healthful fatty acids significantly increased from baseline during the walnutincluded phase as compared with the walnut-excluded phase (1.67±2.32 vs -1.21±3.93; p=0.002 and 1.43±4.53 vs -1.76 ± 4.80 ; p=0.02, respectively). The consumption of vegetables significantly decreased from baseline during walnut-included phase compared with the the walnut-excluded phase (-0.55±1.84 vs 0.80±2.21; p=0.04). Empty calorie consumption non-significantly decreased from baseline during the walnut-included phase as compared with the walnut-excluded phase $(-3.03\pm8.24 \text{ vs})$ 0.70±8.64; p=0.06). However, empty calorie consumption

Cardiovascular and metabolic risk

decreased significantly from baseline in the walnutincluded phase (-3.03 ± 8.24 ; p<0.05). The consumption of greens and beans; fruit; whole fruit; whole grains; dairy products; protein foods; sodium; and refined grains from baseline during the walnut-included phase was comparable to that found during the walnut-excluded phase (p>0.05).

Calorie-regulated arm versus no calorie-regulated arm: The decrease in the consumption of dairy products from baseline observed in participants in the calorie-regulated arm during the walnut-included phase was significant as compared with those in the no calorie-regulated arm $(-2.15\pm3.64 \text{ vs } 1.06\pm4.42; \text{ p=0.02})$. No differential treatment effects were observed (p>0.05) in the consumption of vegetables; greens and beans; fruit; whole fruit; whole grains; protein foods; sodium; refined grains; or empty calories from baseline with the inclusion of walnuts in the diets of participants in the calorie-regulated arm as compared with those in the no calorie-regulated arm.

Our efficacy data are reported in table 2.

DISCUSSION

Our data suggest that the consumption of seafood and plant protein foods and healthful fatty acids significantly increased, while the consumption of sodium, empty calories, vegetables, and dairy foods significantly decreased, with the inclusion of walnuts in the diets of these adults at risk for type 2 diabetes. The consumption of greens and beans, whole fruit, and whole grains did not change significantly with the inclusion of walnuts in the diet.

There was a significant reduction in dairy product consumption in the participants randomized in the calorie-regulated arm as compared with those in the no calorie-regulated arm. All of these dietary changes observed during the walnut-included phase led to an overall significant improvement in diet quality among this group of adults at risk for type 2 diabetes.

Studies have shown that improved diet quality is associated with lower risk for type 2 diabetes.¹⁰¹¹ Further, in a study by Dominguez *et al*,¹⁴ dietary-based diabetes risk score, which integrates optimal food patterns, with the risk of developing type 2 diabetes was inversely associated with the development of type 2 diabetes. The American Diabetes Association (ADA) recommends intake of plant-based protein sources, for example, beans, nuts, seeds, or tofu; fish and seafood; chicken and other poultry; eggs; and low-fat dairy products. The ADA also recommends intake of natural sources of vegetable fats, such as nuts, seeds, or avocados; foods that provide omega-3 fatty acids, such as salmon, tuna, mackerel, or walnuts; and plant-based oils, such as canola, grapeseed, or olive oils as the healthful choices for blood glucose control in diabetes and prediabetes.

We observed a significant increase in the consumption of seafood and plant protein foods with the inclusion of walnuts in the diets of adults at risk for diabetes. In a study by Patel *et al*,¹⁵ consumption of lean fish, total fish and shellfish was not associated with the risk of type 2 diabetes. However, fatty fish consumption was inversely associated with the risk of type 2 diabetes. Wu *et al*¹⁶ also demonstrated no association between the consumption of fish/seafood and the risk of type 2 diabetes, and

Variable	CRW	Calorie regulated/ walnut excluded	p Value	NCRW	No calorie regulated/walnut excluded	p Value	CRW vs NCRW p Value
Healthy Eating Index score	9.14±17.71**	0.40±15.13	0.02	7.02±15.89 [¥]	-5.92±21.84 [¥]	0.001	0.60
Total vegetables	-0.35±1.93	-0.30±1.8	0.93	-0.55±1.84	0.80±2.21	0.04	0.76
Greens and beans	-0.60±2.83	-0.31±2.74	0.76	-0.69±2.71	-0.14±3.72	0.59	0.93
Total fruit	-0.11±2.39	1.52±2.26 [¥]	0.06	-0.55±2.91	-0.13±3.30	0.64	0.62
Whole fruit	-0.36±2.52	1.23±2.40	0.10	-0.51±3.40	-0.34±3.85	0.87	0.88
Whole grains	-1.01±4.64	-0.11±4.95	0.55	0.82±4.87	-0.35±4.56	0.46	0.24
Dairy	–2.15±3.64 [¥]	-0.51±4.39	0.20	1.06±4.42	-0.03±3.57	0.42	0.02
Total protein foods	0.72±1.46	0.90±2.07 [¥]	0.74	0.40±1.00	0.14±2.04	0.64	0.57
Seafood and plant protein foods	2.14±2.06**	-0.49±2.33	0.003	1.67±2.32	-1.21±3.93*	0.002	0.60
Fatty acids	0.93±6.51	0.02±4.26	0.51	1.43±4.53	-1.76±4.80	0.02	0.72
Sodium	-3.20±4.46*	-0.52±5.03	0.07	-1.77±6.27	-1.79±5.71	0.99	0.32
Refined grains	-1.13±3.18	-2.04±4.39 [¥]	0.42	-0.98±3.88	-0.47±5.02	0.64	0.90
Empty calories	-3.15 ± 6.03^{4}	-2.22±5.39	0.63	-3.03±8.24 [¥]	0.70±8.64	0.06	0.95
Total calories	97.0±845.0	-213.4±869.0	0.08	169.8±613.2	-59.6±914.8	0.40	0.70

*<0.05, *<0.01, **<0.001 indicates significant changes from baseline.

CRW, calorie regulated/walnut included; NCRW, no calorie regulated/walnut included.

observed an inverse relationship between the consumption of α -linolenic acid and the risk of diabetes. However, epidemiological studies have shown an association between the consumption of animal proteins and the increased risk of type 2 diabetes.^{17–19} In a meta-analysis by Viguiliouk *et al*²⁰ replacing animal protein sources with plant protein food sources improved glycemic control in individual with diabetes. Seafood and plant-based proteins have a relatively low glycemic index. Foods with a low glycemic index are typically recommended for patients with or at risk for diabetes. Increased consumption of seafood and plant protein foods in the diets of patient at risk for diabetes or diabetes could help in blood glucose control. On the other hand, animal proteins, particularly red meats, are rich in saturated fatty acids that could adversely affect blood glucose levels.

We observed a decrease in the consumption of dairy products with the inclusion of walnuts in the diet. Chen et al^{21} demonstrated that increased vogurt consumption was associated with low risk of type 2 diabetes, while the consumption of other dairy foods or total dairy products was not associated with the incidence of type 2 diabetes. Malik *et al*²² demonstrated that high dairy product consumption during adolescence was associated with lower risk of type 2 diabetes. Meta-analyses²³²⁴ have shown that modest increases in consumption of dairy products such as low fat dairy, cheese and yogurt have been associated with reduced incidence of type 2 diabetes. On the contrary, in a large prospective study by Sluijs *et al*, 25total dairy product consumption was not associated with the risk of type 2 diabetes. However, an inverse relationship was observed between cheese intake and combined fermented dairy product consumption and the risk of type 2 diabetes.

We observed a non-significant decrease in empty calorie consumption from baseline during the walnutincluded phase as compared with the walnut-excluded phase. However, the consumption of empty calories decreased significantly from baseline in the walnutincluded phase. Pan *et al*²⁶ demonstrated that substitution of plain water consumption for sugar sweetened beverages or fruit juice was associated with a lower risk for type 2 diabetes. Fagherazzi *et al*²⁷ demonstrated that the consumption of sugar sweetened and artificially sweetened beverages was associated with increased incidence of type 2 diabetes. These empty calorie foods typically have a relatively high glycemic index. Foods with a high glycemic index are typically not recommended in diabetes and prediabetes due to the risk of adversely affecting blood glucose levels.

In our study, the consumption of whole grains did not differ significantly during the walnut-included phase as compared with the walnut-excluded phase. However, the consumption of refined grains decreased from baseline during the walnut-excluded phase. O'Neil *et al*²⁸ demonstrated that tree nut consumption was associated with improved diet quality and improved nutrient intake.

A meta-analysis by Blanco Mejia *et al*²⁹ demonstrated that tree nuts improved triglycerides and blood glucose levels in individuals with metabolic syndrome. Another meta-analysis by Ye *et al*^{30} also demonstrated that whole grain consumption was associated with vascular disease prevention. In another study by Sahyoun *et al*,³¹ high whole grain consumption was inversely associated with metabolic syndrome and cardiovascular mortality. Aune et al^{32} also showed that high whole grain consumption was associated with reduced risk of type 2 diabetes. Sun et al^{β^3} demonstrated that substituting brown rice for white rice reduced the risk for type 2 diabetes. Whole grains, as compared with refined grains, have a low glycemic index. The decrease in the consumption of refined grains in the walnut-excluded phase in our study may be due to the Hawthorn effect.

Our study showed a significant increase in the consumption of fatty acids with the inclusion of walnuts in the diet, as compared with their exclusion, among participants who did not receive counseling to regulate caloric intake. In a study by Guasch-Ferré et al,³⁴ higher olive oil consumption was associated with a lower risk of type 2 diabetes. Increased animal fat consumption has been associated with an increased risk of type 2 diabetes.³⁵ Diets rich in monounsaturated fatty acids have been shown to decrease fasting insulin and improve the homeostasis model assessment of insulin resistance and the low-density lipoprotein to high-density lipoprotein ratio.³⁶ Salmerón et al⁸⁷ demonstrated that intakes of total fats and saturated and monounsaturated fatty acids are not associated with risk of type 2 diabetes in women, but that trans fatty acids increase the risk, while polyunsaturated fatty acids reduce the risk. Fatty acids affect glucose metabolism by modifying cell membrane function, enzyme activity, insulin signaling, and gene expression.³⁸

We did not observe any difference between the inclusion versus the exclusion of walnuts in the consumption of greens and beans; fruit; and whole fruit. In a study by Cooper *et al*,³⁹ consumption of fruit or vegetables was not associated with the risk of diabetes, while the consumption of green leafy vegetables was inversely associated with the risk of diabetes. Bazzano et al⁴⁰ also demonstrated that the consumption of green leafy vegetables and fruit was associated with a lower risk of diabetes, whereas consumption of fruit juices may be associated with an increased risk among women. In a meta-analysis by Carter *et al*,⁴¹ increased daily intake of green leafy vegetables was associated with a significantly reduced risk of type 2 diabetes. In another meta-analysis by Li *et al*,⁴² the consumption of fruit or green leafy vegetables was associated with a reduced risk of type 2 diabetes. A study by Wu *et al*⁴³ showed that 2–3 servings per day of vegetables and 2 servings per day of fruits reduced the risk of type 2 diabetes when compared with other levels of vegetable and fruit consumption, respectively. Green leafy vegetables are relatively low in saturated and *trans* fatty acids, and have a low glycemic index.

In our study, while the consumption of sodium decreased non-significantly from baseline in the walnutincluded phase compared with the walnut-excluded phase, we observed a significant decrease in the consumption of sodium from baseline with the inclusion of walnuts among our participants in the calorie-regulated arm. In a study by Horikawa et al.⁴⁴ high sodium consumption was associated with increased risk of cardiovascular disease (CVD) in individual with type 2 diabetes. A synergistic effect of the risk of CVD was also identified when high sodium consumption was coupled with poor glucose control. A review by Poggio *et al*⁴⁵ also concluded that high sodium consumption was associated with higher CVD mortality. However, other studies have shown no association between sodium consumption and CVD.^{46 47} High sodium consumption affects the cardiovascular system by adversely affecting blood pressure. The reduction of consumption of sodium with the inclusion of walnuts could therefore reduce our participants risk for CVD.

Limitations

This study had several limitations. First, the participants were predominantly Caucasians and female, which limit us in extrapolating our findings to the general population. Second, the study had a small sample size. This limitation was overcome to some extent by crossing over the participants to the two different treatment assignments, thereby improving the power of the study. Third, the study relied on self-report by the participants for their dietary records, which can introduce measurement and recall biases. Fourth, the participants were not monitored on a daily basis and were not administered a supervised diet. However, this can also be viewed as a strength of the study because it provides a more realistic scenario and potentially increases external validity. In addition, the variation of the dietary patterns among participants may make it difficult to interpret the findings. Fifth, the advice given by the dietitian to the calorie-regulated group may explain some of the dietary changes observed. However, generally similar dietary changes were observed among those who did not receive advice to regulate their calorie intake. Sixth, another limitation is that intake of dairy products was significantly higher in the calorie-regulated arm as compared with the group not receiving guidance for total energy intake. However, we compared the pre/post scores of the calorie-regulated group to the pre/post scores of the group without this guidance to control for baseline differences.

CONCLUSIONS

The inclusion of walnuts in the diets in these adults at risk for type 2 diabetes led to the inclusion in their diets of healthful foods typically recommended for blood glucose control. We observed a significant increase in intake of seafood and plant-based proteins and healthful fatty acids, while the consumption of sodium, empty calorie, vegetables, and dairy foods decreased significantly with the inclusion of walnuts in the diets of these adults. The inclusion of walnuts in the diets of adults with prediabetes facilitated the introduction of healthful foods in their diets that are suitable for blood glucose control. Our study suggests that adding walnuts produces changes in intake of other foods. Future studies should routinely examine the impact of single food interventions on overall dietary patterns, since the addition of any given food almost invariably means changes in the intake of others.

Acknowledgements The authors wish to acknowledge the technical assistance of Susan Acheychek RDMS for assessing the clinical outcomes.

Contributors VYN was involved in study design, project oversight, data analysis, data interpretation, developed manuscript draft, final approval. NY assisted with development of manuscript draft. RGA was the project coordinator. PP conducted statistical analysis, data interpretation. JAT was the study dietitian and provided critical review. DLK was involved in study design, project oversight, data interpretation, critical review of paper, final approval. All authors have reviewed and approve of the completed revised manuscript.

Funding Funding for this study has been provided by the California Walnut Commission.

Competing interests This study was conducted with funding from the Centers for Disease Control and Prevention and the California Walnut Commission. DLK has been compensated for public speaking by the California Walnut Commission.

Ethics approval Griffin Hospital Institutional Review Board.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: http:// creativecommons.org/licenses/by-nc/4.0/

REFERENCES

- 1. http://www.who.int/mediacentre/news/releases/2016/world-healthday/en/
- Imamura F, O'Connor L, Ye Z, *et al.* Consumption of sugar sweetened beverages, artificially sweetened beverages, and fruit juice and incidence of type 2 diabetes: systematic review, meta-analysis, and estimation of population attributable fraction. *Br J Sports Med* 2016;50:496–504.
- Centers for Disease Control and Prevention. National diabetes statistics report: estimates of diabetes and its burden in the United States, 2014. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, 2014. http:// www.niddk.nih.gov/health-information/health-communicationprograms/ndep/am-i-at-risk/diabetes preventable/Pages/ diabetesispreventable.aspx
- Knowler WC, Barrett-Connor E, Fowler SE, et al., Diabetes Prevention Program Research Group. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. N Engl J Med 2002;346:393–403.
- 5. http://www.cdc.gov/diabetes/prevention/index.html
- Knowler WC, Fowler SE, Hamman RF, *et al.*, Diabetes Prevention Program Research Group. 10-year follow-up of diabetes incidence and weight loss in the Diabetes Prevention Program Outcomes Study. *Lancet* 2009;374:1677–86.
- 7. Diabetes Prevention Program Research Group. Long-term effects of lifestyle intervention or metformin on diabetes development and microvascular complications over 15-year follow-up: the Diabetes

Cardiovascular and metabolic risk

Prevention Program Outcomes Study. *Lancet Diabetes Endocrinol* 2015;3:866–75.

- Alouki K, Delisle H, Bermúdez-Tamayo C, *et al.* Lifestyle interventions to prevent type 2 diabetes: a systematic review of economic evaluation studies. *J Diabetes Res* 2016;2016:2159890.
- http://www.diabetes.org/advocacy/advocacy-priorities/prevention/ preventing-diabetes-nutrition.html
- 10. de Koning L, Chiuve SE, Fung TT, *et al.* Diet-quality scores and the risk of type 2 diabetes in men. *Diabetes Care* 2011;34:1150–6.
- Jacobs S, Harmon BE, Boushey CJ, *et al.* A priori-defined diet quality indexes and risk of type 2 diabetes: the Multiethnic Cohort. *Diabetologia* 2015;58:98–112.
- Njike VY, Ayettey R, Petraro P, et al. Walnut ingestion in adults at risk for diabetes: effects on body composition, diet quality, and cardiac risk measures. BMJ Open Diabetes Res Care 2015;3: e000115.
- Guenther PM, Kirkpatrick SI, Reedy J, *et al.* The Healthy Eating Index-2010 is a valid measure of diet quality according to the 2010 Dietary Guidelines for Americans. *J Nutr* 2014;144:399–407.
- Dominguez LJ, Bes-Rastrollo M, Basterra-Gortari FJ, *et al.* Association of a dietary score with incident type 2 diabetes: the Dietary-Based Diabetes-Risk Score (DDS). *PLoS ONE* 2015;10: e0141760.
- Patel PS, Forouhi NG, Kuijsten A, *et al.*, InterAct Consortium. The prospective association between total and type of fish intake and type 2 diabetes in 8 European countries: EPIC-InterAct Study. *Am J Clin Nutr* 2012;95:1445–53.
- Wu JH, Micha R, Imamura F, et al. Omega-3 fatty acids and incident type 2 diabetes: a systematic review and meta-analysis. Br J Nutr 2012;107(Suppl 2):S214–27.
- Pan A, Sun Q, Bernstein AM, *et al.* Changes in red meat consumption and subsequent risk of type 2 diabetes mellitus: three cohorts of US men and women. *JAMA Intern Med* 2013;173:1328–35.
- de Koning L, Fung TT, Liao X, et al. Low-carbohydrate diet scores and risk of type 2 diabetes in men. Am J Clin Nutr 2011;93:844–50.
- Malik VS, Li Y, Tobias DK, *et al.* Dietary protein intake and risk of type 2 diabetes in us men and women. *Am J Epidemiol* 2016;183:715–28.
- Viguiliouk E, Stewart SE, Jayalath VH, et al. Effect of replacing animal protein with plant protein on glycemic control in diabetes: a systematic review and meta-analysis of randomized controlled trials. Nutrients 2015;7:9804–24.
- Chen M, Sun Q, Giovannucci E, et al. Dairy consumption and risk of type 2 diabetes: 3 cohorts of US adults and an updated meta-analysis. *BMC Med* 2014;12:215.
- 22. Malik VS, Sun Q, van Dam RM, *et al.* Adolescent dairy product consumption and risk of type 2 diabetes in middle-aged women. *Am J Clin Nutr* 2011;94:854–61.
- Gao D, Ning N, Wang C, *et al.* Dairy products consumption and risk of type 2 diabetes: systematic review and dose-response meta-analysis. *PLoS ONE* 2013;8:e73965.
- Tong X, Dong JY, Wu ZW, *et al.* Dairy consumption and risk of type 2 diabetes mellitus: a meta-analysis of cohort studies. *Eur J Clin Nutr* 2011;65:1027–31.
- Sluijs I, Forouhi NG, Beulens JW, *et al.*, InterAct Consortium. The amount and type of dairy product intake and incident type 2 diabetes: results from the EPIC-InterAct Study. *Am J Clin Nutr* 2012;96:382–90.
- Pan A, Malik VS, Schulze MB, *et al.* Plain-water intake and risk of type 2 diabetes in young and middle-aged women. *Am J Clin Nutr* 2012;95:1454–60.
- Fagherazzi G, Vilier A, Saes Sartorelli D, *et al.* Consumption of artificially and sugar-sweetened beverages and incident type 2 diabetes in the Etude Epidemiologique aupres des femmes de la Mutuelle Generale de l'Education Nationale-European Prospective Investigation into Cancer and Nutrition cohort. *Am J Clin Nutr* 2013;97:517–23.

- O'Neil CE, Keast DR, Fulgoni VL III, *et al.* Tree nut consumption improves nutrient intake and diet quality in US adults: an analysis of National Health and Nutrition Examination Survey (NHANES) 1999– 2004. *Asia Pac J Clin Nutr* 2010;19:142–50.
- Blanco Mejia S, Kendall CW, Viguiliouk E, *et al.* Effect of tree nuts on metabolic syndrome criteria: a systematic review and meta-analysis of randomised controlled trials. *BMJ Open* 2014;4: e004660.
- Ye EQ, Chacko SA, Chou EL, *et al.* Greater whole-grain intake is associated with lower risk of type 2 diabetes, cardiovascular disease, and weight gain. *J Nutr* 2012;142:1304–13.
- Sahyoun NR, Jacques PF, Zhang XL, *et al.* Whole-grain intake is inversely associated with the metabolic syndrome and mortality in older adults. *Am J Clin Nutr* 2006;83:124–31.
- Aune D, Norat T, Romundstad P, *et al.* Whole grain and refined grain consumption and the risk of type 2 diabetes: a systematic review and dose-response meta-analysis of cohort studies. *Eur J Epidemiol* 2013;28:845–58.
- Sun Q, Spiegelman D, van Dam RM, *et al.* White rice, brown rice, and risk of type 2 diabetes in US men and women. *Arch Intern Med* 2010;170:961–9.
- Guasch-Ferré M, Hruby A, Salas-Salvadó J, *et al.* Olive oil consumption and risk of type 2 diabetes in US women. *Am J Clin Nutr* 2015;102:479–86.
- Thanopoulou AC, Karamanos BG, Angelico FV, *et al.* Dietary fat intake as risk factor for the development of diabetes: multinational, multicenter study of the Mediterranean Group for the Study of Diabetes (MGSD). *Diabetes Care* 2003;26:302–7.
- Due A, Larsen TM, Mu H, *et al.* Comparison of 3 ad libitum diets for weight-loss maintenance, risk of cardiovascular disease, and diabetes: a 6-mo randomized, controlled trial. *Am J Clin Nutr* 2008;88:1232–41.
- Salmerón J, Hu FB, Manson JE, *et al.* Dietary fat intake and risk of type 2 diabetes in women. *Am J Clin Nutr* 2001;73:1019–26.
- 38. Řisérus U, Willett WC, Hu FB. Dietary fats and prevention of type 2 diabetes. *Prog Lipid Res* 2009;48:44–51.
- Cooper AJ, Forouhi NG, Ye Z, *et al.*, InterAct Consortium. Fruit and vegetable intake and type 2 diabetes: EPIC-InterAct prospective study and meta-analysis. *Eur J Clin Nutr* 2012;66:1082–92.
- Bazzano LA, Li TY, Joshipura KJ, et al. Intake of fruit, vegetables, and fruit juices and risk of diabetes in women. *Diabetes Care* 2008;31:1311–17.
- Carter P, Gray LJ, Troughton J, *et al.* Fruit and vegetable intake and incidence of type 2 diabetes mellitus: systematic review and meta-analysis. *BMJ* 2010;341:c4229.
- Li M, Fan Y, Zhang X, *et al.* Fruit and vegetable intake and risk of type 2 diabetes mellitus: meta-analysis of prospective cohort studies. *BMJ Open* 2014;4:e005497.
- Wu Y, Zhang D, Jiang X, *et al.* Fruit and vegetable consumption and risk of type 2 diabetes mellitus: a dose-response meta-analysis of prospective cohort studies. *Nutr Metab Cardiovasc Dis* 2015;25:140–7.
- 44. Horikawa C, Yoshimura Y, Kamada C, et al. Japan Diabetes Complications Study Group Dietary sodium intake and incidence of diabetes complications in Japanese patients with type 2 diabetes: analysis of the Japan Diabetes Complications Study (JDCS). J Clin Endocrinol Metab 2014;99:3635–43.
- Poggio R, Gutierrez L, Matta MG, *et al.* Daily sodium consumption and CVD mortality in the general population: systematic review and meta-analysis of prospective studies. *Public Health Nutr* 2015;18:695–704.
- Kalogeropoulos AP, Georgiopoulou VV, Murphy RA, et al. Dietary sodium content, mortality, and risk for cardiovascular events in older adults: the Health, Aging, and Body Composition (Health ABC) Study. JAMA Intern Med 2015;175:410–19.
- Cohen HW, Hailpern SM, Alderman MH. Sodium intake and mortality follow-up in the Third National Health and Nutrition Examination Survey (NHANES III). J Gen Intern Med 2008;23:1297–302.