

Determinants of physical activity among 6856 individuals with diabetes: a nationwide cross-sectional study

Sofie Rath Mortensen ^{1,2}, Peter Lund Kristensen,¹ Anders Grøntved,¹ Mathias Ried-Larsen,³ Cathrine Lau,⁴ Søren T Skou^{1,2}

To cite: Mortensen SR, Kristensen PL, Grøntved A, *et al.* Determinants of physical activity among 6856 individuals with diabetes: a nationwide cross-sectional study. *BMJ Open Diab Res Care* 2022;**10**:e002935. doi:10.1136/bmjdr-2022-002935

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjdr-2022-002935>).

Received 12 May 2022
Accepted 15 July 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Department of Sports Science and Clinical Biomechanics, University of Southern Denmark, Odense, Denmark

²Department of Physiotherapy and Occupational Therapy, Næstved-Slagelse-Ringsted Hospitals, Slagelse, Denmark

³Centre for Physical Activity Research, Copenhagen University Hospital, Copenhagen, Denmark

⁴Centre for Clinical Research and Prevention, Bispebjerg and Frederiksberg Hospital, Frederiksberg, Denmark

Correspondence to

Sofie Rath Mortensen; smortensen@health.sdu.dk

ABSTRACT

Introduction The aims of this cross-sectional study were to (1) describe habitual physical activity and adherence to WHO recommendations, and (2) investigate the association of comorbidity, obesity, stress, and health-related quality of life (HRQoL) with moderate to vigorous physical activity (MVPA) among individuals with diabetes.

Research design and methods This study included 6856 participants with diabetes from the Danish National Health Survey from 2017. The primary outcome measure was weekly MVPA. Exposures included self-reported number of conditions, body mass index (BMI), perceived stress, and HRQoL. Mean difference in MVPA across exposures was estimated by multiple linear regression analyses.

Results Forty per cent of individuals with diabetes were not adherent to WHO recommendations for physical activity. Individuals with diabetes had higher BMI, more comorbidities, higher perceived stress, and lower HRQoL. Individuals with three or more comorbidities were significantly associated with lower weekly MVPA (−0.48 hours/week, 95% CI −0.88 to −0.07) compared with individuals with no comorbidity. Furthermore, overweight or obese (class I–III) individuals engaged in significantly less weekly MVPA (obese class III vs normal weight: −1.98 hours/week, 95% CI −2.49 to −1.47). Higher perceived stress was significantly associated with lower weekly MVPA (−1.76 hours/week, 95% CI −2.18 to −1.34) versus low perceived stress. Finally, having low physical and mental HRQoL was associated with lower weekly MVPA (−0.93 hours/week, 95% CI −1.19 to −0.66 and −0.39 hours/week, 95% CI −0.71 to −0.08 respectively vs moderate or high HRQoL).

Conclusions We found that 40% of individuals with diabetes do not engage regularly in adequate physical activity. Comorbidities, higher BMI, higher perceived stress, and lower HRQoL were associated with less engagement in physical activity. This study suggests that subgroups of individuals with diabetes are at higher risk of physical inactivity.

INTRODUCTION

Physical inactivity alone is estimated to cause 7% of the burden of type 2 diabetes.¹ On the other hand, engaging in regular physical activity is a cornerstone of diabetes management to prevent long-term diabetes complications, declines in quality of life and premature

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ No previous large-scale study has provided a detailed description of habitual physical activity among individuals with diabetes and associated factors.

WHAT THIS STUDY ADDS

⇒ Forty per cent of individuals with diabetes do not engage regularly in adequate physical activity.
⇒ Having comorbidities, higher body mass index, higher perceived stress, and lower health-related quality of life is significantly associated with lower moderate to vigorous physical activity among individuals with diabetes.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ Subgroups of individuals with diabetes are at higher risk of physical inactivity and might need additional support or tailored interventions to increase their physical activity level.

mortality.^{2,3} According to the WHO and the American Diabetes Association,^{4,5} adults living with diabetes are recommended to do at least 150–300 min of moderate-intensity or 75–150 min of vigorous-intensity (or an equivalent combination) aerobic physical activity weekly.^{4,5} Living with diabetes can be very stressful and significantly affect the individual's quality of life.^{6,7} Individuals with diabetes may worry about existing and future complications and comorbidities, as well as feeling guilty and ashamed of not adhering to lifestyle recommendations, for example, in terms of body mass index (BMI) and physical activity.⁸ Meeting the WHO recommendations of regular physical activity is a major challenge among individuals with diabetes,⁹ but some studies report that about 40%–45% of individuals with diabetes do adhere to the recommendations^{10,11} and their treatment should thus not necessarily revolve around increasing physical activity. Morbidity, obesity, stress, and lower health-related quality of life

(HRQoL) are associated with decreased habitual physical activity in the general population.^{12–15} Given that individuals with diabetes are at high risk of suffering from all these factors,^{6 16 17} it is likely that physical activity among individuals with diabetes is also associated with these factors. Availability of such information would be an important resource when designing and promoting a physical activity intervention taking individual characteristics, needs and preferences into account, and thereby spending the resources on those in most need. However, no previous large-scale studies have provided a detailed description of habitual physical activity among individuals with diabetes and associated factors.

Therefore, the aims of this study were to (1) describe habitual physical activity and adherence to WHO recommendations, and (2) investigate the association of comorbidity, obesity, stress, and HRQoL with moderate to vigorous physical activity (MVPA) among individuals with diabetes based on data from a large nationwide Danish survey.

RESEARCH DESIGN AND METHODS

Reporting of this cross-sectional study followed the ‘Strengthening the Reporting of Observational Studies in Epidemiology’ checklist.¹⁸

Setting and data sources

We used data from the cross-sectional nationwide Danish National Health Survey (DNHS) from 2017. The overall aim of the DNHS is to monitor the status and trends in physical and mental health in the adult Danish population (≥ 16 years). A detailed description of the DNHS design is available on the DNHS web page (<https://www.danskernessundhed.dk/>).^{19 20} In 2017, a total of 312 349 individuals were randomly drawn from the population using the Danish Civil Health Registration System and invited via a secure electronic mail service (Digital Post) or regular postal service to participate in the survey. The questionnaire was fully or partially completed (sex, age, and at least one other question answered) by 183 372 respondents (58.7%).²⁰

Participants

The present study included participants with ‘Diabetes’ and ‘No known diabetes’. Following the categorization of diabetes from DNHS, respondents were defined as ‘Having diabetes’ if they had answered ‘I have diabetes now’ or ‘I have had diabetes’ and ‘I suffer from sequela due to the diabetes’. The question on diabetes did not distinguish between types of diabetes; therefore, the respondents covered the whole spectrum of diabetes. Out of 183 372 respondents, 10 216 individuals reported to have diabetes. Respondents with complete data on outcome, exposure, and confounder variables were considered eligible for the present study.

Outcome variables

The outcomes of interest were self-reported MVPA (hours/week), adherence to WHO recommendations of

physical activity and sedentary behavior, and total sedentary behavior (hours/day). MVPA was assessed with the question: ‘During a regular week, how much time do you spend on moderate and vigorous physical activities, where you can feel your pulse and your breathing increase (eg, walking, cycling as transport or recreational activity, hard gardening, running or exercise sports)? Indicate only activities lasting at least 10 min’. Respondents replied in hours and minutes.

Adherence to WHO recommendations was assessed with the question regarding MVPA and the question: ‘How much time of the above-mentioned physical activities do you spend in a regular week on vigorous physical activities? These are activities where your pulse is substantially increased, you sweat, and which cause you to be out of breath and to find it hard to talk (eg, swimming, running, cycling at high speed, strength training or ball games)’. Adherence to WHO recommendations was dichotomized to ‘Following WHO recommendations’ or ‘Not following WHO recommendations’. Respondents were categorized as ‘Following WHO recommendations’ if they reported at least 150 min/week of moderate intensity, at least 75 min/week of vigorous intensity, or an equivalent combination.⁴

In addition, self-reported MVPA was categorized into four levels of habitual physical activity in accordance with the WHO recommendations: (1) inactive: 0 hour/week; (2) inadequate: participants do not meet recommendations (150–300 min/week of moderate intensity or 75–150 min/week of vigorous intensity); (3) adequate: participants meet recommendations, but not more than 300 min/week of moderate intensity or 150 min/week of vigorous intensity; and (4) optimal: participants with more than 300 min/week of moderate intensity or 150 min/week of vigorous intensity.

Total sedentary behavior was assessed with the question: ‘On a typical weekday/workday, how much time do you spend on sitting down in each of the following situations? Please consider your total sitting time and distribute it in each of the following categories’. Respondents reported minutes and hours spent on (1) transport (eg, in car, bus or train; not cycling); (2) work/school/education (eg, sitting by the desk or at meeting); (3) leisure time: by screen (eg, television, computer, tablet, smartphone); (4) leisure time: other (eg, meals, reading, social gatherings). All questions regarding MVPA and sedentary behavior have been validated.²¹ Motivation for being more physically active was assessed with the question: ‘Do you want to be more physically active?’.

Exposures

Comorbidity

Self-reported information on selected long-term conditions (excluding diabetes) and sequela were used to assess comorbidity. Respondents reported whether they had or have had selected long-term conditions, and whether they were suffering from sequelae due to the specific long-term condition. The definition of comorbidity was based on diagnoses organized in 10 groups of different body systems according to Willadsen *et al.*²² In

this study, there were only seven groups due to lack of information on conditions in all body systems, and since diabetes was the only endocrine condition in the survey, it was excluded from the comorbidity variable. The seven groups were: (1) lung, (2) musculoskeletal, (3) mental, (4) cancer, (5) neurological, (6) cardiovascular, and (7) sensory organs.

The variable was categorized as a count variable, counting the numbers of comorbidities from zero to three or more from different body systems including diabetes. Suffering from several long-term conditions within the same body system, for example, hypertension and myocardial infarction, would still only count as one comorbidity.

Individuals with mental comorbidities engage less regularly in physical activity compared with individuals with only somatic comorbidities.^{23 24} Therefore, a variable differentiating between comorbidities with and without a mental health condition was created.

Obesity

Self-reported data on height and weight were used to calculate BMI (kg/m^2). BMI was categorized into five groups: underweight/normal weight (BMI <25.0), overweight (BMI ≥ 25.0 to <30.0), obese class I (BMI ≥ 30.0 to <35.0), obese class II (BMI ≥ 35.0 to <40.0), and obese class III (BMI ≥ 40), as defined by the WHO.²⁵

Stress

Self-reported psychological stress was assessed using the Danish version of Cohen's 10-item Perceived Stress Scale (PSS).^{26 27} PSS was categorized into three as follows: (1) low perceived stress (scores ranging from 0 to 13), (2) moderate perceived stress (scores ranging from 14 to 26), and (3) high perceived stress (scores ranging from 27 to 40).²⁷

Health-related quality of life

Self-reported HRQoL (physical and mental health) was assessed using the 12-item Short-Form Health Survey and was categorized into a physical and mental scale.^{28 29} A physical score of 50 or less was categorized as 'Low physical HRQoL', and a mental score of 42 or less was categorized as 'Low mental HRQoL'.³⁰

Covariates

The following covariates are suggested to be potential independent risk factors of the four exposures (comorbidity, obesity, stress, and HRQoL) and the outcome (physical activity): age, sex, ethnicity, marital status, educational level, alcohol consumption, smoking, and diet. Directed acyclic graphs³¹ of the assumed causal relations between exposures and outcome of the primary analysis are available in online supplemental file 1. Marital status was dichotomized into 'Married or living with partner' or 'Living alone'. Educational level was assessed using the respondents' highest level of education and categorized into three: (1) primary and lower secondary education, (2) upper secondary or vocational,

and (3) higher education. Ethnicity was categorized into three groups: Danish, other Western, and non-Western background. Smoking was categorized into (1) smoker, (2) ex-smoker, (3) never smoked. Alcohol consumption was measured by the number of drinks of beer, wine and spirits consumed in a typical week and categorized in accordance with the national recommendations from the Danish Health Authority: (1) no alcohol (0 drink), (2) below low risk (men >0 and <14 drinks, women >0 and <7 drinks), (3) above low risk (men ≥ 14 and ≤ 21 drinks, women ≥ 7 and ≤ 14 drinks), (4) high risk (men >21 drinks, women >14 drinks). Eating habits were measured using the diet score dividing the respondents into three health levels based on their diet: (1) healthy diet, (2) medium healthy diet, and (3) unhealthy diet.³²

Statistical analyses

A statistical analysis plan was developed (online supplemental file 2) and openly available (<https://osf.io/25u4g/>) prior to commencing the analyses. Cross-tabulations were conducted to describe habitual physical activity among individuals with diabetes and individuals with no known diabetes and to display potential subgroup differences.

Descriptive statistics are presented as numbers and percentages or means and SDs or medians and IQR. Four multiple linear regression analyses were conducted to investigate the associations between the exposures (comorbidity, obesity, stress, and HRQoL) and the outcome MVPA. Results of the multiple linear regression analyses are presented as mean differences with 95% CIs across levels of exposures with no comorbidity, underweight or normal weight, low perceived stress, and moderate to high HRQoL as reference categories.

To explore the associations between the same four exposures and adherence to WHO recommendations of physical activity, four multiple logistic regression analyses were performed. Results of the logistic regression analyses are presented as prevalence ORs with 95% CI. Furthermore, four multiple linear regression analyses were performed with total sedentary behavior as outcome.

All models are reported crude and adjusted for age, sex, ethnicity, marital status, educational level, alcohol consumption, smoking, and diet. Furthermore, the four exposures were potential independent risk factors in the individual model; therefore, the variables were included as confounders in the individual model (online supplemental file 1). Post hoc analyses of comorbidities and MVPA excluding adjustment of BMI were performed to investigate the size of difference in MVPA between numbers of comorbidities allowing for adiposity levels to be different across comorbidities, because obesity is a strong determinant of a wide range of morbidities. In addition, cross-tabulations were conducted describing proportions and characteristics of participants with diabetes who were not following WHO recommendations of physical activity and their motivation for being more physically




Figure 1 Flow chart of all included participants in the primary analysis and the three secondary analyses from the Danish National Health Survey (DNHS) 2017. BMI, body mass index; SF-12, 12-item Short-Form Health Survey.

active. Statistically, weights provided by the DNHS were included in all models to account for non-response by certain population groups. The weights were computed by Statistics Denmark and account for differences such as age, sex, educational level, income, socioeconomic group, municipality of residence, marital status, ethnic background, healthcare utilization and research protection.³³ All statistical analyses were conducted in STATA/BE V.17.0 using an α level of 0.05, two sided.

RESULTS

Out of 183372 responses, 10216 participants reported they had diabetes. Of these, 6856 participants had complete data on outcome, exposures, and covariates in the primary analysis (figure 1).

Participant characteristics

Characteristics of participants with diabetes with and without complete data and participants with no known diabetes are presented in table 1. Due to differences in age and sex between diabetes status, direct standardization was performed on participants with no known diabetes based on the age and sex distribution among participants with diabetes. Only 60% of those with diabetes met the WHO recommendations compared with 70% of those without diabetes. Participants with diabetes had higher BMI, more comorbidities, higher perceived stress, and lower HRQoL.

Participant characteristics by level of habitual physical activity

Characteristics of participants with diabetes stratified by level of habitual physical activity according to WHO Guidelines of Physical Activity and Sedentary Behavior are presented in table 2. Out of 6341 participants, 2530 (38.8%) had optimal amount of habitual physical activity and 955 (16.4%) were inactive. Inactive participants had lower levels of education and more unhealthy eating habits, higher prevalence of obesity class II and III, higher prevalence of three or more comorbidities, higher perceived stress, and lower HRQoL.

Associations of MVPA and comorbidities

Table 3 represents the results from the primary analysis regarding associations between exposures and MVPA (hours/week) in participants with diabetes. Having diabetes with three or more comorbidities including mental health conditions had significantly lower weekly MVPA after adjustments (-0.48 hours/week, 95% CI -0.88 to -0.07) compared with individuals with no comorbidity, whereas three or more comorbidities excluding mental health conditions were non-significant after adjustments (-0.54 hours/week, 95% CI -0.76 to 0.04) compared with no comorbidity. Post hoc analyses showed significant lower weekly MVPA among individuals with two comorbidities and three or more comorbidities including mental health conditions without adjustment of BMI (-0.58 hours/week, 95% CI -0.98 to -0.18 and

Table 1 Characteristics of participants with diabetes with and without complete responses and no known diabetes

	Diabetes (complete data)	Diabetes (incomplete data)	No known diabetes
n (%) b21	6856 (8.4)*	10216 (8.8)*	173021 (91.2)
Marital status			
Married/living with partner	4928 (63.7)	6977 (61.1)	119029 (66.8)
Living alone	1928 (36.3)	3239 (38.9)	53992 (33.2)
Educational level			
Primary	1113 (19.8)	1848 (22.6)	18478 (16.0)
Upper secondary	4071 (58.9)	5563 (58.3)	86947 (55.3)
Higher	1672 (21.3)	2056 (19.1)	53911 (28.7)
Ethnic background			
Danish	6320 (88.1)	9387 (87.0)	159429 (91.6)
Western	187 (3.9)	273 (4.0)	5847 (4.3)
Non-Western	349 (7.9)	556 (9.0)	7745 (4.1)
Alcohol consumption (drinks consumed weekly)			
No alcohol	2161 (35.5)	3372 (39.4)	35883 (22.0)
Below low risk	3440 (47.3)	4551 (44.4)	88587 (54.0)
Above low risk	753 (10.4)	1027 (9.8)	24361 (16.1)
High risk	502 (6.8)	656 (6.4)	10967 (7.8)
Smoking			
Smoker	1309 (19.8)	1935 (20.8)	33017 (19.3)
Ex-smoker	3023 (42.9)	4388 (41.9)	52149 (39.2)
Never smoked	2524 (37.3)	3693 (37.3)	79529 (41.5)
Diet score			
Unhealthy	997 (14.9)	1522 (16.6)	23609 (16.5)
Medium healthy	4669 (68.2)	6325 (66.3)	105169 (65.5)
Healthy	1190 (16.9)	1667 (17.1)	29405 (17.9)
Obesity			
Underweight/normal	1492 (23.1)	2170 (23.2)	78160 (43.9)
Overweight	2584 (37.2)	3740 (36.9)	57078 (39.5)
Obese class I	1725 (24.6)	2436 (24.3)	18805 (12.6)
Obese class II	691 (9.5)	980 (9.8)	4980 (2.9)
Obese class III	364 (5.6)	511 (5.8)	2146 (1.1)
Comorbidities including mental health conditions			
No comorbidity	839 (11.6)	1205 (11.9)	59297 (27.3)
One comorbidity	1813 (25.1)	2653 (25.0)	53182 (30.2)
Two comorbidities	1989 (28.6)	2980 (28.4)	35944 (23.9)
Three or more comorbidities	2215 (34.5)	3378 (34.7)	24598 (18.6)
Perceived stress			
Low perceived stress	3739 (50.1)	4771 (46.5)	96095 (61.2)
Moderate perceived stress	2826 (44.9)	4207 (48.2)	57962 (36.1)
High perceived stress	291 (5.0)	410 (5.3)	5217 (2.7)
HRQoL			
Low physical HRQoL	4715 (71.8)	6335 (73.7)	60680 (50.6)
Low mental HRQoL	1845 (30.6)	2567 (32.8)	36103 (20.8)
Adherence to WHO recommendations of physical activity			
Following recommendations	2457 (59.1)	4902 (58.4)	102881 (68.9)
Not following recommendations	3884 (40.9)	3224 (41.6)	40578 (31.1)

Data are presented as n (%).

n is different due to variations in complete responses in each variable.

*Proportion of individuals with diabetes compared with proportion of individuals without diabetes.

HRQoL, health-related quality of life.

Table 2 Characteristics of participants with diabetes stratified by level of habitual physical activity

	Level of habitual physical activity*			
	Inactive	Inadequate	Adequate	Optimal
n (%)	955 (16.4)	1502 (23.5)	1354 (21.3)	2530 (38.8)
Sex				
Male	547 (57.8)	912 (60.1)	785 (57.8)	1558 (62.2)
Female	408 (42.2)	590 (39.9)	569 (42.2)	942 (37.8)
Age (years)	64 (53; 74)	64 (54; 72)	62 (53; 71)	63 (52; 71)
Marital status				
Married/living with partner	640 (59.8)	1097 (66.6)	981 (65.6)	1841 (66.1)
Living alone	315 (40.2)	405 (33.4)	373 (34.4)	689 (33.9)
Educational level				
Primary	211 (24.8)	229 (17.2)	191 (16.2)	365 (16.3)
Upper secondary	577 (59.6)	887 (60.0)	794 (60.4)	1488 (58.9)
High	167 (15.6)	386 (22.7)	369 (23.4)	677 (24.8)
Ethnic background				
Danish	877 (86.4)	1394 (88.7)	1256 (87.8)	2347 (88.9)
Western	33 (5.7)	45 (3.9)	27 (2.9)	67 (3.7)
Non-Western	45 (7.8)	63 (7.4)	71 (9.3)	116 (7.3)
Smoking				
Smoker	231 (24.5)	315 (22.8)	253 (19.9)	420 (18.8)
Ex-smoker	431 (43.4)	666 (42.6)	579 (40.0)	1117 (42.1)
Never smoked	293 (32.1)	521 (34.6)	522 (40.1)	993 (37.1)
Diet score				
Unhealthy	255 (26.3)	274 (20.1)	139 (10.5)	255 (10.0)
Medium healthy	594 (62.5)	1044 (68.8)	971 (70.9)	1702 (68.1)
Healthy	106 (11.2)	184 (11.1)	244 (18.7)	573 (21.9)
Alcohol consumption (drinks consumed weekly)				
No alcohol	415 (45.6)	453 (33.8)	369 (31.3)	716 (31.9)
Below low risk	380 (38.0)	736 (46.8)	758 (53.8)	1336 (50.1)
Above low risk	78 (7.7)	168 (10.7)	144 (9.3)	317 (11.9)
High risk	82 (8.7)	145 (8.8)	83 (5.6)	161 (6.1)
MVPA in leisure time (hours/week)	0 (0; 0)	1 (0.5; 1.5)	2.5 (2; 3.25)	6 (4.5; 9)
Total sedentary behavior (hours/day)	9 (6.5; 12)	8.5 (6; 12)	8.25 (6; 11.6)	8.2 (6; 11.7)
Obesity				
Underweight/normal	150 (16.5)	270 (18.4)	271 (21.3)	686 (28.9)
Overweight	331 (33.9)	536 (35.0)	514 (37.4)	1017 (38.6)
Obese class I	238 (24.2)	430 (28.2)	365 (26.1)	548 (21.4)
Obese class II	142 (14.4)	168 (11.0)	133 (9.5)	199 (7.6)
Obese class III	94 (11.0)	98 (7.4)	71 (5.7)	80 (3.5)
Comorbidities including mental health conditions				
No comorbidity	76 (8.3)	140 (8.8)	171 (13.5)	391 (16.3)
One comorbidity	191 (19.3)	373 (24.3)	381 (25.9)	738 (28.2)
Two comorbidities	272 (27.9)	461 (30.2)	420 (29.5)	689 (26.7)
Three or more comorbidities	416 (44.5)	528 (36.7)	382 (31.1)	712 (28.8)
Perceived stress				
Low perceived stress	411 (40.2)	778 (46.3)	759 (50.6)	1566 (58.9)

Continued

Table 2 Continued

	Level of habitual physical activity*			
	Inactive	Inadequate	Adequate	Optimal
Moderate perceived stress	455 (50.0)	657 (47.8)	546 (44.9)	897 (37.7)
High perceived stress	89 (9.8)	67 (5.9)	49 (4.5)	67 (3.3)
HRQoL				
Low physical HRQoL	820 (86.3)	1137 (76.8)	930 (70.0)	1450 (58.8)
Low mental HRQoL	955 (43.6)	1502 (34.1)	1354 (28.1)	2530 (23.4)

n=6341.
 Data are presented as number and proportion (%) except from age, MVPA and sedentary behavior, which are presented as median and IQR.
 *Level of habitual physical activity was categorized in accordance with the 'WHO Guidelines of Physical Activity and Sedentary Behavior': (1) inactive=0 hour/week, (2) inadequate=participants do not meet recommendations (150–300 min/week of moderate intensity or 75–150 min/week of vigorous intensity), (3) adequate=participants meet recommendations, but not more than 300 min/week of moderate intensity or 150 min/week of vigorous intensity, and (4) optimal=participants with more than 300 min/week of moderate intensity or 150 min/week of vigorous intensity of physical activity.
 HRQoL, health-related quality of life; MVPA, moderate to vigorous physical activity.

–0.85 hours/week, 95% CI –1.25 to –0.45) compared with individuals with no comorbidity. The analyses excluding mental health conditions showed also significant results without adjustment of BMI (–0.51 hours/week, 95% CI –0.89 to –0.12 and –0.72 hours/week, 95% CI –1.12 to –0.32).

Associations of MVPA and obesity

Overweight or obese (class I–III) individuals engaged in significantly less MVPA after adjustments (overweight: –0.61 hours/week, 95% CI –0.94 to –0.29; obese class I: –1.08 hours/week, 95% CI –1.43 to –0.74; obese class II: –1.43 hours/week, 95% CI –1.85 to –1.01; obese class III: –1.98 hours/week, 95% CI –2.49 to –1.47) compared with underweight and normal weight individuals.

Associations of MVPA and perceived stress

Moderate and high perceived stress were significantly associated with lower weekly MVPA after adjustments (–0.59 hours/week, 95% CI –0.83 to –0.34 and –1.76 hours/week, 95% CI –2.18 to –1.34) compared with low perceived stress.

Associations of MVPA and HRQoL

At last, low physical and mental HRQoL were associated with lower weekly MVPA after adjustments (–0.93 hours/week, 95% CI –1.19 to –0.66 and –0.39 hours/week, 95% CI –0.71 to –0.08) compared with moderate or high HRQoL. The pattern of association of comorbidity, obesity, and stress with MVPA appeared consistent with an inverse graded relationship.

Adherence to WHO recommendations

The estimated associations of the logistic regression models between exposures and adherence to WHO recommendations appeared similar to the linear models (online supplemental file 3).

Associations of sedentary behavior and determinants

Results of sedentary behavior showed significant associations between obese class II and III and higher weekly sedentary behavior compared with normal weight individuals with diabetes (for more details see online supplemental file 4).

Motivation for being more physically active

Table 4 shows the distribution and characteristics of participants with diabetes who are not following WHO recommendations of physical activity and their motivation for being more physically active. Younger participants were more motivated for being more physically active. In addition, the proportion of motivated participants was highest among obese class II and III compared with obese class I, overweight, and underweight/normal. Distribution and characteristics of participants with diabetes who are following the WHO recommendations of physical activity and their motivation for being more physically active are available in online supplemental file 5.

DISCUSSION

We found that 40% of individuals with diabetes from this nationwide population were not adherent to WHO recommendations for physical activity as measured by self-report. Inactive individuals with diabetes were less educated, had more unhealthy eating and smoking habits, higher BMI, more comorbidities, higher perceived stress, and lower HRQoL. The primary analysis showed that higher BMI, higher perceived stress, lower HRQoL, and having two comorbidities and three or more comorbidities were associated with lower weekly MVPA. However, when comorbidities only included somatic conditions, the association was only significant when participants had three or more comorbidities. Obesity, high perceived stress, and low physical HRQoL were the most remarkable determinants of low physical activity with mean differences of at least

Table 3 Linear regressions on the associations of determinants with MVPA in participants with diabetes**Therefore, the aims of this**

	MVPA (hours/week)					
	Crude		Multivariable adjusted		Multivariable adjusted (excluding BMI)†	
	β (95% CI)*	P value	β (95% CI)	P value	β (95% CI)	P value
Comorbidities including mental health conditions‡						
No comorbidity	(Reference)		(Reference)		(Reference)	
One comorbidity	-0.38 (-0.79 to 0.02)	0.064	-0.08 (-0.48 to 0.34)	0.688	-0.29 (-0.69 to 0.11)	0.161
Two comorbidities	-0.73 (-1.13 to -0.34)	0.001§	-0.31 (-0.70 to 0.09)	0.135	-0.58 (-0.98 to -0.18)	0.004§
Three or more comorbidities	-1.04 (-1.44 to -0.65)	0.001§	-0.48 (-0.88 to -0.07)	0.021§	-0.85 (-1.25 to -0.45)	0.001§
Comorbidities excluding mental health conditions‡						
No comorbidity	(Reference)		(Reference)		(Reference)	
One comorbidity	-0.27 (-0.66 to 0.12)	0.0175	-0.04 (-0.34 to 0.43)	0.823	-0.16 (-0.55 to 0.23)	0.417
Two comorbidities	-0.65 (-1.03 to -0.27)	0.001§	-0.23 (-0.61 to 0.16)	0.243	-0.51 (-0.89 to -0.12)	0.010§
Three or more comorbidities	-0.91 (-1.29 to -0.53)	0.001§	-0.54 (-0.76 to 0.04)	0.081	-0.72 (-1.12 to -0.32)	0.001§
Obesity¶						
Underweight/normal	(Reference)		(Reference)			
Overweight	-0.61 (-0.93 to -0.29)	0.001§	-0.61 (-0.94 to -0.29)	0.001§		
Obese class I	-1.12 (-1.47 to -0.77)	0.001§	-1.08 (-1.43 to -0.74)	0.001§		
Obese class II	-1.49 (-1.92 to -1.07)	0.001§	-1.43 (-1.85 to -1.01)	0.001§		
Obese class III	-2.05 (-2.58 to -1.52)	0.001§	-1.98 (-2.49 to -1.47)	0.001§		
Stress**						
Low perceived stress	(Reference)		(Reference)			
Moderate perceived stress	-0.68 (-0.92 to -0.44)	0.001§	-0.59 (-0.83 to -0.34)	0.001§		
High perceived stress	-1.90 (-2.29 to -1.51)	0.001§	-1.76 (-2.18 to -1.34)	0.001§		
HRQoL††						
<i>Physical score</i>						
Moderate to high physical HRQoL	(Reference)		(Reference)			
Low physical HRQoL	-1.35 (-1.59 to -1.09)	0.001§	-0.93 (-1.19 to -0.66)	0.001§		
<i>Mental score</i>						
Moderate to high mental HRQoL	(Reference)		(Reference)			
Low mental HRQoL	-0.90 (-1.15 to -0.65)	0.001§	-0.39 (-0.71 to -0.08)	0.015§		

n=6856.
 Crude and adjusted models were weighted for non-response.
 *β coefficients and 95% CIs represent mean difference in MVPA (hours/week) compared with the reference.
 †Post hoc analyses of comorbidities and MVPA excluding adjustment of BMI.
 ‡Each multivariable model was adjusted as follows: age, sex, ethnicity, educational level, marital status, diet, smoking, alcohol consumption, BMI.
 §Significant results (p<0.05).
 ¶Each multivariable model was adjusted as follows: age, sex, ethnicity, educational level, marital status, diet, smoking, alcohol consumption, comorbidity.
 **Each multivariable model was adjusted as follows: age, sex, educational level, marital status, smoking, BMI, comorbidity.
 ††Each multivariable model was adjusted as follows: age, sex, educational level, marital status, smoking, BMI, comorbidity, stress.
 BMI, body mass index; HRQoL, health-related quality of life; MVPA, moderate to vigorous physical activity.

1 hour of lower weekly MVPA compared with references. Post hoc analyses showed significant lower weekly MVPA among individuals with two comorbidities and three or more comorbidities.

Among individuals with diabetes who did not meet the WHO recommendations, 60% were motivated for being more physically active, with higher motivation among young, middle-aged and obese individuals.

The prevalence of adherence to the WHO recommendations found in this study is inconsistent with two previous cross-sectional studies outside Denmark. A study by Martinez-Harvell *et al*¹¹ from 2020 found that 53.5% of

25 980 participants with diabetes from the USA did not meet the recommendations of physical activity (150 min/week of MVPA, or 75 min/week of vigorous intensity at least three times/week). A Scottish study by Salman *et al*³⁴ from 2019 found that 65.9% of 1259 participants did not meet the recommended levels of physical activity (150 min/week moderate or 75 min/week vigorous physical activity or equivalent combination). The inconsistency might be explained by differences in how physical activity was quantified and the populations included. Our study included a large population with diabetes from all regions of Denmark, which increases the generalizability to the

Table 4 Distribution of insufficiently physically active participants and their motivation for being more physically active

	Motivated	Not motivated	Don't know
n (%)	1914 (59.8)	604 (19.2)	631 (21.0)
Sex			
Male	1089 (59.6)	389 (20.3)	360 (20.1)
Female	825 (60.1)	215 (17.7)	271 (22.2)
Age (categories)			
16–34 years	59 (72.5)	6 (9.2)	13 (18.3)
35–54 years	414 (77.1)	47 (9.4)	72 (13.6)
55–64 years	545 (65.6)	131 (14.9)	147 (19.4)
65–74 years	592 (56.1)	211 (21.6)	206 (22.3)
≥75 years	304 (41.4)	209 (30.8)	193 (27.8)
Obesity			
Underweight/normal	285 (52.3)	138 (26.7)	106 (21.0)
Overweight	642 (57.6)	244 (22.6)	214 (19.8)
Obese class I	489 (58.4)	141 (16.6)	192 (25.0)
Obese class II	289 (71.2)	44 (11.2)	65 (17.6)
Obese class III	177 (72.6)	27 (9.6)	41 (18.9)
Comorbidities including mental health conditions			
No comorbidity	162 (61.5)	64 (23.0)	43 (15.5)
One comorbidity	412 (59.7)	141 (18.9)	151 (21.3)
Two comorbidities	556 (57.5)	184 (20.2)	193 (22.3)
Three or more comorbidities	784 (61.0)	215 (18.0)	244 (21.0)
Stress			
Low perceived stress	820 (59.5)	296 (21.5)	258 (19.0)
Moderate perceived stress	871 (60.6)	248 (18.1)	281 (21.0)
High perceived stress	121 (62.0)	22 (11.9)	47 (26.1)
HRQoL			
<i>Physical score</i>			
Moderate to high physical HRQoL	359 (66.9)	107 (18.9)	83 (14.2)
Low physical HRQoL	1395 (59.8)	415 (18.2)	473 (22.0)
<i>Mental score</i>			
Moderate to high mental HRQoL	1103 (59.3)	373 (20.5)	353 (20.2)
Low mental HRQoL	651 (63.8)	149 (15.1)	203 (21.1)

n=3149 (n differs due to variations in complete responses in each variable).
 Data are presented as n (%).
 HRQoL, health-related quality of life.

whole country thereby extending previous findings. Even though our results are inconsistent with previous studies, our results still indicate that physical activity interventions targeting individuals with diabetes are needed.

We found that individuals with higher perceived stress, lower HRQoL and comorbidities including mental health conditions are less likely to engage in regular physical activity. These findings support previous studies showing an association between mental health conditions, distress, and physical activity among individuals with diabetes.^{35 36} Individuals suffering from mental health conditions or distress may experience challenges

concerning self-care in managing the diabetes condition compared with individuals with only somatic conditions. We also found that higher BMI and lower HRQoL were associated with lower MVPA, which support the results from other studies.^{11 34 37} Individuals with poor HRQoL and elevated BMI may suffer from discomfort, fatigue, shame and fear of exercise as barriers to engage in regular physical activity.^{38–40}

No other studies have described and investigated the characteristics and correlates of physical activity among individuals with diabetes who are completely inactive corresponding to 0 hour of weekly MVPA. Results showed

that inactive individuals with diabetes had lower levels of education, which indicates that social inequality in engagement in physical activity is present among individuals with diabetes. Social inequality is highly associated with prevalence and complications of type 2 diabetes.^{41 42} This includes a 10% higher risk of developing severe diabetes-related complications, and 26% higher risk of premature mortality among individuals with lower levels of education.⁴³ With a large sociodemographic and geographic represented population, the results of the present study highlight the nationwide challenges of social inequality among individuals with diabetes and their ability to engage in regular physical activity.

Clinical implications

Based on our results, healthcare providers may need more targeted approaches to reach those in most need of physical activity interventions to increase the reach and motivation as well as the success of the interventions. Tailored interventions that are adapted to individuals with lower levels of education, higher BMI, more comorbidities, higher perceived stress, and lower HRQoL might be needed to help improve diabetes self-management and physical activity level among all individuals with diabetes and thereby help prevent the development of other chronic conditions, diabetes-related complications and premature mortality. It seems important to develop or adapt current diabetes management and physical interventions to increase reach and motivation on an individual level among insufficiently active individuals to increase the health of the population and prevent the development of other chronic conditions. In our study, most individuals with diabetes who were not sufficiently active were motivated to increase their physical activity level (60%), which suggests that other factors prevent them from increasing their physical activity level. Previous studies have found that mental health, lack of time and energy, working schedule, economic circumstances, and social support are reasons for not being more physically active.^{38 44–46} These reasons are also important to consider when designing and promoting a physical activity intervention.

Strengths and limitations

The present study is based on a relatively large nationwide sample, which strengthens the generalizability of the results. The sample of participants with self-reported diabetes was 5.6%, which is slightly higher than the prevalence of diabetes in a recently published study in Denmark that found an overall prevalence of 0.5% of type 1 diabetes and 4.4% of type 2 diabetes in 2016.⁴⁷ Another strength of this study is the data on perceived stress, HRQoL and weekly MVPA, which were provided with validated questionnaires.^{21 26 28} Furthermore, this study applied statistically weights provided by the DNHS to account for non-response and ensure a nationally representative analysis.³³

The results of this study should be interpreted with the following limitations in mind. It was not possible to distinguish between type 1 and type 2 diabetes among participants,

which would have been preferable due to etiological differences. Since type 2 diabetes accounts for 85%–95% of diabetes cases in high-income countries,⁴⁸ the findings of this study will primarily be applicable to this patient group. In addition, information regarding diabetes complications and duration of the diabetes disease was not available, and the sample answering could potentially be skewed. Data were collected through self-reported questionnaires, which introduces information bias, including social desirability bias regarding lifestyle questions, such as diet, alcohol, smoking, and physical activity. Individuals with certain negative health conditions and behaviors are more likely to refrain from responding to health surveys.⁴⁹ Thus, the absolute prevalence of diabetes might be underestimated, and the presented characteristics might not be representative for all subgroups of individuals with diabetes, while the estimated levels of MVPA might be overestimated due to desirability bias. Despite the use of statistical weights to reduce the possible impact of non-response bias on the estimates, citizens living alone, or with non-Western background or at younger age were not highly represented in the study population affecting the generalizability of the findings.

The comorbidity variable created in this study does not include all the conditions, and not all conditions affect the individual's ability to be physically active. Questions regarding weekly MVPA are reported to provide an acceptable but weak estimate of participants' level of physical activity, which questions the absolute distribution of adherence to WHO recommendations of physical activity.²¹ In addition, results might be different if assessment of MVPA was following the WHO recommendations from 2020, where physical activity of less than 10 min bouts has been removed.⁴ Finally, the cross-sectional study design is a limitation, as reverse causation, uncontrolled confounding, and information bias cannot be ruled out. A prospective study design and/or objective measurement tools are suggested for further research to support the findings of this study.

CONCLUSION

In this nationwide survey, we found that having comorbidities, higher BMI, higher perceived stress, and lower HRQoL were significantly associated with lower MVPA. Suffering from obesity, high perceived stress, and low physical HRQoL were the most remarkable determinants of low MVPA. Furthermore, we found that 40% of individuals with diabetes do not engage regularly in adequate physical activity, and 60% of those who do not meet the WHO recommendations for physical activity are motivated for being more physically active.

Our study suggests that subgroups of individuals with diabetes are at risk of physical inactivity and might need additional support or tailored interventions to increase their physical activity level.

Contributors All authors contributed to formulation of the study question and design. PLK and AG supervised the statistical analyses, and SRM conducted the statistical analyses. All authors contributed to the interpretation of the study results.

SRM wrote the first draft of the manuscript. STS reviewed and commented on all drafts of the manuscript. All authors contributed to the critical revision and final manuscript. All authors read and approved the final manuscript. SRM is the guarantor and takes responsibility for the conduct and the overall content of the article.

Funding SRM is funded by a program grant from Region Zealand (Exercise First), a research grant from Næstved-Slagelse-Ringsted Hospitals' Research Fund (A986), a research grant from the Health Science Research Fund of Region Zealand (A1136), and a research grant from Steno Diabetes Center Sjælland. STS is currently funded by a program grant from Region Zealand (Exercise First) and two grants from the European Union's Horizon 2020 research and innovation program: one from the European Research Council (MOBILIZE, grant agreement number: 801790) and the other from ESCAPE (grant agreement number: 945377). AG is funded by the European Research Council (SCREENS, grant agreement number: 716657), the Novo Nordisk Foundation (grant number: NNF20SH0062965), and TrygFonden (grant numbers: 130081 and 115606). MR-L is funded by TrygFonden. The Danish National Health Survey was funded by the Capital Region, Region Zealand, the South Denmark Region, the Central Denmark Region, the North Denmark Region, Ministry of Health and the National Institute of Public Health, University of Southern Denmark.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants and was approved by the Danish Data Protection Agency through the University of Southern Denmark (journal number: 11.397). Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. All data used in the current study were derived from the DNHS. Data are available through application to the National Institute of Public Health, University of Southern Denmark.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Sofie Rath Mortensen <http://orcid.org/0000-0002-9550-1507>


REFERENCES

- World Health Organization. *Physical activity strategy for the who European region 2016-2025*. Regional Office for Europe, 2015.
- Tielemans SMAJ, Soedamah-Muthu SS, De Neve M, et al. Association of physical activity with all-cause mortality and incident and prevalent cardiovascular disease among patients with type 1 diabetes: the EURODIAB prospective complications study. *Diabetologia* 2013;56:82–91.
- Colberg SR, Sigal RJ, Yardley JE, et al. Physical activity/exercise and diabetes: a position statement of the American diabetes association. *Diabetes Care* 2016;39:2065–79.
- World Health Organization. *Who guidelines on physical activity and sedentary behaviour*; 2020.
- American Diabetes Association. *Standards of medical care in diabetes—2011*. *Diabetes Care* 2011;34 Suppl 1:S11–61.
- Huang ES, Brown SES, Ewigman BG, et al. Patient perceptions of quality of life with diabetes-related complications and treatments. *Diabetes Care* 2007;30:2478–83.
- Jing X, Chen J, Dong Y, et al. Related factors of quality of life of type 2 diabetes patients: a systematic review and meta-analysis. *Health Qual Life Outcomes* 2018;16:189.
- Liu NF, Brown AS, Foliass AE, et al. Stigma in people with type 1 or type 2 diabetes. *Clin Diabetes* 2017;35:27–34.
- van der Berg JD, Stehouwer CDA, Bosma H, et al. Associations of total amount and patterns of sedentary behaviour with type 2 diabetes and the metabolic syndrome: the maastricht study. *Diabetologia* 2016;59:709–18.
- Zhao G, Ford ES, Li C, et al. Compliance with physical activity recommendations in US adults with diabetes. *Diabet Med* 2008;25:221–7.
- Martinez-Harvell G, Goluboff F, Rodriguez P, et al. Predictors of adherence to physical activity guidelines in patients with diabetes mellitus in the US in 2017: an exploratory analysis. *Prim Care Diabetes* 2020;14:645–53.
- Stults-Kolehmainen MA, Sinha R. The effects of stress on physical activity and exercise. *Sports Med* 2014;44:81–121.
- Cooper AR, Page A, Fox KR, et al. Physical activity patterns in normal, overweight and obese individuals using minute-by-minute accelerometry. *Eur J Clin Nutr* 2000;54:887–94.
- Durstine JL, Gordon B, Wang Z, et al. Chronic disease and the link to physical activity. *J Sport Health Sci* 2013;2:3–11.
- Anokye NK, Trueman P, Green C, et al. Physical activity and health related quality of life. *BMC Public Health* 2012;12:624.
- Daousi C, Casson IF, Gill GV, et al. Prevalence of obesity in type 2 diabetes in secondary care: association with cardiovascular risk factors. *Postgrad Med J* 2006;82:280–4.
- Centers for Disease Control and Prevention. Risk factors for diabetes-related complications; 2020. <https://www.cdc.gov/diabetes/data/statistics-report/risks-complications.html> [Accessed 13 Apr 2021].
- Vandenbroucke JP, von Elm E, Altman DG, et al. Strengthening the reporting of observational studies in epidemiology (STROBE): explanation and elaboration. *Int J Surg* 2014;12:1500–24.
- Christensen AI, Ekholm O, Glümer C, et al. The danish national health survey 2010. Study design and respondent characteristics. *Scand J Public Health* 2012;40:391–7.
- Christensen AI, Lau CJ, Kristensen PL, et al. The danish national health survey: study design, response rate and respondent characteristics in 2010, 2013 and 2017. *Scand J Public Health* 2022;50:180–188.
- Danquah IH, Skov SS, Callesen BR, et al. *Validering af spørgsmålt om fysisk aktivitet og siddetid - Til brug for national monitoring*. Statens Institut for Folkesundhed, 2016.
- Willadsen TG, Siersma V, Nicolaisdóttir DR, et al. Multimorbidity and mortality: A 15-year longitudinal registry-based nationwide danish population study. *J Comorb* 2018;8:2235042X18804063.
- Difrancesco S, Lamers F, Riese H, et al. Sleep, circadian rhythm, and physical activity patterns in depressive and anxiety disorders: A 2-week ambulatory assessment study. *Depress Anxiety* 2019;36:975–86.
- Izawa KP, Watanabe S, Oka K, et al. Association between mental health and physical activity in patients with chronic heart failure. *Disabil Rehabil* 2014;36:250–4.
- World Health Organization. *Obesity: preventing and managing the global epidemic : report of a WHO consultation*. Geneva; 2000.
- Eskildsen A, Dalgaard VL, Nielsen KJ, et al. Cross-cultural adaptation and validation of the danish consensus version of the 10-item perceived stress Scale. *Scand J Work Environ Health* 2015;41:486–90.
- Cohen S, Kamarck T, Mermelstein R. A global measure of perceived stress. *J Health Soc Behav* 1983;24:385–96.
- Ware J, Kosinski M, Keller SD. A 12-Item short-form health survey: construction of scales and preliminary tests of reliability and validity. *Med Care* 1996;34:220–33.
- Gandek B, Ware JE, Aaronson NK, et al. Cross-Validation of item selection and scoring for the SF-12 health survey in nine countries: results from the IQOLA project. international quality of life assessment. *J Clin Epidemiol* 1998;51:1171–8.
- Ware J, Jr, Kosinski M, Keller SD. SF-12 : how to score the SF-12 physical and mental health summary scales; 1995.
- Shrier I, Platt RW. Reducing bias through directed acyclic graphs. *BMC Med Res Methodol* 2008;8:70.
- Toft U, Kristoffersen LH, Lau C, et al. The dietary quality score: validation and association with cardiovascular risk factors: the Inter99 study. *Eur J Clin Nutr* 2007;61:270–8.
- Särndal C-E, Lundström S. *Estimation in surveys with nonresponse*. Hoboken, 2005.
- Salman A, Ukwaja KN, Alkhatib A. Factors associated with meeting current recommendation for physical activity in Scottish adults with diabetes. *Int J Environ Res Public Health* 2019;16. doi:10.3390/ijerph16203857. [Epub ahead of print: 12 10 2019].


- 35 Nyboe L, Lund H. Low levels of physical activity in patients with severe mental illness. *Nord J Psychiatry* 2013;67:43–6.
- 36 Mulligan K, McBain H, Lamontagne-Godwin F, *et al.* Barriers to effective diabetes management - a survey of people with severe mental illness. *BMC Psychiatry* 2018;18:165.
- 37 Plotnikoff RC, Taylor LM, Wilson PM, *et al.* Factors associated with physical activity in Canadian adults with diabetes. *Med Sci Sports Exerc* 2006;38:1526–34.
- 38 Korkiakangas EE, Alahuhta MA, Laitinen JH. Barriers to regular exercise among adults at high risk or diagnosed with type 2 diabetes: a systematic review. *Health Promot Int* 2009;24:416–27.
- 39 Shultz JA, Sprague MA, Branen LJ, *et al.* A comparison of views of individuals with type 2 diabetes mellitus and diabetes educators about barriers to diet and exercise. *J Health Commun* 2001;6:99–115.
- 40 Egan AM, Mahmood WAW, Fenton R, *et al.* Barriers to exercise in obese patients with type 2 diabetes. *QJM* 2013;106:635–8.
- 41 Karter AJ, Stevens MR, Brown AF, *et al.* Educational disparities in health behaviors among patients with diabetes: the translating research into action for diabetes (triad) study. *BMC Public Health* 2007;7:308.
- 42 Andersen MB, Bjørkman A-SD, Pedersen M, *et al.* Social inequality in lifestyle, motivation to change lifestyle and received health advice in individuals with diabetes: a nationwide study. *Scand J Public Health* 2020;48:847–54.
- 43 Sortsø C, Lauridsen J, Emneus M, *et al.* Social inequality in diabetes patients' morbidity patterns from diagnosis to death - a danish register-based investigation. *Scand J Public Health* 2018;46:92–101.
- 44 Martin CG, Pomares ML, Muratore CM, *et al.* Level of physical activity and barriers to exercise in adults with type 2 diabetes. *AIMS Public Health* 2021;8:229–39.
- 45 Cradock KA, Quinlan LR, Finucane FM, *et al.* Identifying barriers and facilitators to diet and physical activity behaviour change in type 2 diabetes using a design probe methodology. *J Pers Med* 2021;11. doi:10.3390/jpm11020072. [Epub ahead of print: 26 01 2021].
- 46 Lidegaard LP, Schwennesen N, Willaing I, *et al.* Barriers to and motivators for physical activity among people with type 2 diabetes: patients' perspectives. *Diabet Med* 2016;33:1677–85.
- 47 Carstensen B, Rønn PF, Jørgensen ME. Prevalence, incidence and mortality of type 1 and type 2 diabetes in Denmark 1996–2016. *BMJ Open Diabetes Res Care* 2020;8:e001071.
- 48 International Diabetes Federation. *IDF diabetes atlas*. 10th edn. Brussels, Belgium: International Diabetes Federation, 2021.
- 49 Thygesen LC, Johansen C, Keiding N, *et al.* Effects of sample attrition in a longitudinal study of the association between alcohol intake and all-cause mortality. *Addiction* 2008;103:1149–59.

Supplement 1


Fig. s1. DAG of the assumed causal relations between comorbidity and MVPA.




Directed Acyclic Graph (DAG) of the assumed causal relations between exposure (comorbidity), outcome (MVPA) and covariates in the primary analyses. Green line reflects the assumed causal paths. Red line and variables reflect possible confounding paths.

Fig. s2. DAG of the assumed causal relations between obesity and MVPA.

Directed Acyclic Graph (DAG) of the assumed causal relations between exposure (obesity), outcome (MVPA), and covariates in the primary analyses. Green line reflects the assumed causal paths. Red line and variables reflect possible confounding paths.

Fig. s3. DAG of the assumed causal relations between stress and MVPA.

Directed Acyclic Graph (DAG) of the assumed causal relations between exposure (stress), outcome (MVPA), and covariates in the primary analyses. Green line reflects the assumed causal paths. Red line and variables reflect possible confounding paths.

Fig. s4. DAG of the assumed causal relations between HRQoL and MVPA.

Directed Acyclic Graph (DAG) of the assumed causal relations between exposure (HRQoL), outcome (MVPA), and covariates in the primary analyses. Green line reflects the assumed causal paths. Red line and variables reflect possible confounding paths.

Supplement 2

Statistical analysis plan

Running title

Determinants of physical activity among individuals with diabetes: a cross-sectional study

Authors

Sofie Rath Mortensen, Peter Lund Kristensen, Cathrine Juel Lau, Mathias Ried-Larsen, Anders Grøntved, Søren Thorgaard Skou

Contributors and roles in the SAP

Responsible for writing the SAP: PhD-student Sofie Rath Mortensen (PT, MSc)

Brief background

Engagement in regular physical activity is a cornerstone of type 2 diabetes management to prevent long-term diabetes complications and premature mortality. Morbidity, obesity, stress, and health-related quality of life are all factors that may determine decreased habitual physical activity in general population. Given that individuals with diabetes are at high risk of suffering from these factors, it is likely that individuals with diabetes have different physical activity patterns. No previous large-scale studies have provided a detailed description of habitual physical activity among individuals with diabetes. Availability of such information would be an important resource for planning future treatment courses taking individual characteristics, needs and preferences into account when designing and promoting a physical activity intervention.

Primary aim

To provide descriptive data on habitual physical activity and investigate the association of morbidity, obesity, stress, and health-related quality of life with physical activity among individuals with diabetes based on data from The Danish National Health Survey from 2017.

Secondary aim

To investigate the association of morbidity, obesity, stress, and health-related quality of life with sedentary behavior among individuals with diabetes based on data from The Danish National Health Survey from 2017.

Methods

Study design

The study design is cross-sectional. Reporting of the study will be followed by the STROBE checklist (1).

Data sources

Responders of The Danish National Health Survey (DNHS) in 2017 will be included. The DNHS was based on six mutually exclusive random subsamples; one in each of the five Danish administrative regions, and one national sample. 312,349 individuals were invited via a secure electronical mail service (Digital Post) or regular postal service to participate in the survey. Mandatory questions of the survey for all subsamples will be used in this project (appendix 5).

Variables

Exposures

Comorbidity (excluding diabetes) (categorical)

- Self-reported information on selected long-term conditions (excluding diabetes) and sequela from question 10-11 were used to assess comorbidity. Respondents reported whether they have or have had selected long-term conditions. If respondents reported they have had a long-term condition, they reported whether they were suffering from sequelae due to the specific long-term condition.
- The definition of multimorbidity will be based on diagnoses organized in 10 groups of different body systems according to Willadsen et al. (2). In this study there will only be 7 groups due to lack of information on gastrointestinal and genitourinary diseases from the survey, and the endocrine body system will be excluded since diabetes is not a part of the comorbidity-variable in the present study. The 7 groups will be: 1) Lung (asthma and bronchitis), 2) Musculoskeletal (osteoarthritis, rheumatoid arthritis, osteoporosis, and back diagnoses), 3) Mental (temporary mental disease and long-term mental disease), 4) Cancer, 5) Neurological (stroke and migraine), 6) Cardiovascular (hypertension, angina pectoris, and myocardial infarction), and 7) Sensory organs (tinnitus and cataract).
- The variable will be categorized as follows: 1) Have diabetes and no comorbidities, 2) Have diabetes and one comorbidity from one body system, 3) Have diabetes and two comorbidities from two different body systems, 4) Have diabetes and three or more comorbidities from at least three different body systems. Suffering from several long-term conditions within the same body system, e.g., hypertension and myocardial infarction, will still only count as one comorbidity.
- Studies have found that a decreased mental health status among individuals with comorbidities is associated with lower physical activity (3, 4). Therefore, a variable differentiating between having diabetes and comorbidities with and without a mental disease will be created.

Obesity (categorical)

- Self-reported data on body weight and height were obtained from question 38-39.
- Body mass index (BMI) (kg/m^2) will be calculated and categorized into the following groups: underweight/normal weight ($\text{BMI} < 25.0$), overweight ($\text{BMI} \geq 25.0 - < 30.0$), obese class I ($\text{BMI} \geq 30.0 - < 35.0$), obese class II ($\text{BMI} \geq 35.0 - < 40.0$), and obese class III ($\text{BMI} \geq 40$), as defined by the World Health Organization (5).

Stress (categorical)

- Self-reported psychological stress was obtained with Cohen's 10-item Perceived Stress Scale (PSS) (6) from question 8.
- The PSS will be categorized into three as follows:
 - o Scores ranging from 0-13 will be considered low perceived stress
 - o Scores ranging from 14-26 will be considered moderate perceived stress
 - o Scores ranging from 27-40 will be considered high perceived stress

Health-related quality of life (categorical)

- Self-reported health-related quality of life was obtained with the 12-item Short-Form Health Survey (SF-12) from question 1-7 (7).
- The SF-12 score is divided into a physical and mental score and each variable was dichotomized following the recommendations (8):
 - o Physical: A score of 50 or less determines a physical condition

- Mental: A score of 42 or less may be indicative of clinical depression

Primary outcome

Moderate to vigorous physical activity (MVPA) h/week (continuous)

- Respondents reported hours and minutes spent weekly on moderate and vigorous physical activities in question D-E.
- MVPA in hours and minutes weekly will be calculated into hours.

Secondary outcomes

Adherence to the WHO recommendations of physical activity and sedentary behavior (binary)

- Adherence to the WHO recommendations of physical activity and sedentary behavior was assessed with information regarding hours and minutes spent on weekly MVPA from question D-E.
- Following the WHO recommendations: ≥ 150 mins/week of moderate intensity or ≥ 75 mins/week of vigorous intensity or an appropriate combination hereby.
- Not following the WHO recommendations: < 150 - 300 mins/week of moderate intensity or < 75 mins/week of vigorous intensity (9).

Total sedentary behavior h/day (continuous)

- In question F, respondents reported hours and minutes spent daily sedentary on the following: work, transport, screen time and other sedentary activities, such as eating, social gatherings etc.
- Total sedentary behavior h/day will be calculated by adding hours and minutes from the abovementioned categories and then calculated into hours.

Covariates

The listed covariates below are suggested to adjust for in the analyses due to possible confounding: Age, sex, ethnicity, marital status, educational level, alcohol consumption, smoking, and diet may be independent risk factors of the four exposures (comorbidity, obesity, stress, and health-related quality of life) and the outcome (physical activity).

Furthermore, the four exposures may be independent risk factors in the individual model, therefore, the variables comorbidity, obesity, stress, and health-related quality of life will be included as potential confounders in each model. Directed Acyclic Graphs (DAGs) of the assumed causal relations between exposures and outcome of the primary analysis have been created (appendix 1-4).

Age (continuous)

- Calculated age at Jan 11th, 2017, from CPR-register

Sex (binary)

- From CPR-register
- Categorizations: Male, Female

Ethnicity (categorical)

- From CPR-register
- Categorizations: Born in Denmark, Western, Non-Western

Marital status (binary)

- Based on question G and CPR-register

- Marital status was categorized into: Married or Living with partner, Living alone

Educational level (categorical)

- Educational level was defined as the highest completed education, and the variable was obtained from question 50-51
- Educational level was categorized into the following: Primary (<10 years), Upper secondary or vocational (10-12 years), Higher education (≥ 13 years)

Alcohol consumption (categorical)

- Respondents were asked about their weekly alcohol consumption in question 24
- Alcohol consumption was categorized in accordance with the national recommendations from the Danish Health Authority and the risk of alcohol-related diseases for men and women:
 - o No alcohol (0 drinks – both men and women)
 - o Below low risk (men >0 & <14 drinks, women >0 & <7 drinks)
 - o Above low risk (men ≥ 14 & ≤ 21 drinks, women ≥ 7 & ≤ 14 drinks)
 - o High risk (men >21 drinks, women >14 drinks)

Smoking (categorical)

- Information regarding respondents smoking habits was obtained from question 13
- Smoking was categorized into the following: Smoker, Ex-smoker, Never smoked

Diet (categorical)

- Frequency and self-rated dietary habits were obtained from question 27-32
- Diet was calculated and categorized in accordance with the Dietary Quality Score: Unhealthy, Medium healthy, Healthy (10)

Statistical methods

Statistical software

STATA/BE 17.0

Sample size

The survey was fully or partially completed by 183,372 respondents (58.7 %). Respondents with diabetes and complete data on outcome, exposures and covariates will be considered as eligible for the analytical sample of the primary aim. Following the categorization of diabetes from DNHS, respondents were defined as “Having diabetes” if they had answered “I have diabetes now” or “I have had diabetes” and “I suffer from sequelae due to the diabetes”. According to this categorization, 10,216 respondents have diabetes. Respondents with complete data on primary outcome, exposures and covariates will be 6,856. The secondary analysis will have a smaller sample size due to fewer complete responses.

Missing data

The percentage of missing data for the primary outcome (MVPA) among individuals with diabetes is 11.8 %. For the secondary outcome (WHO recommendations) the percentage of missing data is 20.5 %, and for the outcome total sedentary behavior the percentage of missing data is 20.0 %. To reduce the possible impact of non-response bias on the estimates, calibration weighting from NATSUP will be applied.

A supplementary table comparing characteristics of responders with non-responders among individuals with diabetes will be conducted.

Primary analyses

Cross-tabulations will be conducted to describe habitual physical activity among individuals with diabetes and to display potential subgroup differences.

Table 1 (participant characteristics) will be standardized on age and sex due to expected large differences between “Individuals with diabetes” and “Individuals with no known diabetes”.

Both crude and multivariable adjusted associations will be estimated.

Four multivariable linear regression analyses will be conducted to investigate the association between selected determinants and MVPA. Adjustments will differentiate from each analysis according to DAGs (appendix 1-4). Model assumptions of linear regressions will be assessed:

- 1) Exposure: Comorbidity with and without mental diseases
Adjustments: Age, alcohol consumption, diet, educational level, ethnicity, sex, marital status, obesity, and smoking
- 2) Exposure: Obesity
Adjustments: **Age, alcohol consumption, diet, educational level, ethnicity, sex, marital status, comorbidity, and smoking**
- 3) Exposure: Stress
Adjustments: Age, educational level, sex, marital status, obesity, comorbidity, and smoking
- 4) Exposure: Health-related quality of life
Adjustments: Age, educational level, sex, marital status, obesity, comorbidity, smoking, and stress

Secondary analyses

Four multivariable logistic regression analyses will be conducted with the same exposures and adjustments as the abovementioned. Using multivariable logistic regression analyses with the WHO guidelines of physical activity as outcome (binary). Model assumptions of logistic regressions will be examined by linearity between independent variables and log odds of outcome.

Furthermore, four multivariable linear regression analyses will be conducted to investigate the secondary aim with sedentary behavior in leisure time as outcome and comorbidity, obesity, stress, and health-related quality of life as exposures. Adjustments in the secondary analyses will be the same as the primary analyses. Model assumptions will be assessed.

At last, cross-tabulations will be conducted to describe proportions of whether inactive participants with diabetes who are motivated for being more physically active. The cross-tabulations will be conducted in subgroups of age and sex due to expected differences in motivation of behavioral changes.

References

1. Vandenbroucke JP, von Elm E, Altman DG, Gøtzsche PC, Mulrow CD, Pocock SJ, et al. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE): explanation and elaboration. *Int J Surg*. 2014;12(12):1500-24.
2. Willadsen TG, Siersma V, Nicolaisdóttir DR, Køster-Rasmussen R, Jarbøl DE, Reventlow S, et al. Multimorbidity and mortality: A 15-year longitudinal registry-based nationwide Danish population study. *Journal of Comorbidity*. 2018;8(1):2235042X18804063.
3. Difrancesco S, Lamers F, Riese H, Merikangas KR, Beekman ATF, van Hemert AM, et al. Sleep, circadian rhythm, and physical activity patterns in depressive and anxiety disorders: A 2-week ambulatory assessment study. *Depress Anxiety*. 2019;36(10):975-86.
4. Izawa KP, Watanabe S, Oka K, Hiraki K, Morio Y, Kasahara Y, et al. Association between mental health and physical activity in patients with chronic heart failure. *Disabil Rehabil*. 2014;36(3):250-4.
5. World Health Organization. Obesity: preventing and managing the global epidemic : report of a WHO consultation. Geneva: World Health Organization; 2000.
6. Cohen S, Kamarck T, Mermelstein R. A global measure of perceived stress. *J Health Soc Behav*. 1983;24(4):385-96.
7. Ware J, Jr., Kosinski M, Keller SD. A 12-Item Short-Form Health Survey: construction of scales and preliminary tests of reliability and validity. *Med Care*. 1996;34(3):220-33.
8. Ware J, Jr., Kosinski M, Keller SD. SF-12 : how to score the SF-12 physical and mental health summary scales 1995.
9. WHO guidelines on physical activity and sedentary behaviour. World Health Organization; 2020.
10. Toft U, Kristoffersen LH, Lau C, Borch-Johnsen K, Jørgensen T. The Dietary Quality Score: validation and association with cardiovascular risk factors: the Inter99 study. *European Journal of Clinical Nutrition*. 2007;61(2):270-8.

Supplement 3

Supplementary table 1. Logistic regressions on the association of determinants with adherence to recommendations^a in participants with diabetes

	Crude		Multivariable adjusted	
	POR [95 % CI]	<i>p</i> value	POR [95 % CI]	<i>p</i> value
Comorbidities including mental conditions¹				
No comorbidities	1.00 (Reference)		1.00 (Reference)	
One comorbidity	0.69 [0.55; 0.86]	0.001*	0.84 [0.67; 1.06]	0.133
Two comorbidities	0.53 [0.43; 0.66]	0.001*	0.70 [0.56; 0.88]	0.002*
Three or more comorbidities	0.42 [0.34; 0.51]	0.001*	0.59 [0.47; 0.74]	0.001*
Comorbidities excluding mental conditions¹				
No comorbidities	1.00 (Reference)		1.00 (Reference)	
One comorbidity	0.76 [0.62; 0.94]	0.011*	0.94 [0.75; 1.17]	0.573
Two comorbidities	0.57 [0.47; 0.70]	0.001*	0.76 [0.61; 0.94]	0.011*
Three or more comorbidities	0.46 [0.37; 0.56]	0.001*	0.65 [0.52; 0.81]	0.001*
Obesity²				
Underweight /Normal	1.00 (Reference)		1.00 (Reference)	
Overweight	0.74 [0.63; 0.88]	0.001*	0.75 [0.63; 0.89]	0.001*
Obese class I	0.58 [0.49; 0.69]	0.001*	0.62 [0.52; 0.75]	0.001*
Obese class II	0.45 [0.36; 0.56]	0.001*	0.48 [0.38; 0.61]	0.001*
Obese class III	0.33 [0.25; 0.43]	0.001*	0.36 [0.27; 0.48]	0.001*
Stress³				
Low perceived stress	1.00 (Reference)		1.00 (Reference)	
Moderate perceived stress	0.65 [0.57; 0.73]	0.001*	0.71 [0.62; 0.80]	0.001*
High perceived stress	0.39 [0.29; 0.52]	0.001*	0.44 [0.32; 0.59]	0.001*
HRQoL⁴				
<i>Physical score</i>				
Moderate to high physical HRQoL	1.00 (Reference)		1.00 (Reference)	
Low physical HRQoL	0.40 [0.35; 0.46]	0.001*	0.56 [0.48; 0.65]	0.001*
<i>Mental score</i>				
Moderate to high mental HRQoL	1.00 (Reference)		1.00 (Reference)	
Low mental HRQoL	0.54 [0.48; 0.62]	0.001*	0.69 [0.59; 0.82]	0.001*

n = 6,341

^aFollowing WHO recommendations = at least 150 mins/week of moderate physical activity or at least 75 mins/week of vigorous physical activity or an appropriate combination hereby

POR = prevalence odds ratio; CI = confidence interval

Crude and multivariable adjusted prevalence odds ratios of following WHO recommendations of physical activity in relation to comorbidities, obesity, stress, and health-related quality of life. Significant results (*p*<0.05) are marked with *Crude and adjusted models were weighted for non-response. Each multivariable model was adjusted as follows: ¹Age, sex, ethnicity, educational level, marital status, diet, smoking, alcohol consumption, BMI. ²Age, sex, ethnicity, educational level, marital status, diet, smoking, alcohol consumption, comorbidity.³Age, sex, educational level, marital status, smoking, BMI, comorbidity. ⁴Age, sex, educational level, marital status, smoking, BMI, comorbidity, stress.

Supplement 4

Supplementary table 2. Linear regression on the associations of determinants with total sedentary behavior in participants with diabetes

	Total sedentary behavior h/day			
	Crude β^a [95 % CI]	<i>p</i> value	Multivariable adjusted β [95 % CI]	<i>p</i> value
Comorbidities including mental conditions¹				
No comorbidities	(Reference)		(Reference)	
One comorbidity	-0.30 [-0.75; 0.15]	0.187	-0.01 [-0.45; 0.43]	0.973
Two comorbidities	-0.46 [-0.89; -0.02]	0.038*	-0.0004 [-0.43; 0.43]	0.998
Three or more comorbidities	-0.09 [-0.53; 0.35]	0.691	0.33 [-0.11; 0.77]	0.139
Comorbidities excluding mental conditions¹				
No comorbidities	(Reference)		(Reference)	
One comorbidity	-0.47 [-0.90; -0.03]	0.037*	-0.12 [-0.55; 0.31]	0.577
Two comorbidities	-0.50 [-0.92; -0.08]	0.020*	-0.004 [-0.42; 0.41]	0.984
Three or more comorbidities	-0.29 [-0.72; 0.15]	0.197	0.23 [-0.23; 0.65]	0.316
Obesity²				
Underweight /Normal	(Reference)		(Reference)	
Overweight	-0.16 [-0.51; 0.18]	0.352	0.02 [-0.32; 0.35]	0.920
Obese class I	-0.02 [-0.39; 0.35]	0.898	0.15 [-0.22; 0.51]	0.427
Obese class II	0.66 [0.16; 1.15]	0.009*	0.63 [0.14; 1.12]	0.011*
Obese class III	1.15 [0.49; 1.80]	0.001*	1.05 [0.39; 1.71]	0.002*
Stress³				
Low perceived stress	(Reference)		(Reference)	
Moderate perceived stress	0.15 [-0.11; 0.41]	0.270	-0.01 [-0.28; 0.25]	0.923
High perceived stress	0.92 [0.22; 1.62]	0.010*	0.55 [-0.14; 1.24]	0.116
HRQoL⁴				
<i>Physical score</i>				
Normal physical HRQoL	(Reference)		(Reference)	
Low physical HRQoL	-0.09 [-0.36; 0.18]	0.529	0.12 [-0.18; 0.41]	0.438
<i>Mental score</i>				
Normal mental HRQoL	(Reference)		(Reference)	
Low mental HRQoL	0.42 [0.12; 0.71]	0.006*	0.14 [-0.21; 0.49]	0.426

n = 5,967

CI = confidence interval

^a β coefficients and 95 % confidence intervals (95 % CI) represent mean difference in sedentary behavior h/day compared with the reference. Significant results (p<0.05) are marked with *Crude and adjusted models were weighted for non-response. Each multivariable model was adjusted as follows: ¹Age, sex, ethnicity, educational level, marital status, diet, smoking, alcohol consumption, BMI. ²Age, sex, ethnicity, educational level, marital status, diet, smoking, alcohol consumption, comorbidity. ³Age, sex, educational level, marital status, smoking, BMI, comorbidity. ⁴Age, sex, educational level, marital status, smoking, BMI, comorbidity, stress.

Supplement 5

Supplementary table 3. Distribution of sufficiently physically active participants and their motivation for being more physically active

	Motivated	Not motivated	Don't know
n (%)	2,681 (55.6)	1,341 (27.3)	798 (17.3)
Sex			
Male	1,544 (53.0)	902 (30.1)	478 (16.9)
Female	1,137 (59.5)	439 (22.9)	320 (17.6)
Age (categories)			
16-34 years	127 (64.2)	40 (22.6)	22 (13.2)
35-54 years	607 (69.5)	153 (17.5)	92 (13.0)
55-64 years	785 (61.5)	278 (22.0)	191 (16.5)
65-74 years	851 (47.2)	576 (33.1)	343 (19.7)
≥75 years	311 (41.4)	294 (38.6)	150 (20.1)
Obesity			
Underweight /Normal	531 (47.3)	437 (36.2)	185 (16.5)
Overweight	1,034 (53.3)	573 (29.0)	320 (17.7)
Obese class I	678 (60.6)	239 (21.6)	192 (17.8)
Obese class II	276 (69.1)	58 (14.5)	66 (16.4)
Obese class III	144 (80.2)	17 (10.5)	16 (9.3)
Comorbidities including mental conditions			
No comorbidities	364 (54.0)	208 (29.2)	114 (16.8)
One comorbidity	752 (53.6)	434 (31.0)	208 (15.4)
Two comorbidities	771 (55.4)	381 (27.7)	226 (16.9)
Three or more comorbidities	794 (58.4)	318 (22.4)	250 (19.2)
Stress			
Low perceived stress	1,382 (50.7)	853 (32.7)	415 (16.6)
Moderate perceived stress	1,058 (60.5)	382 (21.3)	319 (18.3)
High perceived stress	105 (72.5)	23 (15.2)	17 (12.3)
HRQoL			
<i>Physical score</i>			
Normal physical HRQoL	1,822 (52.3)	1,043 (30.5)	564 (17.2)
Low physical HRQoL	636 (66.3)	182 (17.9)	150 (15.8)
<i>Mental score</i>			
Normal mental HRQoL	850 (52.3)	564 (34.1)	233 (23.6)
Low mental HRQoL	1,608 (57.9)	661 (23.6)	481 (18.6)

n = 4,820 (n differs due to variations in complete responses in each variable)

Data are presented as number (%)

Abbreviations: HRQoL = Health-related quality of life.