

Food insecurity, diet quality, and suboptimal diabetes management among US adults with diabetes

Sarah S Casagrande ¹, Kai McKeever Bullard,² Karen R Siegel,² Jean M Lawrence ³

To cite: Casagrande SS, Bullard KMCK, Siegel KR, *et al.* Food insecurity, diet quality, and suboptimal diabetes management among US adults with diabetes. *BMJ Open Diab Res Care* 2022;**10**:e003033. doi:10.1136/bmjdr-2022-003033

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjdr-2022-003033>).

Parts of this study were presented in abstract form at the 82nd Scientific Sessions of the American Diabetes Association, New Orleans, Louisiana, 3–7 June 2022.

Received 7 July 2022
Accepted 8 October 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Sarah S Casagrande;
scasagrande@s-3.com

ABSTRACT

Introduction A healthy diet is recommended to support diabetes management, including HbA1c, blood pressure, and cholesterol (ABC) control, but food insecurity is a barrier to consuming a healthy diet. We determined the prevalence of food insecurity and diet quality among US adults with diabetes and the associations with ABC management.

Research design and methods Cross-sectional analyses were conducted among 2075 adults ≥20 years with diagnosed diabetes who participated in the 2013–2018 National Health and Nutrition Examination Surveys. Food insecurity was assessed using a standard questionnaire and diet quality was assessed using quartiles of the 2015 Healthy Eating Index. Adjusted ORs (aOR, 95% CI) were calculated from logistic regression models to determine the association between household food insecurity/diet quality and the ABCs while controlling for sociodemographic characteristics, healthcare utilization, smoking, medication for diabetes, blood pressure, or cholesterol, and body mass index.

Results Overall, 17.6% of adults had food insecurity/low diet quality; 14.2% had food insecurity/high diet quality; 33.1% had food security/low diet quality; and 35.2% had food security/high diet quality. Compared with adults with food security/high diet quality, those with food insecurity/low diet quality were significantly more likely to have HbA1c ≥7.0% (aOR=1.85, 95% CI 1.23 to 2.80) and HbA1c ≥8.0% (aOR=1.79, 95% CI 1.04 to 3.08); food insecurity/high diet quality was significantly associated with elevated HbA1c; and food security/low diet quality with elevated A1c.

Conclusions Food insecurity, regardless of diet quality, was significantly associated with elevated A1c. For people with food insecurity, providing resources to reduce food insecurity could strengthen the overall approach to optimal diabetes management.

INTRODUCTION

The American Diabetes Association (ADA) recommends that individuals with diabetes eat a healthy diet that includes a variety of nutrient-dense foods such as fruits, vegetables, and whole grains in appropriate portion sizes and is limited in added sugars and fats.¹ Consuming a healthy diet may help people with diabetes maintain or achieve a desired

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ While a healthy diet is important for persons with diabetes to manage their disease, food insecurity may negatively impact healthy eating. Few national studies have assessed the combination of food insecurity and diet quality on the management of diabetes.

WHAT THIS STUDY ADDS

⇒ Among a national sample of US adults with diabetes, food insecurity, regardless of diet quality, was significantly associated with elevated A1c.
⇒ Low diet quality, even in the presence of food security, was significantly associated with elevated A1c.
⇒ There were no statistically significant associations between food insecurity and low diet quality with high blood pressure or elevated low-density lipoprotein cholesterol.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ Identifying persons with food insecurity in healthcare settings and referring them to interventions and resources to reduce food insecurity could strengthen the overall approach to optimal diabetes management.

body weight; achieve individualized HbA1c, blood pressure (BP), and cholesterol (ABCs) targets; and delay or prevent comorbidities and diabetes-related complications.² However, there are barriers to consuming a healthy diet, including adequate disposable income to purchase food, access to healthy food, education on preparing and cooking healthy meals, and healthy eating support from friends and family.^{3–5}

Food security, a state of having reliable access to sufficient quantity of affordable and nutritious food, is one factor that may affect healthy eating among people with diabetes.⁶ A report based on the 1999–2002 National Health and Nutrition Examination Surveys (NHANES) demonstrated that diabetes prevalence was higher among people with severe food insecurity than people without food

insecurity; the association remained after adjusting for body mass index (BMI).⁷ Another NHANES study showed that while diet quality trends among adults with diabetes improved overall between 1999 and 2014, there were substantial disparities in consuming a high-quality diet; adults with low education, low income, and food insecurity were less likely to consume a healthy diet and showed no improvement in diet quality over time.⁸ Further, results from the 1999–2008 NHANES data showed that food insecurity among adults with diabetes was significantly associated with HbA1c $\geq 9.0\%$ and low-density lipoprotein cholesterol (LDL-C) ≥ 100 mg/dL after controlling for several sociodemographic and diabetes-related factors.⁹ A prospective cohort study of adult patients with diabetes in a primary care network also found that food insecurity was associated with higher HbA1c levels.¹⁰ A 2011–2016 NHANES study among adults with diabetes (diagnosed and undiagnosed) found an interaction between diet quality and food insecurity on the association with HbA1c at 8.0% to $<9.0\%$.¹¹

Given the gaps in the literature, the objective of this study was to determine the prevalence of combined food insecurity and healthy diet quality status among US adults with diabetes, overall and stratified by sociodemographic characteristics and health behaviors. In addition, we examined how food insecurity/diet quality status was associated with ABC management while accounting for factors related to achieving these ABC targets.

RESEARCH DESIGN AND METHODS

Study population

The NHANES is a stratified multistage probability cluster survey conducted in the non-institutionalized civilian US population.¹² Participants were interviewed in their homes to obtain demographic and health information and then visited a mobile examination center (MEC) for physical examinations and laboratory measures.^{13 14}

Data were analyzed from three cycles of NHANES (2013–2014, 2015–2016, 2017–2018). Unweighted response rates ranged from 51.9% to 71.0% for the interviewed sample and from 48.8% to 68.5% for the examination sample. Study participants selected for inclusion in these analyses which combined the data from the three NHANES cycles included 2075 adults aged ≥ 20 years with diagnosed diabetes and at least one reliable 24-hour dietary recall.

Measures

Diagnosed diabetes was considered present if study participants answered ‘yes’ when asked if a physician or other healthcare professional ever told them that they had diabetes or sugar diabetes. Adults with HbA1c indicative of diabetes without a physician diagnosis of diabetes were not included in these analyses since this study focuses on diabetes management of the ABCs. Participants self-reported sociodemographic characteristics (current age, race and ethnicity, household income, and household

size, to determine poverty income ratio (PIR), with PIR <1.0 indicating below the household poverty threshold), highest education level, receipt of Supplemental Nutrition Assistance Program (SNAP) benefits, health insurance status and type of health insurance, healthcare utilization (seen a physician or diabetes specialist in the past year, HbA1c checked in the past year), smoking status (current, former, or never), and diabetes, BP, and cholesterol medication use. Height and weight were measured by a trained interviewer to determine the BMI (kg/m^2), defined as: normal <25.0 kg/m^2 , overweight 25.0–29.9 kg/m^2 ; obesity ≥ 30 kg/m^2 .

Main independent measures: food insecurity and diet quality

To assess the food insecurity status for adults in their household, participants responded to either the first 10 items (if no children in the household) or full 18 items (if children in the household) of the US Department of Agriculture’s household food security questionnaire, which assessed running out of food and not being able to afford more, cutting meal size, skipping meals, eating less or not at all, or going hungry due to lack of money for food in the past 12 months.^{15 16} Participants were categorized as having full food security, marginal food security, or low/very low food security based on the frequency of responses to the questionnaire items and the standard scoring methods.¹⁷

Diet quality was determined by the Healthy Eating Index 2015 (HEI-2015),¹⁸ which measures how well an individual’s diet aligns with federal dietary guidelines and is generally considered a good marker of overall diet quality. The HEI-2015 includes 13 food group components that reflect the key recommendations in the 2015–2020 Dietary Guidelines for Americans—nine ‘adequacy’ components represent food groups whose consumption is encouraged (total fruits, whole fruits, total vegetables, greens and beans, whole grains, total protein foods, seafood and plant proteins dairy, fatty acids) and four ‘moderation’ components to represent food groups for which there are recommended limits to consumption (refined grains, sodium, added sugars, saturated fats). The composite HEI-2015 scores range from 0 to 100, with higher scores denoting better alignment with dietary recommendations (ie, diet quality).¹⁸ In NHANES, HEI-2015 scores were determined using participant data from at least one reliable 24-hour dietary recall and the simple HEI scoring algorithm-per person method.^{19 20} Participants’ diet quality was classified into four categories based on HEI-2015 score quartiles, with the first quartile indicating low diet quality, second quartile indicating low-moderate diet quality, third quartile indicating moderate-high diet quality, and the fourth quartile indicating high diet quality.

A combined food insecurity/diet quality variable was defined as (1) food secure (full food security) and moderate-high diet quality (third and fourth HEI-2015 quartiles, referred hereafter as ‘high diet quality’), (2) food secure and low-moderate diet quality (first and

second HEI-2015 quartiles, referred hereafter as 'low diet quality'), (3) food insecure (low/very low or marginal food security) and high diet quality, and (4) food insecure and low diet quality. In a sensitivity analysis, we assessed a combined food insecurity/diet quality variable that defined high diet quality as only the fourth quartile of the HEI-2015 and low diet quality as the first to third quartiles of the HEI-2015; categorization of food insecurity remained the same.

Main dependent (outcome) variables: ABCs

A phlebotomist obtained a blood sample from all participants during the MEC visit using a standardized protocol.¹⁴ BP was measured using a standardized mercury sphygmomanometer after the participant rested quietly for 5 min.¹³ Up to four readings were taken, and the readings were averaged, excluding the first measure. HbA1c was directly measured.¹⁴ LDL-C levels were calculated for people who had fasted ≥ 8 to < 24 hours using the Friedewald formula, which is not valid when triglycerides are ≥ 400 mg/dL.²¹ Outcomes for the ABC measures were based on clinically elevated values. The cut-points for elevated HbA1c ($\geq 7.0\%$, $\geq 8.0\%$, $\geq 9.0\%$), hypertension (BP $\geq 130/80$ mm Hg, BP $\geq 140/90$ mm Hg), and cholesterol (LDL-C ≥ 100 mg/dL, LDL-C ≥ 130 mg/dL) were based on current guidelines from the ADA Standards of Care.²²

Statistical analysis

We used descriptive statistics to describe the characteristics of the study population overall and by food insecurity and diet quality categories. Bivariate associations between food insecurity and diet quality categories and demographic characteristics and health behaviors were tested using a χ^2 test for significance. Logistic regression was used to calculate ORs (95% CI) for the association between (1) food insecurity, (2) diet quality, and (3) food insecurity/diet quality (four-level combined variable) and elevated ABC levels. Models were (1) unadjusted, (2) adjusted for age, sex, race, and ethnicity, (3) additionally adjusted for PIR, education, health insurance, (4) additionally adjusted for healthcare utilization (seeing a physician or specialist in the past year, having HbA1c checked), (5) additionally adjusted for smoking status, diabetes, BP, and cholesterol medications, and BMI, and (6) additionally adjusted for food insecurity or diet quality when these variables were not the main covariates. P values < 0.05 were considered statistically significant. All statistical analyses used sampling weights to account for the NHANES' complex survey design using SUDAAN (SUDAAN User's Manual, Release 11, 2012; Research Triangle Institute). A 6-year dietary weight was used for all analyses except for analyses among the fasting sample (LDL-C and triglycerides) where a 6-year fasting weight was used.

RESULTS

Participant characteristics

Among US adults with diagnosed diabetes, the mean age was 60.4 years, 53.0% were men, 15.3% were living below the poverty threshold, and 19.3% had low/very low food security (food insecure) (online supplemental appendix table 1). Mean HEI score was 53.6, with 66.7% having an HEI score < 60 . The mean HEI-2015 scores were 36.5 (range, 10.0–43.2) for the first quartile, 48.3 (range, 43.2–52.7) for the second quartile, 57.9 (range, 52.7–62.9) for the third quartile, and 71.8 (range, 62.8–99.5) for the fourth quartile. Nearly half (49.8%) had HbA1c $\geq 7.0\%$, 28.5% had BP $\geq 140/90$ mm Hg, and 18.0% had LDL-C ≥ 130 mg/dL. Several sociodemographic characteristics and metabolic control varied by food insecurity and diet quality.

Participant characteristics by food insecurity and diet quality

When food insecurity and diet quality were combined into a four-level variable, 17.6% of persons with diabetes had food insecurity (low/very low or marginal)/low diet quality (first/second HEI-2015 quartile); 14.2% had food insecurity/high diet quality (third/fourth HEI-2015 quartile); 33.1% had food security/low diet quality; and 35.2% had food security/high diet quality (table 1). Among adults with diabetes, the distribution of age, sex, race and ethnicity, PIR, education, receiving SNAP benefits, and health insurance status all varied by food insecurity/diet quality category ($p < 0.01$ for all). In addition, the per cent with uncontrolled HbA1c, elevated LDL-C, and having any one elevated ABC measure varied by food insecurity/diet quality category ($p < 0.03$ for all) (figure 1).

Adjusted associations between food insecurity and ABCs

Adults with diabetes who were food insecure were significantly more likely to have elevated HbA1c compared with those with full food security after adjusting for sociodemographic characteristics, healthcare utilization, smoking status, diabetes, BP, and cholesterol medication use, BMI, and diet quality (adjusted OR (aOR)=1.62, 95% CI 1.07 to 2.47 for HbA1c $\geq 7.0\%$; aOR=1.71, 95% CI 1.14 to 2.56 for HbA1c $\geq 8.0\%$; aOR=1.59, 95% CI 1.04 to 2.43 for HbA1c $\geq 9.0\%$) (online supplemental appendix table 2). There were no significant associations between food insecurity and elevated BP and LDL-C except for the unadjusted association between food insecurity and LDL-C ≥ 100 mg/dL (OR=1.50, 95% CI 1.05 to 2.15).

Adjusted associations between diet quality and ABCs

Adults with diabetes who had low diet quality (lowest, first quartile) were significantly more likely to have elevated HbA1c (aOR=1.61, 95% CI 1.06 to 2.46 for $\geq 7.0\%$; aOR=1.60, 95% CI 1.01 to 2.53 for $\geq 8.0\%$) compared with those with high diet quality (highest, fourth quartile) after full adjustment (online supplemental appendix table 2). There were no significant associations between food insecurity and elevated BP and LDL-C.

Table 1 Characteristics of US adults with diagnosed diabetes by food insecurity and diet quality

	Food insecurity and diet quality (HEI-2015)*				P value, χ^2
	Food secure and high diet quality	Food secure and low diet quality	Food insecure and high diet quality	Food insecure and low diet quality	
Unweighted n	681	551	400	443	
	% (SE)				
Total	35.2 (1.5)	33.1 (1.5)	14.2 (1.1)	17.6 (1.3)	
Age (years)					
Mean	62.5 (0.7)	60.3 (0.7)	59.9 (0.94)	56.3 (0.8)	
20–44	6.9 (1.4)	13.0 (1.3)	15.6 (2.5)	19.9 (1.9)	0.001
45–64	45.1 (2.7)	48.2 (2.5)	43.3 (3.6)	50.5 (3.7)	
≥65	48.0 (2.8)	38.8 (2.6)	41.1 (4.0)	29.7 (3.5)	
Sex					
Men	58.1 (2.8)	57.5 (3.3)	36.8 (3.3)	49.3 (2.6)	<0.001
Women	41.9 (2.8)	42.5 (3.3)	63.2 (3.3)	50.7 (2.6)	
Race and ethnicity					
Non-Hispanic White	67.4 (3.0)	74.1 (2.5)	45.8 (4.7)	59.0 (3.8)	<0.001
Non-Hispanic Black	12.7 (1.9)	11.0 (1.3)	19.2 (2.9)	16.7 (2.4)	
Hispanic/Latino	11.3 (1.8)	12.5 (2.2)	29.7 (4.0)	21.6 (2.7)	
Non-Hispanic Asian	8.6 (1.5)	2.4 (0.6)	5.3 (1.6)	2.7 (0.8)	
Poverty income ratio					
<1.00	6.9 (1.3)	8.0 (1.2)	26.5 (2.8)	35.6 (3.6)	<0.001
1.00–2.00	13.8 (1.9)	16.8 (1.9)	35.0 (3.2)	29.5 (3.5)	
>2.00	72.5 (2.9)	69.8 (2.3)	31.0 (3.4)	29.6 (3.7)	
Missing	6.8 (1.4)	5.5 (0.7)	7.5 (1.5)	5.3 (1.7)	
Education					
Less than high school	12.4 (1.7)	16.1 (1.7)	31.3 (2.9)	27.3 (3.1)	<0.001
High school graduate	23.0 (2.2)	23.6 (2.6)	32.5 (3.0)	30.4 (3.2)	
Some college	33.4 (2.9)	37.2 (3.1)	26.2 (2.7)	35.7 (3.2)	
College graduate or above	31.2 (3.3)	23.1 (2.3)	10.1 (1.6)	6.6 (1.9)	
SNAP benefit					
Yes	9.3 (1.6)	13.2 (1.7)	34.1 (3.3)	43.9 (3.5)	<0.001
No	90.7 (1.6)	86.8 (1.7)	65.9 (3.3)	56.1 (3.5)	
Health insurance					
Uninsured	4.8 (0.8)	7.7 (1.5)	12.5 (1.9)	12.9 (2.2)	0.002
Insured	95.2 (0.8)	92.3 (1.5)	87.5 (1.9)	87.1 (2.2)	
Health insurance type					
Private	65.3 (3.6)	68.3 (3.1)	53.6 (4.0)	42.7 (3.3)	<0.001
Medicare	48.2 (2.7)	42.8 (2.7)	48.6 (4.4)	46.8 (3.4)	0.599
Medicaid	7.7 (1.7)	9.1 (1.3)	22.7 (2.8)	27.0 (2.8)	<0.001
Military/other government	16.4 (2.1)	13.5 (2.4)	12.4 (2.4)	17.3 (2.5)	0.409
Seen physician in the past year					
Yes	98.1 (0.8)	97.6 (0.9)	94.0 (1.5)	95.2 (1.1)	0.124
No	1.9 (0.8)	2.4 (0.9)	6.0 (1.5)	4.8 (1.1)	

Continued

Table 1 Continued

	Food insecurity and diet quality (HEI-2015)*				P value, χ^2
	Food secure and high diet quality	Food secure and low diet quality	Food insecure and high diet quality	Food insecure and low diet quality	
Seen diabetes specialist in the past year					
Yes	33.3 (2.9)	24.3 (2.4)	34.8 (2.7)	26.6 (2.9)	0.021
No	35.6 (3.2)	35.5 (2.4)	28.3 (2.5)	36.2 (3.3)	
Never	31.1 (2.4)	40.2 (2.6)	36.9 (2.4)	37.2 (3.6)	
HbA1c checked in the past year					
Yes	89.7 (1.2)	81.2 (2.5)	78.2 (2.5)	74.9 (2.9)	0.001
No	7.4 (1.0)	13.1 (2.1)	18.0 (2.1)	19.7 (2.8)	
Do not know	2.9 (0.7)	5.7 (1.5)	3.8 (1.1)	5.4 (1.3)	
Smoking status					
Current	5.9 (1.4)	15.1 (2.2)	13.6 (2.2)	25.6 (3.1)	<0.001
Former	43.8 (3.3)	35.2 (2.8)	33.7 (2.7)	32.8 (2.9)	
Never	50.3 (3.0)	49.7 (3.3)	52.7 (3.2)	41.6 (3.1)	
Diabetes medication					
Insulin only	11.7 (1.8)	9.0 (1.7)	9.3 (1.7)	18.7 (2.7)	0.003
Oral medication only	61.4 (2.7)	58.1 (3.3)	54.1 (2.6)	42.8 (3.0)	
Both insulin and oral	11.2 (1.8)	15.5 (2.3)	18.8 (2.6)	18.4 (1.9)	
None	15.8 (2.1)	17.4 (2.7)	17.9 (2.1)	20.2 (2.3)	
Taking antihypertensive medication	65.7 (2.7)	63.5 (3.0)	57.4 (3.3)	60.8 (2.7)	0.094
Taking antilipidemic medication	61.3 (2.9)	64.9 (2.2)	50.1 (3.4)	56.3 (2.7)	0.007
BMI (kg/m ²)					
Mean	32.0 (0.4)	34.0 (0.5)	33.1 (0.6)	34.8 (0.6)	
<25.0	11.7 (1.7)	8.5 (1.4)	10.4 (1.7)	8.0 (1.5)	0.027
25.0–29.9	30.1 (2.8)	23.2 (2.1)	32.1 (2.6)	21.1 (2.7)	
≥30	58.2 (3.3)	68.3 (1.7)	57.5 (3.0)	70.9 (3.1)	

Boldface p values indicate statistical significance of $p < 0.05$.
 *Food insecurity is defined as marginal or low/very low food security; food secure is defined as full food security; low diet quality is defined as HEI-2015 score in the 1st or 2nd quartile; high diet quality is defined as HEI-2015 score in the 3rd or 4th quartile.
 BMI, body mass index; HEI-2015, Healthy Eating Index 2015; SNAP, Supplemental Nutrition Assistance Program.

Adjusted joint associations between food insecurity, diet quality, and ABCs

Adults with diabetes who had food insecurity (low/very low or marginal food security)/low diet quality (first/second HEI-2015 quartile) were significantly more likely to have HbA1c $\geq 7.0\%$ (aOR=1.85, 95% CI 1.23 to 2.80) and HbA1c $\geq 8.0\%$ (aOR=1.79, 95% CI 1.04 to 3.08) compared with those who had food security (full food security)/high diet quality (third/fourth HEI-2015 quartile) in fully adjusted models (table 2). A similar association for HbA1c $\geq 7.0\%$ and HbA1c $\geq 8.0\%$ was shown for those who had food insecurity/high diet quality versus food security/high diet quality. Adults who had food security/low diet quality were significantly more likely to have HbA1c $\geq 7.0\%$ (aOR=1.55, 95% CI 1.07 to 2.24).

In unadjusted analysis, adults with diabetes who had food insecurity/low diet quality were significantly more likely to have A1c $\geq 9.0\%$ compared with those who had food security/high diet quality (OR=2.10, 95% CI 1.24 to 3.55), but this association became non-significant after adjusting for age, sex, and race/ethnicity; a similar finding was shown for those with food insecurity/high diet quality, but the association became non-significant after adjusting for all sociodemographic characteristics. There were no significant interactions between food insecurity and diet quality for any of the other ABC outcomes.

When high diet quality was defined as only the fourth quartile of the HEI-2015 (vs low/moderate diet quality, first to third HEI-2015 quartiles), the results remained largely unchanged for those with food insecurity and low/




Figure 1 Prevalence of suboptimal diabetes ABC management by food insecurity and diet quality. Error bars represent 95% CIs. ABC, HbA1c, blood pressure, and cholesterol; BP, blood pressure; LDL-C, low-density lipoprotein cholesterol.

moderate diet quality (online supplemental appendix table 3). In fully adjusted models, adults who had food insecurity and low/moderate diet quality (first to third HEI-2015 quartiles) were significantly more likely to have elevated HbA1c compared with those who had food security/high diet quality (fourth HEI-2015 quartile).

DISCUSSION

In this national study of US adults with diabetes, the combination of food insecurity and low diet quality was significantly associated with poorer glycemic management compared with those who were food secure and had high diet quality, independent of sociodemographic characteristics, healthcare utilization, diabetes medications, and BMI; findings were similar for those with food insecurity and high diet quality. In addition, low diet quality, even in the presence of food security, was significantly associated with A1c $\geq 7.0\%$. We did not find any statistically significant associations between food insecurity and low diet quality and high BP or LDL-C.

Lower socioeconomic status, which, by definition, is strongly associated with food insecurity, has been previously associated with poorer ABC control.^{23 24} Indeed, in the current study among adults with food insecurity, living below the poverty threshold or having Medicaid was five times more prevalent compared with those adults with full food security; having less than a high school education was nearly 2.5 times more prevalent among those with food insecurity compared with those adults with full food security. Diet quality was also worse for those with less income, but the difference was not as large in magnitude as it was for food insecurity. Previous studies have found that diet quality is related to the affordability of healthy foods,^{25–27} but access to healthy food and the ability to afford these foods does not directly equate to a high-quality diet; diet quality is a modifiable behavior, more so than food insecurity. However, it is well

established that behavior change is difficult and challenging to maintain.²⁸

Previous studies in various populations have shown that food insecurity is associated with poorer glycemic control. Among low-income urban residents with type 2 diabetes, food insecurity was associated with poorer glucose control; however, among those who were food insecure, the receipt of SNAP benefits was associated with a significantly reduced odds of having poor glucose control.²⁹ Thus, economic food assistance appears helpful for those with diabetes; these data suggest that food insecure adults were eating healthier with SNAP than those without assistance. SNAP assistance has also been shown to reduce psychological distress related to food insecurity.³⁰ In a different study of low-income patients with diabetes, those who were food insecure had higher mean HbA1c levels compared with those who were food secure.³¹ A previous study using NHANES 2011–2016 data among adults with diagnosed and undiagnosed diabetes found an interaction between food insecurity and diet quality with elevated HbA1c (8.0% to $<9.0\%$) but the relationship between food insecurity and diet quality was not fully elucidated.¹¹ In another earlier study using NHANES data from 1999 to 2008, food insecurity was associated with HbA1c $>9.0\%$ and poor LDL control after adjustment for sociodemographic characteristics and diabetes-related factors; however, dietary intake was not included in that study.⁹ The results were similar to those from our study, where we found that food insecurity was associated with all categories of poorer glucose control after adjusting for sociodemographic characteristics, diabetes-related factors, including prescription medications, and diet quality. Further, we also found that food insecurity, regardless of whether diet quality was high or low, was associated with elevated HbA1c before and after adjustment for other covariates. Given that diet quality did not account for the association between food insecurity and

Table 2 OR (95% CI) for suboptimal diabetes ABC management by food security and diet quality

	Prevalence, % (SE)	Food security and diet quality (HEI-2015)*			
		Food secure and high diet quality	Food secure and low diet quality	Food insecure and high diet quality	Food insecure and low diet quality
HbA1c ≥7.0%	49.8 (1.8)				
Unadjusted		1.00	1.49 (1.08 to 2.07)	1.51 (1.06 to 2.14)	1.73 (1.26 to 2.36)
Model 1		1.00	1.50 (1.08 to 2.09)	1.59 (1.08 to 2.34)	1.76 (1.25 to 2.47)
Model 2		1.00	1.52 (1.09 to 2.12)	1.64 (1.11 to 2.42)	1.89 (1.34 to 2.66)
Model 3		1.00	1.55 (1.10 to 2.18)	1.64 (1.11 to 2.44)	1.92 (1.36 to 2.71)
Model 4		1.00	1.55 (1.07 to 2.24)	1.69 (1.14 to 2.51)	1.85 (1.23 to 2.80)
HbA1c ≥8.0%	26.1 (1.3)				
Unadjusted		1.00	1.39 (0.93 to 2.07)	2.10 (1.37 to 2.32)	1.97 (1.30 to 2.99)
Model 1		1.00	1.29 (0.86 to 1.94)	1.80 (1.15 to 2.82)	1.67 (1.07 to 2.59)
Model 2		1.00	1.28 (0.85 to 1.93)	1.74 (1.08 to 2.79)	1.61 (0.99 to 2.61)
Model 3		1.00	1.30 (0.87 to 1.95)	1.71 (1.06 to 2.75)	1.62 (1.00 to 2.65)
Model 4		1.00	1.37 (0.91 to 2.07)	1.83 (1.12 to 2.98)	1.79 (1.04 to 3.08)
HbA1c ≥9.0%	14.4 (1.0)				
Unadjusted		1.00	1.27 (0.75 to 2.16)	2.22 (1.40 to 3.52)	2.10 (1.24 to 3.55)
Model 1		1.00	1.11 (0.65 to 1.88)	1.70 (1.07 to 2.71)	1.58 (0.90 to 2.77)
Model 2		1.00	1.10 (0.66 to 1.84)	1.56 (0.94 to 2.60)	1.43 (0.79 to 2.59)
Model 3		1.00	1.11 (0.67 to 1.84)	1.48 (0.88 to 2.47)	1.42 (0.77 to 2.65)
Model 4		1.00	1.18 (0.70 to 1.98)	1.54 (0.92 to 2.60)	1.47 (0.80 to 2.70)
Blood pressure ≥140/90 mm Hg	28.5 (1.5)				
Unadjusted		1.00	1.10 (0.78 to 1.56)	1.12 (0.78 to 1.62)	1.13 (0.80 to 1.59)
Model 1		1.00	1.18 (0.82 to 1.71)	1.18 (0.80 to 1.74)	1.22 (0.84 to 1.79)
Model 2		1.00	1.16 (0.80 to 1.67)	1.11 (0.75 to 1.65)	1.16 (0.77 to 1.74)
Model 3		1.00	1.15 (0.80 to 1.65)	1.12 (0.75 to 1.67)	1.16 (0.78 to 1.74)
Model 4		1.00	1.21 (0.83 to 1.77)	1.14 (0.76 to 1.70)	1.14 (0.74 to 1.75)
Blood pressure ≥130/80 mm Hg	52.0 (1.5)				
Unadjusted		1.00	1.11 (0.81 to 1.53)	1.17 (0.76 to 1.83)	0.97 (0.70 to 1.35)
Model 1		1.00	1.15 (0.82 to 1.62)	1.32 (0.86 to 2.04)	1.01 (0.71 to 1.45)
Model 2		1.00	1.14 (0.80 to 1.61)	1.17 (0.73 to 1.88)	0.90 (0.62 to 1.31)
Model 3		1.00	1.15 (0.81 to 1.63)	1.15 (0.72 to 1.83)	0.92 (0.64 to 1.32)
Model 4		1.00	1.17 (0.78 to 1.75)	1.15 (0.74 to 1.79)	0.87 (0.58 to 1.31)
LDL-C ≥100mg/dL	41.8 (2.4)				
Unadjusted		1.00	1.39 (0.82 to 2.36)	1.93 (1.20 to 3.09)	1.89 (1.15 to 3.11)
Model 1		1.00	1.12 (0.64 to 1.97)	1.36 (0.79 to 2.35)	1.40 (0.78 to 2.51)
Model 2		1.00	1.13 (0.64 to 1.99)	1.19 (0.69 to 2.05)	1.25 (0.68 to 2.29)
Model 3		1.00	1.04 (0.59 to 1.82)	1.11 (0.65 to 1.88)	1.25 (0.66 to 2.34)
Model 4		1.00	1.00 (0.62 to 1.62)	0.93 (0.53 to 1.65)	1.08 (0.52 to 2.23)
LDL-C ≥130mg/dL	18.0 (1.7)				
Unadjusted		1.00	1.63 (0.77 to 3.41)	2.69 (1.31 to 5.52)	1.39 (0.71 to 2.74)
Model 1		1.00	1.24 (0.60 to 2.54)	1.82 (0.84 to 3.94)	1.01 (0.50 to 2.05)
Model 2		1.00	1.27 (0.66 to 2.45)	1.70 (0.72 to 4.02)	0.97 (0.42 to 2.25)
Model 3		1.00	1.22 (0.65 to 2.30)	1.65 (0.73 to 3.72)	0.99 (0.44 to 2.22)

Continued

Table 2 Continued

	Prevalence, % (SE)	Food security and diet quality (HEI-2015)*			
		Food secure and high diet quality	Food secure and low diet quality	Food insecure and high diet quality	Food insecure and low diet quality
Model 4		1.00	1.37 (0.76 to 2.48)	1.24 (0.61 to 2.55)	0.77 (0.31 to 1.90)
At least one elevated ABC (HbA1c ≥7.0%, BP ≥130/80 mm Hg, LDL ≥100 mg/dL)	84.3 (2.1)				
Unadjusted		1.00	1.07 (0.48 to 2.39)	2.44 (1.37 to 4.35)	2.90 (1.24 to 6.82)
Model 1		1.00	0.90 (0.39 to 2.09)	1.87 (1.04 to 3.36)	2.00 (0.82 to 4.89)
Model 2		1.00	0.88 (0.38 to 2.03)	1.70 (0.96 to 3.01)	1.87 (0.81 to 4.32)
Model 3		1.00	0.88 (0.37 to 2.05)	1.67 (0.93 to 3.00)	1.86 (0.80 to 4.34)
Model 4		1.00	0.80 (0.34 to 1.90)	1.59 (0.73 to 3.43)	1.98 (0.72 to 5.49)
At least one elevated ABC (HbA1c ≥8.0%, BP ≥140/90 mm Hg, LDL ≥130 mg/dL)	55.0 (2.6)				
Unadjusted		1.00	1.01 (0.57 to 1.80)	2.38 (1.52 to 3.73)	1.19 (0.78 to 1.83)
Model 1		1.00	0.96 (0.52 to 1.76)	1.89 (1.18 to 3.02)	1.08 (0.67 to 1.76)
Model 2		1.00	0.95 (0.52 to 1.72)	1.75 (1.04 to 2.93)	1.03 (0.61 to 1.76)
Model 3		1.00	0.96 (0.53 to 1.72)	1.72 (1.04 to 2.86)	1.05 (0.60 to 1.86)
Model 4		1.00	0.99 (0.55 to 1.79)	1.58 (0.87 to 2.86)	1.05 (0.57 to 1.96)
Model 1: adjusted for age, sex, race/ethnicity.					
Model 2: Model 1 and additionally adjusted for PIR, education, health insurance.					
Model 3: Model 2 and additionally adjusted for healthcare utilization (seeing physicians in the past year, having HbA1c checked).					
Model 4: Model 3 and additionally adjusted for smoking status, diabetes medication, blood pressure medication, cholesterol medication, BMI.					
Boldface indicates statistical significance of p<0.05.					
*Food insecurity is defined as marginal or low/very low food security; food secure is defined as full food security; low diet quality is defined as HEI-2015 score in the 1st or 2nd quartile; high diet quality is defined as HEI-2015 in the 3rd or 4th quartile.					
ABC, HbA1c, blood pressure, and cholesterol; BMI, body mass index; BP, blood pressure; HEI-2015, Healthy Eating Index 2015; LDL, low-density lipoprotein; PIR, poverty income ratio.					

poor glucose control, future research could investigate the intricacies of food insecurity and the pathways by which it is associated with metabolic outcomes. Potential pathways highlighted in a recent scientific review of social determinants of health as they relate to diabetes included nutritional, compensatory, and psychological pathways, each of which warrants further investigation including the intersectionality of these pathways.³²

While food insecurity was significantly associated with the highest level of uncontrolled HbA1c (≥9.0%) after full adjustment including diet quality, when food insecurity and diet quality were combined into the four-category variable, only the unadjusted associations for food insecurity/low diet quality and food insecurity/high diet quality were significant. There may be other unmeasured factors contributing to the lack of significant associations for the highest level of uncontrolled HbA1c. In addition, 14% of

our sample had HbA1c ≥9.0%, thus our sample size was limited to detect an association for this outcome.

Our study suggests that food insecurity may have a direct impact on metabolic outcomes such as HbA1c. Previous research has shown that stress from food insecurity may result in consuming high-carbohydrate, high-sugar, or high-fat foods which may offer temporary feelings of comfort.³³ Securing these palatable foods may be easier in areas with high food insecurity due to a higher prevalence of food swamps, areas with a high density of establishments selling high-calorie fast food and junk food relative to healthier food items.⁵ However, food assistance programs such as food pantries, Women, Infants, and Children program, and SNAP benefits are resources available to those with food insecurity and may support a higher quality diet.^{34–36}

Consuming a high-quality diet, as recommended by the ADA,² is strongly associated with a variety of factors including community-level resources (eg, access to healthy foods), support for a healthy diet from friends and family, and individual-level sociodemographic characteristics (eg, income barriers), behaviors, and habits. At the community level, lower availability of healthy food has been associated with lower quality dietary intake and poorer glycemic control in some studies^{10 37–39}; other studies have been inconclusive.^{40 41} Considering the many contextual factors that influence diet quality is important when developing interventions and programs to support people with diabetes and improve metabolic outcomes.⁴²

We found no significant association between food insecurity/diet quality and BP or LDL-C. Many of the adults in our study had controlled BP and LDL-C, with 63% on antihypertensive medication and 60% on antilipidemic medication. Therefore, there may have been less of an association for food insecurity and diet quality on these outcomes. In addition, the associations between food insecurity/diet quality may be mediated through obesity, a condition that effected nearly two-thirds of the population of adults with diabetes. This may also partially explain the lack of an association for BP and LDL-C. While LDL-C is considered the hallmark measure for cholesterol management among people with diabetes, triglycerides and high-density lipoprotein cholesterol (HDL-C) are also important markers for hyperlipidemia. Of note, in analyses that were not included in the paper, we found that compared with adults with food security/high diet quality, those with food insecurity/low diet quality were statistically significantly more likely to have high triglycerides and low HDL-C; the significant association with high triglycerides was also found for those with food insecurity/high diet quality. When any of these lipid measures are suboptimal, current diabetes management guidelines recommend optimizing glycemic control.⁴³

Limitations

NHANES is a cross-sectional survey, thus causal associations between our exposures of interest, food insecurity and diet quality, and outcomes of ABC management cannot be determined. Trends in associations of food insecurity/diet quality and ABC management outcomes were not assessed. However, declines in A1c control after 2010 may be associated with the notable prevalence of food insecurity and suboptimal dietary intake among adults with diabetes during 2013–2018. Dietary intake was self-reported, which is subject to measurement error similar to other self-reported data⁴⁴; however, these data still provide valuable information on food intake and eating patterns.⁴⁵ Although we adjusted for many sociodemographic factors, residual confounding may have occurred and the true association between food insecurity/low diet quality and poor ABC management may be associated with other unmeasured variables. Additionally, we do not distinguish between type 1 and type 2 diabetes mellitus in this analysis; however, given that the NHANES

is nationally representative, we can assume that 90%–95% of diagnosed diabetes in this study is type 2 diabetes.⁴⁶ In supplemental analysis, we found no interaction between continuous insulin use, an indicator of type 1 diabetes mellitus, and food insecurity/diet quality. However, this study used a nationally representative sample allowing generalization to the US adult non-institutionalized population with diabetes. Multiple clinical outcomes were assessed using standardized measures to characterize diabetes control.

CONCLUSIONS

In this nationally representative study among US adults with diabetes, food insecurity plus low diet quality was associated with higher HbA1c levels even after accounting for sociodemographic characteristics and health behaviors, with food insecurity having a greater impact on metabolic outcomes than low diet quality. Healthcare providers treating adults with diabetes may consider including an assessment of food insecurity as a part of their overall approach to diabetes-related healthcare, particularly for their patients with suboptimal HbA1c and those living in low-income or under-resourced communities. Future research on food insecurity among adults with diabetes might focus on how food insecurity affects metabolic outcomes and address the multiple pathways that may contribute to these outcomes. Healthcare delivery systems could link patients to social or food services as part of clinical care, in addition to point-of-care interventions that promote healthier lifestyles.

Author affiliations

¹Public Health & Scientific Research, DLH Holdings, Atlanta, Georgia, USA

²Division of Diabetes Translation, Centers for Disease Control and Prevention, Atlanta, Georgia, USA

³Division of Diabetes, Endocrinology, and Metabolic Diseases, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, Maryland, USA

Acknowledgements The authors would like to acknowledge Pamela L Thornton for her review and comments on this paper.

Contributors SSC: Conceptualization and design; Acquired, analyzed, and interpreted the data; Drafted the manuscript; Provided critical revision for important intellectual content. KMB and KRS: Interpreted the data; Provided critical revision for important intellectual content. JML: Conceptualization and design; Provided critical revision for important intellectual content. SSC is the guarantor of this work and takes responsibility for the integrity of the data and the accuracy of the analysis.

Funding This study was supported by a contract with the National Institute of Diabetes and Digestive and Kidney Diseases (contract number: 75N94022F00050).

Disclaimer The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention or the National Institutes of Health.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants and was approved by the National Center for Health Statistics (NCHS) Research Ethics Review Board (protocol numbers: 2018-01 and 2011-17). Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available in a public, open access repository. Data for the NHANES are publicly available at <https://www.cdc.gov/nchs/nhanes/>.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Sarah S Casagrande <http://orcid.org/0000-0002-6293-383X>

Jean M Lawrence <http://orcid.org/0000-0002-2480-8480>

REFERENCES

- American Diabetes Association. 5. Facilitating Behavior Change and Well-being to Improve Health Outcomes: *Standards of Medical Care in Diabetes-2021*. *Diabetes Care* 2021;44:S53–72.
- Evert AB, Dennison M, Gardner CD, *et al*. Nutrition therapy for adults with diabetes or prediabetes: a consensus report. *Diabetes Care* 2019;42:731–54.
- Lehr AL, Driver SL, Stone NJ. The ABCDs of lifestyle counseling. *JAMA Cardiol* 2016;1:505–6.
- Seguin R, Connor L, Nelson M, *et al*. Understanding barriers and facilitators to healthy eating and active living in rural communities. *J Nutr Metab* 2014;2014:1–8.
- Cooksey-Stowers K, Schwartz MB, Brownell KD. Food swamps predict obesity rates better than food Deserts in the United States. *Int J Environ Res Public Health* 2017;14. doi:10.3390/ijerph14111366. [Epub ahead of print: 14 11 2017].
- Gucciardi E, Vahabi M, Norris N, *et al*. The intersection between food insecurity and diabetes: a review. *Curr Nutr Rep* 2014;3:324–32.
- Seligman HK, Bindman AB, Vittinghoff E, *et al*. Food insecurity is associated with diabetes mellitus: results from the National health examination and nutrition examination survey (NHANES) 1999–2002. *J Gen Intern Med* 2007;22:1018–23.
- Orr CJ, Keyserling TC, Ammerman AS, *et al*. Diet quality trends among adults with diabetes by socioeconomic status in the U.S.: 1999–2014. *BMC Endocr Disord* 2019;19:54.
- Berkowitz SA, Baggett TP, Wexler DJ, *et al*. Food insecurity and metabolic control among U.S. adults with diabetes. *Diabetes Care* 2013;36:3093–9.
- Berkowitz SA, Karter AJ, Corbie-Smith G, *et al*. Food Insecurity, Food "Deserts," and Glycemic Control in Patients With Diabetes: A Longitudinal Analysis. *Diabetes Care* 2018;41:1188–95.
- Shaheen M, Kibe LW, Schrode KM. Dietary quality, food security and glycemic control among adults with diabetes. *Clin Nutr ESPEN* 2021;46:336–42.
- Centers for Disease Control and Prevention (CDC), National Center for Health Statistics (NCHS). National health and nutrition examination survey; 2013–2018. Hyattsville, MD U.S. Department of Health and Human Services, Centers for Disease Control and Prevention; 2018.
- Centers for Disease Control and Prevention (CDC), National Center for Health Statistics (NCHS). National health and nutrition examination survey examination protocol: 2017–2018. Hyattsville, MD Department of Health and Human Services, Centers for Disease Control and Prevention; 2018.
- Centers for Disease Control and Prevention (CDC), National Center for Health Statistics (NCHS). National health and nutrition examination survey laboratory protocol: glucose, cholesterol and triglycerides. 2017–2018 2018.
- National Center for Health Statistics. National health and nutrition examination survey, data documentation, food security: 2017–2018, 2021. Available: https://www.cdc.gov/Nchs/Nhanes/2017-2018/FSQ_J.htm
- Bickel G, Nord M, Price C. Guide to measuring household food security USDA Food and Nutrition Service; 2000.
- National Center for Health Statistics. National health and nutrition examination survey: food security questionnaire, 2021. Available: https://www.cdc.gov/Nchs/Nhanes/2017-2018/FSQ_J.htm
- Division of Cancer Control & Population Sciences. Developing the healthy eating index, 2015, 2021. Available: <https://epi.grants.cancer.gov/hei/developing.html#2015c>
- Division of Cancer Control & Population Sciences. Healthy eating index SAS code, 2022. Available: <https://epi.grants.cancer.gov/hei/sas-code.html>
- National Center for Health Statistics. NHANES 2013–2018 dietary data: Analytic documentation. 2013–2018, 2021. Available: <https://www.cdc.gov/nchs/nhanes/search/datapage.aspx?Component=Dietary&CycleBeginYear=2017>
- Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem* 1972;18:499–502.
- ADA's standards of medical care in diabetes. *Clin Diabetes* 2021;39:128.
- Stark Casagrande S, Fradkin JE, Saydah SH, *et al*. The prevalence of meeting A1C, blood pressure, and LDL goals among people with diabetes, 1988–2010. *Diabetes Care* 2013;36:2271–9.
- Wang L, Li X, Wang Z, *et al*. Trends in prevalence of diabetes and control of risk factors in diabetes among US adults, 1999–2018. *JAMA* 2021;326:704.
- Darmon N, Drewnowski A. Does social class predict diet quality? *Am J Clin Nutr* 2008;87:1107–17.
- Darmon N, Drewnowski A. Contribution of food prices and diet cost to socioeconomic disparities in diet quality and health: a systematic review and analysis. *Nutr Rev* 2015;73:643–60.
- Pechey R, Monsivais P. Socioeconomic inequalities in the healthiness of food choices: exploring the contributions of food expenditures. *Prev Med* 2016;88:203–9.
- Kelly MP, Barker M. Why is changing health-related behaviour so difficult? *Public Health* 2016;136:109–16.
- Mayer VL, McDonough K, Seligman H, *et al*. Food insecurity, coping strategies and glucose control in low-income patients with diabetes. *Public Health Nutr* 2016;19:1103–11.
- Oddo VM, Mabli J. Association of participation in the supplemental nutrition assistance program and psychological distress. *Am J Public Health* 2015;105:e30–5.
- Lyles CR, Wolf MS, Schillinger D, *et al*. Food insecurity in relation to changes in hemoglobin A1c, self-efficacy, and fruit/vegetable intake during a diabetes educational intervention. *Diabetes Care* 2013;36:1448–53.
- Hill-Briggs F, Adler NE, Berkowitz SA, *et al*. Social determinants of health and diabetes: a scientific review. *Diabetes Care* 2020. doi:10.2337/dci20-0053. [Epub ahead of print: 02 Nov 2020].
- Ulrich-Lai YM, Fulton S, Wilson M, *et al*. Stress exposure, food intake and emotional state. *Stress* 2015;18:381–99.
- Aiyer JN, Raber M, Bello RS, *et al*. A pilot food prescription program promotes produce intake and decreases food insecurity. *Transl Behav Med* 2019;9:922–30.
- Caspi CE, Davey C, Barsness CB, *et al*. Applying the healthy eating Index-2015 in a sample of Choice-Based Minnesota food Pantries to test associations between food Pantry inventory, client food selection, and client diet. *J Acad Nutr Diet* 2021;121:2242–50.
- Ratcliffe C, McKernan S-M, Zhang S. How much does the supplemental nutrition assistance program reduce food insecurity? *Am J Agric Econ* 2011;93:1082–98.
- Franco M, Diez-Roux AV, Nettleton JA, *et al*. Availability of healthy foods and dietary patterns: the multi-ethnic study of atherosclerosis. *Am J Clin Nutr* 2009;89:897–904.
- Kanchi R, Lopez P, Rummo PE, *et al*. Longitudinal analysis of neighborhood food environment and diabetes risk in the Veterans administration diabetes risk cohort. *JAMA Netw Open* 2021;4:e2130789.
- Tabaie BP, Rundle AG, Wu WY, *et al*. Associations of residential socioeconomic, food, and built environments with glycemic control in persons with diabetes in New York City from 2007–2013. *Am J Epidemiol* 2018;187:736–45.
- Hager ER, Cockerham A, O'Reilly N, *et al*. Food swamps and food deserts in Baltimore City, MD, USA: associations with dietary behaviours among urban adolescent girls. *Public Health Nutr* 2017;20:2598–607.
- Travert A-S, Sidney Annerstedt K, Daivadanam M. Built environment and health behaviors: Deconstructing the black box of interactions-a review of reviews. *Int J Environ Res Public Health* 2019;16. doi:10.3390/ijerph16081454. [Epub ahead of print: 24 04 2019].

- 42 Stokols D, Allen J, Bellingham RL. The social ecology of health promotion: implications for research and practice. *Am J Health Promot* 1996;10:247–51.
- 43 American Diabetes Association Professional Practice Committee. 10. cardiovascular disease and risk management: standards of medical care in Diabetes-2022. *Diabetes Care* 2022;45:S144–74.
- 44 Moshfegh AJ, Rhodes DG, Baer DJ, *et al*. The US department of agriculture automated Multiple-Pass method reduces bias in the collection of energy intakes. *Am J Clin Nutr* 2008;88:324–32.
- 45 Subar AF, Freedman LS, Tooze JA, *et al*. Addressing current criticism regarding the value of self-report dietary data. *J Nutr* 2015;145:2639–45.
- 46 Centers for Disease Control and Prevention. Diabetes basics: quick facts 2021, 2022. Available: <https://www.cdc.gov/diabetes/basics/type2.html>

Appendix Table 1. Characteristics (% , SE) of U.S. Adults with Diagnosed Diabetes by Food Insecurity and Diet Quality

	Total	Food Insecurity			p-value	Diet Quality (HEI-2015) Quartiles				p-value
		Food secure	Marginal	Food insecure		4 th (High)	3 rd	2 nd	1 st (Low)	
Unweighted N		1356	330	589		504	512	555	560	
Total	100.0	68.0 (1.4)	12.7 (1.0)	19.3 (1.3)		26.2 (1.5)	23.3 (1.0)	25.8 (1.2)	24.7 (1.1)	
DEMOGRAPHICS										
Age, years										
Mean	60.4 (0.4)	61.6 (0.4)	58.8 (1.2)	57.1 (0.7)		62.4 (0.8)	60.9 (0.8)	60.0 (0.6)	57.9 (0.8)	
20-44	13.0 (0.7)	10.1 (0.9)	15.6 (3.0)	21.1 (1.8)	<0.001	7.9 (1.7)	11.9 (1.4)	14.3 (1.7)	16.3 (2.3)	0.006
45-64	45.6 (1.5)	45.8 (1.8)	45.4 (4.6)	46.7 (3.3)		44.7 (3.0)	43.4 (3.3)	47.3 (3.3)	51.3 (3.1)	
≥65	41.5 (1.5)	44.1 (1.7)	39.0 (4.3)	32.2 (2.9)		47.4 (3.3)	44.6 (3.8)	38.5 (2.8)	32.5 (2.7)	
Sex										
Men	53.0 (1.7)	57.3 (2.2)	43.8 (3.3)	43.2 (2.0)	<0.001	54.6 (3.7)	49.1 (3.2)	52.2 (3.7)	57.6 (2.9)	0.150
Women	47.0 (1.7)	42.7 (2.1)	56.2 (3.3)	56.8 (2.0)		45.4 (3.7)	50.9 (3.2)	47.8 (3.7)	42.4 (2.9)	
Race and ethnicity										
Non-Hispanic White	63.4 (2.2)	69.1 (2.2)	54.2 (5.0)	49.3 (4.0)	<0.001	61.6 (3.4)	60.0 (4.0)	69.6 (2.7)	68.0 (3.0)	<0.001
Non-Hispanic Black	14.0 (1.4)	12.4 (1.4)	18.7 (3.1)	17.4 (2.2)		12.3 (1.8)	17.0 (2.3)	12.6 (1.7)	13.2 (2.0)	
Hispanic/Latino	16.5 (1.8)	12.1 (1.6)	21.1 (3.5)	29.3 (3.2)		17.0 (2.2)	16.5 (2.6)	15.4 (2.4)	16.2 (2.1)	
Non-Hispanic Asian	6.1 (0.8)	6.5 (0.9)	6.0 (1.4)	4.0 (0.8)		9.2 (1.3)	6.5 (1.6)	2.4 (0.7)	2.7 (0.6)	
SOCIODEMOGRAPHIC CHARACTERISTICS										
Poverty Income Ratio										
<1.00	15.3 (1.2)	8.0 (1.0)	17.7 (3.2)	41.3 (3.1)	<0.001	10.6 (1.6)	14.1 (2.1)	16.1 (2.2)	18.6 (2.5)	0.010
1.00-2.00	20.0 (1.2)	15.5 (1.2)	30.1 (3.2)	31.9 (2.8)		18.1 (2.1)	20.9 (2.4)	20.7 (2.1)	21.0 (2.4)	
>2.00	55.7 (1.9)	69.9 (1.9)	45.1 (4.8)	20.4 (2.7)		61.9 (3.4)	56.4 (3.7)	56.7 (3.2)	53.2 (3.4)	
Missing	9.1 (0.9)	6.6 (0.7)	7.2 (2.0)	6.4 (1.3)		9.5 (1.9)	8.6 (1.9)	6.6 (1.8)	7.2 (1.9)	
Education										
<High school	19.9 (1.4)	15.0 (1.3)	18.4 (2.7)	37.1 (2.7)	<0.001	16.8 (1.9)	19.3 (2.3)	18.1 (2.1)	22.9 (2.8)	0.130
High school graduate	25.0 (1.2)	23.1 (1.6)	34.2 (4.1)	27.2 (2.1)		23.6 (2.6)	27.7 (3.1)	25.1 (2.9)	26.4 (2.7)	
Some college	33.8 (1.5)	35.0 (2.2)	34.6 (4.0)	30.1 (2.3)		29.8 (3.0)	33.3 (3.0)	37.4 (3.3)	35.7 (3.7)	
College graduate or	21.4 (1.5)	27.0 (2.1)	12.8 (2.6)	5.6 (1.2)		29.8 (3.4)	19.7 (3.1)	19.5 (2.6)	15.0 (3.2)	

Above										
Received SNAP benefits, past 12 mo.										
Yes	20.7 (1.3)	11.8 (1.3)	25.5 (3.0)	48.9 (3.2)	<0.001	25.0 (2.2)	22.9 (2.3)	17.0 (1.9)	15.9 (2.0)	0.008
No	79.3 (1.3)	88.2 (1.3)	74.5 (3.2)	51.1 (3.2)		75.0 (2.2)	77.1 (2.3)	83.0 (1.9)	84.1 (2.0)	
Health Insurance										
Uninsured	8.1 (0.8)	6.0 (2.1)	9.8 (2.1)	14.8 (1.8)	<0.001	6.0 (1.1)	8.0 (1.3)	10.0 (1.8)	9.1 (1.7)	0.170
Insured	91.9 (0.8)	94.0 (0.8)	90.2 (2.1)	85.3 (1.8)		94.0 (1.1)	92.0 (1.3)	90.0 (1.8)	91.0 (1.7)	
Health Insurance Type										
Private	60.0 (1.8)	65.4 (2.2)	62.4 (3.9)	36.0 (3.0)	<0.001	60.2 (3.6)	64.2 (3.6)	60.1 (3.9)	59.9 (2.5)	0.650
Medicare	46.7 (1.8)	46.0 (1.1)	43.7 (4.3)	50.0 (3.1)	0.350	47.9 (3.2)	49.0 (4.0)	47.2 (3.0)	40.7 (3.3)	0.310
Medicaid	13.9 (1.1)	5.7 (1.2)	16.4 (3.0)	32.5 (2.6)	<0.001	11.4 (1.8)	12.5 (2.0)	15.1 (2.5)	15.0 (1.8)	0.260
Military/Other Government	15.2 (1.1)	15.4 (1.56)	9.5 (1.87)	19.5 (2.6)	0.006	14.1 (2.0)	16.1 (2.3)	18.3 (3.0)	11.0 (1.5)	0.150
HEALTHCARE UTILIZATION										
Seen a Physician in past year										
Yes	96.9 (0.4)	97.9 (0.6)	96.4 (1.4)	93.2 (1.2)	0.010	96.8 (1.2)	97.1 (0.9)	97.8 (1.3)	95.6 (1.3)	0.510
No	3.1 (0.4)	2.1 (0.6)	3.6 (1.4)	6.8 (1.2)		3.2 (1.2)	2.9 (0.9)	2.21 (0.6)	4.4 (1.3)	
Seen a Diabetes specialist in past year										
Yes	29.9 (1.2)	30.1 (1.8)	27.3 (3.1)	31.6 (2.1)	0.820	33.7 (3.5)	33.3 (2.7)	23.1 (2.2)	26.8 (3.2)	0.030
No	34.8 (1.7)	35.0 (2.1)	34.6 (3.3)	33.3 (2.4)		33.5 (2.5)	34.5 (3.3)	36.7 (3.2)	34.8 (3.1)	
Never	35.3 (1.5)	34.9 (1.7)	38.2 (3.7)	35.1 (2.6)		32.8 (2.6)	32.3 (2.8)	40.2 (3.0)	38.3 (3.7)	
A1c Checked in past year										
Yes	82.5 (1.1)	85.2 (1.3)	79.6 (3.1)	73.9 (2.9)	0.001	88.0 (1.6)	84.5 (1.7)	81.6 (2.0)	76.3 (2.9)	0.040
No	13.0 (0.9)	10.3 (1.0)	16.3 (2.8)	21.2 (2.5)		9.0 (1.5)	11.9 (1.7)	13.8 (2.0)	16.9 (2.4)	
Do not know	4.6 (0.6)	4.6 (0.8)	4.1 (1.0)	4.9 (1.2)		3.0 (0.6)	3.56 (1.1)	4.6 (0.9)	6.9 (1.8)	
CLINICAL CHARACTERISTICS										
Smoking Status										

Current	14.4 (1.1)	10.9 (1.3)	17.0 (3.1)	24.4 (2.4)	0.001	5.3 (1.3)	12.0 (2.0)	16.1 (2.4)	21.8 (2.7)	<0.001
Former	36.5 (1.4)	39.0 (2.0)	35.4 (3.8)	30.0 (2.7)		42.3 (3.6)	39.0 (3.3)	36.1 (2.6)	32.1 (3.1)	
Never	49.1 (1.3)	50.2 (1.9)	47.7 (4.1)	45.5 (2.8)		52.4 (3.5)	49.0 (3.5)	47.8 (3.2)	46.2 (3.4)	
Diabetes Medication										
Insulin only	12.8 (1.0)	11.1 (1.1)	15.9 (3.2)	15.4 (2.6)	0.020	12.1 (2.2)	10.5 (1.8)	13.0 (1.9)	11.5 (1.8)	0.110
Oral medication only	55.5 (1.4)	59.2 (2.0)	50.8 (4.0)	44.5 (2.4)		59.1 (2.9)	59.4 (3.2)	52.8 (2.6)	53.7 (3.7)	
Both insulin and oral medications	14.8 (1.0)	13.6 (1.3)	16.8 (2.6)	18.7 (1.8)		11.8 (1.7)	14.8 (2.1)	17.2 (2.0)	15.4 (2.6)	
None reported	17.0 (0.9)	16.1 (1.4)	16.5 (2.4)	21.3 (1.9)		17.1 (2.5)	15.3 (2.2)	17.1 (2.6)	19.5 (2.5)	
Taking Antihypertensive Medication	62.9 (1.7)	65.1 (2.1)	57.1 (4.3)	60.5 (2.1)	0.050	64.8 (3.0)	61.3 (3.2)	64.7 (3.4)	60.3 (3.1)	0.550
Taking Antilipidemic Medication	59.4 (1.3)	62.4 (1.7)	50.9 (4.1)	55.2 (2.7)	0.008	60.5 (3.6)	54.0 (2.7)	63.3 (3.3)	60.8 (3.2)	0.070
BMI, kg/m²										
Mean	33.0 (0.3)	32.8 (0.3)	33.3 (0.6)	34.1 (0.5)		31.5 (0.4)	33.2 (0.5)	34.0 (0.6)	34.6 (0.6)	
<25.0 (Underweight/healthy weight)	10.5 (0.9)	10.8 (1.1)	8.7 (1.9)	10.2 (1.4)	0.606	10.9 (1.8)	11.5 (1.8)	9.6 (1.4)	6.7 (1.8)	0.035
25.0-29.9 (Overweight)	27.7 (1.2)	27.5 (1.6)	30.9 (4.4)	24.6 (2.1)		33.0 (3.1)	28.8 (3.1)	22.2 (2.4)	23.0 (3.1)	
≥30 (Obesity)	61.8 (1.7)	61.6 (2.2)	60.5 (4.3)	65.1 (2.4)		56.2 (3.6)	59.7 (3.1)	68.2 (2.7)	70.2 (3.7)	
FOOD INSECURITY AND DIET QUALITY										
Food Insecurity Status										
Food secure	68.0 (1.4)	NA	NA	NA		74.4 (2.7)	67.8 (2.7)	67.4 (2.8)	63.3 (2.9)	0.100
Marginal	12.7 (1.0)	NA	NA	NA		11.6 (1.7)	13.1 (1.8)	12.2 (2.3)	12.5 (1.7)	
Food insecure	19.3 (1.3)	NA	NA	NA		14.1 (1.8)	19.2 (2.4)	20.5 (2.3)	24.2 (2.7)	
Healthy Eating Index										
Mean (SE)	53.6 (0.4)	54.3 (0.5)	53.0 (1.2)	51.0 (0.7)		NA	NA	NA	NA	
Quartile 4 (High)	26.2 (1.5)	28.6 (1.9)	24.7 (3.9)	19.0 (2.4)	0.100	NA	NA	NA	NA	
Quartile 3	23.3 (1.0)	22.9 (1.6)	24.5 (2.6)	22.8 (2.1)		NA	NA	NA	NA	
Quartile 2	25.8 (1.2)	25.5 (1.6)	25.6 (3.87)	27.3 (2.58)		NA	NA	NA	NA	
Quartile 1 (Low)	24.7 (1.1)	22.9 (1.4)	25.2 (3.3)	30.9 (2.5)		NA	NA	NA	NA	
METABOLIC										

OUTCOMES										
A1c ≥7.0%	49.8 (1.8)	47.7 (2.3)	52.2 (3.8)	54.8 (2.6)	0.120	43.9 (3.4)	46.9 (3.0)	51.7 (3.2)	54.9 (4.1)	0.050
A1c ≥8.0%	26.1 (1.3)	22.8 (1.8)	29.8 (3.8)	34.2 (2.1)	0.002	20.1 (2.6)	25.8 (2.7)	25.4 (2.5)	28.2 (2.6)	0.120
A1c ≥9.0%	14.4 (1.0)	12.2 (1.3)	15.0 (2.7)	21.9 (2.2)	0.003	11.9 (2.2)	14.3 (2.1)	12.6 (1.7)	18.1 (2.4)	0.250
Blood Pressure ≥140/90 mmHg	28.5 (1.5)	27.8 (2.0)	30.1 (3.4)	29.4 (2.8)	0.800	29.6 (2.7)	25.4 (2.7)	29.7 (3.0)	28.3 (2.7)	0.550
Blood Pressure ≥130/80 mmHg	52.0 (1.5)	51.5 (2.3)	52.9 (4.7)	51.6 (3.1)	0.970	51.9 (3.1)	51.6 (3.2)	53.4 (2.8)	50.9 (3.3)	0.950
LDL-C ≥100 mg/dL	41.8 (2.4)	38.3 (3.2)	50.2 (5.6)	48.3 (3.2)	0.060	35.9 (4.6)	43.3 (4.5)	45.7 (5.0)	44.0 (4.3)	0.470
LDL-C ≥130 mg/dL	18.0 (1.7)	16.2 (2.4)	23.6 (5.3)	20.1 (3.1)	0.430	16.9 (3.0)	17.9 (3.0)	22.9 (3.8)	14.2 (3.9)	0.250
Elevated ABCs (A1c ≥7.0, BP ≥130/80, and LDL ≥100)										
None	15.7 (2.1)	19.7 (3.2)	10.9 (2.8)	7.4 (1.8)	0.017	18.1 (4.9)	15.9 (3.6)	12.9 (3.2)	18.6 (5.0)	0.567
One elevated ABC	36.4 (2.7)	34.9 (3.7)	37.9 (5.4)	40.2 (3.3)	0.585	35.7 (4.9)	29.6 (4.3)	41.4 (5.8)	33.5 (4.9)	0.437
Two elevated ABCs	35.3 (2.9)	33.3 (3.7)	37.0 (5.6)	40.8 (4.1)	0.411	36.5 (6.1)	42.0 (4.8)	34.8 (6.0)	33.0 (4.6)	0.571
Three elevated ABCs	12.3 (1.6)	12.2 (2.0)	14.2 (4.3)	11.6 (2.2)	0.808	9.8 (2.4)	12.5 (3.0)	10.9 (2.1)	15.0 (4.5)	0.732
Any one elevated ABC	84.3 (2.1)	80.9 (3.0)	89.7 (2.6)	92.0 (1.8)	0.015	81.4 (4.8)	84.8 (3.5)	88.0 (2.9)	82.3 (4.4)	0.424
Elevated ABCs (A1c ≥8.0, BP ≥140/90 and LDL ≥130)										
None	45.0 (2.6)	48.5 (3.6)	35.7 (5.3)	39.3 (3.7)	0.055	45.6 (5.2)	42.5 (4.5)	43.2 (5.8)	49.7 (4.9)	0.776
One elevated ABC	40.6 (2.4)	38.4 (3.0)	47.6 (6.3)	43.4 (4.6)	0.228	44.1 (5.2)	44.1 (4.5)	41.5 (6.0)	32.2 (4.4)	0.286
Two elevated ABCs	11.4 (1.6)	9.8 (1.9)	15.4 (3.6)	14.4 (3.0)	0.135	7.6 (2.1)	11.5 (2.1)	10.3 (2.5)	16.5 (4.6)	0.320
Three elevated ABCs	3.0 (0.6)	3.4 (0.75)	³	2.9 (1.6) ²	0.238	2.8 (1.5) ²	2.0 (0.7) ¹	5.1 (2.0) ¹	1.6 (0.8) ²	0.327
Any one elevated ABC	55.2 (2.6)	51.6 (3.5)	64.1 (5.3)	61.3 (3.7)	0.034	54.1 (5.1)	59.5 (4.4)	57.4 (5.5)	49.4 (4.8)	0.572

Health insurance categories are not mutually exclusive; estimates are among those with insurance

LDL-C were determined among the fasting sample only

Relative Standard Error (RSE): ¹RSE 30-40%; ²RSE 40-50%; ³RSE>50%, data do not meet the National Center for Health Statistics Data

Presentation Standards for Proportions

ABC, A1c, blood pressure, cholesterol

BP, blood pressure

BMI, body mass index

HEI, Healthy Eating Index

LDL-C, low-density lipoprotein cholesterol

Appendix Table 2. Odds Ratio (95% CI) for Sub-Optimal Diabetes ABC Management by Food Insecurity and Diet Quality

	Total % (SE)	Food Insecurity			Diet Quality (HEI-2015) Quartiles			
		Food secure	Marginal	Food insecure	Q4 (High)	Q3	Q2	Q1 (Low)
HbA1c ≥7.0%	49.8 (1.8)							
Unadjusted		1.00	1.20 (0.85, 1.68)	1.33 (1.01, 1.75)	1.00	1.13 (0.81, 1.57)	1.36 (0.94, 1.97)	1.56 (1.04, 2.32)
Model 1		1.00	1.24 (0.85, 1.80)	1.36 (1.01, 1.83)	1.00	1.10 (0.78, 1.55)	1.36 (0.95, 1.95)	1.47 (0.96, 2.26)
Model 2		1.00	1.23 (0.85, 1.80)	1.44 (1.06, 1.96)	1.00	1.09 (0.77, 1.56)	1.39 (0.97, 1.98)	1.49 (0.95, 2.33)
Model 3		1.00	1.24 (0.85, 1.80)	1.44 (1.06, 1.97)	1.00	1.11 (0.78, 1.58)	1.42 (0.99, 2.03)	1.53 (0.97, 2.41)
Model 4		1.00	1.16 (0.78, 1.73)	1.47 (1.01, 2.16)	1.00	1.10 (0.76, 1.59)	1.27 (0.86, 1.88)	1.63 (1.10, 2.42)
Model 5		1.00	1.21 (0.78, 1.87)	1.62 (1.07, 2.47)	1.00	1.06 (0.72, 1.55)	1.29 (0.87, 1.92)	1.61 (1.06, 2.46)
HbA1c ≥8.0%	26.1 (1.3)							
Unadjusted		1.00	1.44 (0.92, 2.23)	1.76 (1.30, 2.37)	1.00	1.38 (0.93, 2.04)	1.36 (0.90, 2.06)	1.56 (1.06, 2.31)
Model 1		1.00	1.30 (0.82, 2.07)	1.52 (1.12, 2.05)	1.00	1.31 (0.89, 1.93)	1.27 (0.84, 1.93)	1.39 (0.92, 2.11)
Model 2		1.00	1.26 (0.77, 2.06)	1.48 (1.04, 2.10)	1.00	1.30 (0.88, 1.91)	1.24 (0.82, 1.89)	1.37 (0.90, 2.07)
Model 3		1.00	1.26 (0.77, 2.05)	1.45 (1.01, 2.09)	1.00	1.33 (0.91, 1.95)	1.29 (0.85, 1.95)	1.41 (0.93, 2.12)
Model 4		1.00	1.33 (0.81, 2.17)	1.54 (1.07, 2.23)	1.00	1.46 (0.89, 2.39)	1.31 (0.80, 2.15)	1.67 (1.09, 2.57)
Model 5		1.00	1.31 (0.81, 2.14)	1.71 (1.14, 2.56)	1.00	1.42 (0.85, 2.36)	1.35 (0.80, 2.26)	1.60 (1.01, 2.53)
HbA1c ≥9.0%	14.4 (1.0)							
Unadjusted		1.00	1.27 (0.77, 2.11)	2.03 (1.41, 2.91)	1.00	1.24 (0.76, 2.03)	1.07 (0.64, 1.80)	1.64 (0.93, 2.91)
Model 1		1.00	1.05 (0.61, 1.81)	1.61 (1.11, 2.34)	1.00	1.16 (0.70, 1.92)	0.91 (0.53, 1.55)	1.44 (0.78, 2.65)
Model 2		1.00	0.96 (0.55, 1.67)	1.40 (0.91, 2.17)	1.00	1.15 (0.70, 1.91)	0.89 (0.51, 1.52)	1.42 (0.77, 2.62)
Model 3		1.00	0.93 (0.54, 1.62)	1.34 (0.86, 2.11)	1.00	1.20 (0.73, 1.96)	0.93 (0.54, 1.63)	1.47 (0.81, 2.68)
Model 4		1.00	1.02 (0.56, 1.85)	1.46 (0.97, 2.21)	1.00	1.33 (0.74, 2.41)	0.97 (0.53, 1.78)	1.76 (0.97, 3.19)
Model 5		1.00	1.16 (0.61, 2.21)	1.59 (1.04, 2.43)	1.00	1.37 (0.76, 2.47)	0.98 (0.52, 1.84)	1.66 (0.91, 3.00)
Blood Pressure ≥140/90 mmHg	28.5 (1.5)							
Unadjusted		1.00	1.12 (0.76, 1.65)	1.08 (0.79, 1.50)	1.00	0.81 (0.56, 1.16)	1.00 (0.69, 1.46)	0.94 (0.68, 1.29)
Model 1		1.00	1.14 (0.81, 1.60)	1.17 (0.84, 1.64)	1.00	0.81 (0.54, 1.22)	1.08 (0.71, 1.64)	1.00 (0.68, 1.48)
Model 2		1.00	1.10 (0.79, 1.55)	1.10 (0.74, 1.63)	1.00	0.80 (0.53, 1.19)	1.03 (0.68, 1.57)	0.99 (0.66, 1.48)
Model 3		1.00	1.11 (0.79, 1.55)	1.11 (0.74, 1.64)	1.00	0.79 (0.53, 1.19)	1.03 (0.68, 1.55)	0.98 (0.66, 1.46)
Model 4		1.00	1.10 (0.80, 1.63)	1.05 (0.69, 1.60)	1.00	0.85 (0.55, 1.31)	1.05 (0.69, 1.60)	1.08 (0.68, 1.72)
Model 5		1.00	1.07 (0.75, 1.54)	0.99 (0.64, 1.53)	1.00	0.81 (0.52, 1.24)	1.01 (0.66, 1.54)	1.07 (0.67, 1.69)
Blood Pressure ≥130/80 mmHg	52.0 (1.5)							
Unadjusted		1.00	1.06 (0.67, 1.67)	1.01 (0.73, 1.40)	1.00	0.99 (0.69, 1.42)	1.06 (0.75, 1.50)	0.96 (0.69, 1.34)
Model 1		1.00	1.10 (0.71, 1.71)	1.08 (0.77, 1.52)	1.00	1.02 (0.70, 1.49)	1.11 (0.76, 1.63)	0.96 (0.66, 1.40)
Model 2		1.00	1.02 (0.66, 1.60)	0.96 (0.66, 1.40)	1.00	1.00 (0.69, 1.46)	1.10 (0.74, 1.63)	0.94 (0.63, 1.39)

Model 3		1.00	1.01 (0.66, 1.56)	0.96 (0.66, 1.38)	1.00	1.01 (0.69, 1.48)	1.13 (0.77, 1.67)	0.95 (0.64, 1.40)
Model 4		1.00	1.01 (0.65, 1.58)	0.91 (0.61, 1.35)	1.00	1.03 (0.67, 1.58)	1.12 (0.73, 1.74)	0.95 (0.62, 1.45)
Model 5		1.00	0.96 (0.59, 1.54)	0.88 (0.58, 1.33)	1.00	1.00 (0.65, 1.55)	1.12 (0.72, 1.74)	0.93 (0.60, 1.44)
LDL-C ≥100 mg/dL	41.8 (2.4)							
Unadjusted		1.00	1.62 (0.98, 2.68)	1.50 (1.05, 2.15)	1.00	1.37 (0.79, 2.36)	1.50 (0.91, 2.50)	1.40 (0.78, 2.52)
Model 1		1.00	1.36 (0.78, 2.39)	1.19 (0.76, 1.85)	1.00	1.15 (0.64, 2.06)	1.20 (0.71, 2.02)	1.12 (0.61, 2.06)
Model 2		1.00	1.21 (0.66, 2.20)	1.02 (0.65, 1.60)	1.00	1.19 (0.66, 2.15)	1.26 (0.76, 2.08)	1.25 (0.74, 2.04)
Model 3		1.00	1.18 (0.62, 2.21)	1.08 (0.69, 1.69)	1.00	1.16 (0.65, 2.06)	1.25 (0.74, 2.12)	1.10 (0.59, 2.02)
Model 4		1.00	0.98 (0.49, 1.99)	1.07 (0.63, 1.80)	1.00	1.17 (0.65, 2.10)	1.23 (0.72, 2.09)	1.05 (0.56, 1.97)
Model 5		1.00	0.95 (0.47, 1.95)	1.01 (0.60, 1.71)	1.00	1.06 (0.55, 2.02)	1.20 (0.70, 2.05)	0.97 (0.54, 1.75)
LDL-C ≥ 130mg/dL	18.0 (1.7)							
Unadjusted		1.00	1.60 (0.76, 3.37)	1.31 (0.77, 2.24)	1.00	1.07 (0.60, 1.91)	1.46 (0.75, 2.85)	0.82 (0.36, 1.88)
Model 1		1.00	1.34 (0.63, 2.83)	1.06 (0.60, 1.85)	1.00	0.85 (0.45, 1.61)	1.09 (0.55, 2.18)	0.64 (0.29, 1.44)
Model 2		1.00	1.17 (0.56, 2.47)	0.94 (0.46, 1.92)	1.00	0.88 (0.48, 1.63)	1.18 (0.59, 2.36)	0.65 (0.30, 1.37)
Model 3		1.00	1.16 (0.54, 2.48)	0.99 (0.50, 1.97)	1.00	0.91 (0.50, 1.67)	1.21 (0.60, 2.44)	0.63 (0.29, 1.35)
Model 4		1.00	0.77 (0.38, 1.55)	0.87 (0.42, 1.80)	1.00	0.75 (0.41, 1.38)	1.32 (0.63, 2.73)	0.56 (0.27, 1.18)
Model 5		1.00	0.70 (0.35, 1.40)	0.82 (0.40, 1.69)	1.00	0.76 (0.41, 1.40)	1.31 (0.61, 2.84)	0.57 (0.28, 1.18)
Any one elevated ABC [HbA1c ≥7.0%, BP ≥130/80mmHg, LDL ≥100mg/dL]	84.3 (2.1)							
Unadjusted		1.00	2.06 (1.06, 4.03)	2.71 (1.46, 5.06)	1.00	1.27 (0.55, 2.96)	1.68 (0.77, 3.67)	1.06 (0.45, 2.52)
Model 1		1.00	1.77 (0.95, 3.31)	2.09 (1.13, 3.88)	1.00	0.98 (0.41, 2.34)	1.30 (0.57, 2.96)	0.76 (0.30, 1.92)
Model 2		1.00	1.67 (0.91, 3.07)	1.88 (1.07, 3.30)	1.00	0.96 (0.40, 2.31)	1.28 (0.56, 2.97)	0.70 (0.28, 1.78)
Model 3		1.00	1.66 (0.90, 3.07)	1.89 (1.05, 3.39)	1.00	0.97 (0.41, 2.33)	1.30 (0.56, 3.00)	0.70 (0.28, 1.77)
Model 4		1.00	1.97 (0.91, 4.28)	1.91 (0.94, 3.89)	1.00	1.08 (0.40, 2.92)	1.41 (0.62, 3.22)	0.64 (0.24, 1.70)
Model 5		1.00	1.89 (0.84, 4.26)	1.89 (0.91, 4.94)	1.00	1.04 (0.38, 2.85)	1.30 (0.57, 2.99)	0.64 (0.23, 1.76)
Any one elevated ABC [HbA1c ≥8.0% and BP ≥140/90mmHg and LDL ≥130mg/dL]	55.2 (2.6)							
Unadjusted		1.00	1.57 (0.99, 2.82)	1.49 (1.00, 2.22)	1.00	1.24 (0.74, 2.08)	1.14 (0.65, 1.99)	0.83 (0.49, 1.39)
Model 1		1.00	1.45 (0.87, 2.41)	1.34 (0.91, 1.97)	1.00	1.08 (0.64, 1.82)	1.07 (0.59, 1.95)	0.76 (0.43, 1.34)
Model 2		1.00	1.28 (0.74, 2.21)	1.25 (0.85, 1.84)	1.00	1.06 (0.62, 1.81)	1.06 (0.58, 1.96)	0.75 (0.41, 1.36)
Model 3		1.00	1.26 (0.73, 2.19)	1.28 (0.85, 1.93)	1.00	1.10 (0.66, 1.86)	1.12 (0.61, 2.08)	0.76 (0.42, 1.37)
Model 4		1.00	1.18 (0.65, 2.12)	1.33 (0.84, 2.11)	1.00	1.15 (0.62, 2.11)	1.25 (0.67, 2.33)	0.78 (0.42, 1.42)
Model 5		1.00	1.14 (0.63, 2.06)	1.25 (0.76, 2.07)	1.00	1.10 (0.58, 2.06)	1.19 (0.61, 2.33)	0.72 (0.38, 1.37)

Model 1: adjusted for age, sex, race/ethnicity

Model 2: additionally adjusted for poverty income ratio, education, health insurance

Model 3: additionally adjusted for healthcare utilization (seeing physicians in past year, having HbA1c checked)

Model 4: additionally adjusted for smoking status, diabetes medications, blood pressure medication, cholesterol medication, BMI

Model 5: additionally adjusted for food security (when HEI is main covariate) or HEI (when food security is main covariate),

Boldface p-value indicates statistical significance of $p < 0.05$

SE, standard error

Appendix Table 3. Odds Ratio (95% CI) for Sub-Optimal Diabetes ABC Management by Food Security and Diet Quality

	Food Security & Diet Quality Categories (HEI-2015)^a			
	Food Secure & High Diet Quality (Q4)	Food Secure & Low/Moderate Diet Quality (Q1-3)	Food Insecure & High Diet Quality (Q4)	Food Insecure & Low/Moderate Diet Quality (Q1-3)
A1c \geq7.0%				
Unadjusted	1.00	1.31 (0.88, 1.96)	1.33 (0.75, 2.34)	1.71 (1.18, 2.48)
Model 1	1.00	1.21 (0.79, 1.86)	1.19 (0.69, 2.07)	1.71 (1.15, 2.56)
Model 2	1.00	1.22 (0.79, 1.89)	1.23 (0.70, 2.15)	1.82 (1.21, 2.73)
Model 3	1.00	1.24 (0.80, 1.90)	1.18 (0.67, 2.08)	1.86 (1.24, 2.78)
Model 4	1.00	1.30 (0.86, 1.97)	1.45 (0.82, 2.57)	1.82 (1.21, 2.75)
A1c \geq8.0%				
Unadjusted	1.00	1.48 (0.88, 2.47)	2.07 (1.08, 3.87)	2.36 (1.49, 3.73)
Model 1	1.00	1.30 (0.77, 2.19)	1.54 (0.80, 2.96)	1.93 (1.20, 3.09)
Model 2	1.00	1.29 (0.77, 2.14)	1.49 (0.78, 2.83)	1.86 (1.13, 3.06)
Model 3	1.00	1.30 (0.79, 2.17)	1.41 (0.75, 2.67)	1.89 (1.13, 3.15)
Model 4	1.00	1.50 (0.85, 2.64)	1.66 (0.91, 3.01)	2.22 (1.30, 3.79)
A1c \geq9.0%				
Unadjusted	1.00	1.19 (0.61, 2.33)	1.75 (0.81, 3.76)	2.28 (1.17, 4.43)
Model 1	1.00	0.98 (0.49, 1.92)	1.12 (0.52, 2.38)	1.65 (0.82, 3.30)
Model 2	1.00	0.96 (0.49, 1.88)	1.01 (0.46, 2.24)	1.50 (0.74, 3.06)
Model 3	1.00	0.98 (0.51, 1.90)	0.91 (0.41, 1.99)	1.51 (0.72, 3.16)
Model 4	1.00	1.14 (0.54, 2.40)	1.03 (0.45, 2.40)	1.71 (0.82, 3.57)
Blood Pressure \geq140/90 mmHg				
Unadjusted	1.00	0.82 (0.57, 1.18)	0.85 (0.52, 1.40)	0.96 (0.64, 1.44)
Model 1	1.00	0.86 (0.57, 1.30)	0.91 (0.53, 1.57)	1.02 (0.65, 1.61)
Model 2	1.00	0.85 (0.56, 1.27)	0.87 (0.51, 1.50)	0.95 (0.59, 1.55)
Model 3	1.00	0.84 (0.56, 1.26)	0.88 (0.51, 1.53)	0.96 (0.59, 1.55)
Model 4	1.00	0.89 (0.58, 1.36)	0.87 (0.51, 1.51)	0.98 (0.60, 1.60)
Blood Pressure \geq130/80 mmHg				
Unadjusted	1.00	0.99 (0.67, 1.47)	1.01 (0.58, 1.76)	0.99 (0.68, 1.46)
Model 1	1.00	1.04 (0.67, 1.63)	1.16 (0.65, 2.08)	1.08 (0.71, 1.65)
Model 2	1.00	1.04 (0.66, 1.63)	1.06 (0.57, 1.96)	0.96 (0.62, 1.47)
Model 3	1.00	1.04 (0.67, 1.64)	1.02 (0.56, 1.86)	0.97 (0.64, 1.47)
Model 4	1.00	1.08 (0.66, 1.75)	1.06 (0.59, 1.88)	0.94 (0.61, 1.43)
LDL-C \geq100 mg/dL				
Unadjusted	1.00	1.50 (0.80, 2.81)	2.12 (0.88, 5.13)	2.18 (1.23, 3.86)
Model 1	1.00	1.22 (0.65, 2.32)	1.61 (0.60, 4.29)	1.50 (0.83, 2.70)
Model 2	1.00	1.24 (0.66, 2.33)	1.35 (0.52, 3.48)	1.34 (0.75, 2.40)
Model 3	1.00	1.17 (0.63, 2.18)	1.18 (0.45, 3.12)	1.33 (0.74, 2.39)
Model 4	1.00	1.06 (0.56, 1.98)	0.96 (0.26, 2.58)	1.07 (0.53, 2.18)

LDL-C \geq130 mg/dL				
Unadjusted	1.00	1.36 (0.58, 3.17)	3.02 (0.92, 9.96)	1.67 (0.75, 3.71)
Model 1	1.00	1.01 (0.45, 2.28)	2.17 (0.61, 7.71)	1.07 (0.49, 2.36)
Model 2	1.00	1.03 (0.48, 2.20)	1.90 (0.50, 7.24)	1.03 (0.43, 2.47)
Model 3	1.00	1.02 (0.49, 2.11)	1.80 (0.47, 6.97)	1.05 (0.47, 2.37)
Model 4	1.00	1.00 (0.48, 2.10)	1.39 (0.34, 5.67)	0.73 (0.30, 1.75)
At least one elevated ABC (A1c \geq7.0, BP \geq130/80, LDL \geq100)				
Unadjusted	1.00	1.14 (0.52, 2.52)	2.60 (1.06, 6.41)	2.91 (1.26, 6.72)
Model 1	1.00	0.89 (0.39, 2.02)	1.95 (0.79, 4.85)	1.86 (0.79, 4.38)
Model 2	1.00	0.87 (0.38, 1.98)	1.75 (0.65, 4.68)	1.72 (0.75, 3.99)
Model 3	1.00	0.87 (0.38, 1.96)	1.63 (0.60, 4.37)	1.73 (0.74, 4.01)
Model 4	1.00	0.91 (0.38, 2.15)	1.63 (0.63, 4.24)	1.91 (0.69, 5.31)
At least one elevated ABC (A1c \geq8.0 and BP \geq140/90 and LDL \geq130)				
Unadjusted	1.00	1.10 (0.61, 1.96)	2.78 (1.41, 5.49)	1.54 (0.93, 2.56)
Model 1	1.00	0.99 (0.53, 1.84)	2.18 (1.06, 4.49)	1.28 (0.75, 2.16)
Model 2	1.00	0.97 (0.51, 1.82)	1.96 (0.93, 4.15)	1.20 (0.69, 2.07)
Model 3	1.00	0.98 (0.53, 1.81)	1.76 (0.83, 3.72)	1.23 (0.71, 2.14)
Model 4	1.00	1.06 (0.53, 2.09)	1.68 (0.77, 3.70)	1.25 (0.66, 2.34)

Model 1: adjusted for age, sex, race/ethnicity

Model 2: additionally adjusted for PIR, education, health insurance

Model 3: additionally adjusted for healthcare utilization (seeing physicians in past year, having HbA1c checked)

Model 4: additionally adjusted for smoking status, diabetes medications, blood pressure medication, cholesterol medication, BMI

^a Food insecurity is defined as marginal or low/very low food security; food secure is defined as full food security; Low/moderate diet quality is defined as HEI-2015 score in the 1st-3rd quartiles; High diet quality is defined as HEI-2015 in the 4th quartile

Boldface p-value indicates statistical significance of $p < 0.05$