

# Higher risk of severe hypoglycemia in children and adolescents with a rapid loss of C-peptide during the first 6 years after type 1 diabetes diagnosis

Annika Grönberg <sup>1</sup>, Daniel Espes <sup>2,3</sup>, Per-Ola Carlsson,<sup>2,3</sup> Johnny Ludvigsson<sup>4</sup>

**To cite:** Grönberg A, Espes D, Carlsson P-O, *et al.* Higher risk of severe hypoglycemia in children and adolescents with a rapid loss of C-peptide during the first 6 years after type 1 diabetes diagnosis. *BMJ Open Diab Res Care* 2022;**10**:e002991. doi:10.1136/bmjdr-2022-002991

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjdr-2022-002991>).

Received 16 June 2022  
Accepted 18 October 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

<sup>1</sup>Department of Women's and Children's Health, Uppsala Universitet Institutionen för kvinnors och barns hälsa, Uppsala, Sweden

<sup>2</sup>Department of Medical Sciences, Uppsala University, Uppsala, Sweden

<sup>3</sup>Science for Life Laboratory, Department of Medical Cell Biology, Uppsala University, Uppsala, Sweden

<sup>4</sup>Department of Biomedical and Clinical Sciences, Linköping Universitet, Linköping, Sweden

## Correspondence to

Dr Annika Grönberg;  
annika.gronberg@kbh.uu.se

## ABSTRACT

**Introduction** The progression to insulin deficiency in type 1 diabetes is heterogenous. This study aimed to identify early characteristics associated with rapid or slow decline of beta-cell function and how it affects the clinical course.

**Research design and methods** Stimulated C-peptide was assessed by mixed meal tolerance test in 50 children (<18 years) during 2004–2017, at regular intervals for 6 years from type 1 diabetes diagnosis. 40% of the children had a rapid decline of stimulated C-peptide defined as no measurable C-peptide (<0.03 nmol/L) 30 months after diagnosis.

**Results** At diagnosis, higher frequencies of detectable glutamic acid decarboxylase antibodies (GADA) and IA-2A ( $p=0.027$ ) were associated with rapid loss of beta-cell function. C-peptide was predicted positively by age at 18 months ( $p=0.017$ ) and 30 months duration ( $p=0.038$ ). BMI SD scores (BMISDS) at diagnosis predicted higher C-peptide at diagnosis ( $p=0.006$ ), 3 months ( $p=0.002$ ), 9 months ( $p=0.005$ ), 30 months ( $p=0.022$ ), 3 years ( $p=0.009$ ), 4 years ( $p=0.016$ ) and 6 years ( $p=0.026$ ), whereas high HbA1c and blood glucose at diagnosis predicted a lower C-peptide at diagnosis ( $p<0.001$ ) for both comparisons. Both GADA and IA-2A were negative predictors of C-peptide at 9 months ( $p=0.011$ ), 18 months ( $p=0.008$ ) and 30 months ( $p<0.001$ ). Ten children had 22 events of severe hypoglycemia, and they had lower mean C-peptide at 18 months ( $p=0.025$ ), 30 months ( $p=0.008$ ) and 6 years ( $p=0.018$ ) compared with others. Seven of them had a rapid decline of C-peptide ( $p=0.030$ ), and the odds to experience a severe hypoglycemia were nearly fivefold increased (OR=4.846,  $p=0.04$ ).

**Conclusions** Low age and presence of multiple autoantibodies at diagnosis predicts a rapid loss of beta-cell function in children with type 1 diabetes. Low C-peptide is associated with an increased risk of severe hypoglycemia and higher Hemoglobin A1C. A high BMISDS at diagnosis is predictive of remaining beta-cell function during the 6 years of follow-up.

## INTRODUCTION

C-peptide can be detected in most children and adolescents at diagnosis of type 1 diabetes,<sup>1 2</sup> but the disease is characterized by a progressive decline in beta-cell function.<sup>1 3 4</sup> Several factors, including age at

## WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Low age, ketoacidosis at onset and high HbA1c are factors associated with a rapid loss of beta-cell function after onset of type 1 diabetes, while lower HbA1c during the first years and higher frequency of the HLADR3 genotype are factors associated with a long-term preservation of C-peptide.

## WHAT THIS STUDY ADDS

⇒ High BMI SD scores (BMISDS) at diagnosis predicts a slower decline of endogenous beta-cell function.  
⇒ A decline of stimulated C-peptide to undetectable levels (<0.03 nmol/L) within 30 months after diagnosis increases the risk of severe hypoglycemia (five times higher OR for severe hypoglycemia), and low C-peptide is associated with higher HbA1c and insulin requirement.  
⇒ Higher frequencies of detectable glutamic acid decarboxylase antibodies and islet cell autoantibodies islet antigen 2 at diagnosis were associated with rapid loss of beta-cell function.  
⇒ Measurements of residual C-peptide in children and adolescents during the progression of type 1 diabetes provide valuable information about clinical course.

## HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ Repeated measurements of C-peptide during follow-up of type 1 diabetes are warranted and are important for identifying the children and adolescents at highest risk of severe hypoglycemia.

## WHAT ARE THE NEW FINDINGS?

⇒ A rapid loss of C-peptide increases the risk of severe hypoglycemia.  
⇒ High BMISDS at diagnosis predicts a slower decline of beta-cell function.  
⇒ Higher frequencies of GADA and IA-2A at diagnosis were associated with rapid loss of beta-cell function.

diagnosis,<sup>5</sup> female gender,<sup>6</sup> body mass index (BMI) and ketoacidosis at diagnosis<sup>1 4 7–11</sup> as well as the occurrence of islet cell autoantibodies islet antigen 2 (IA-2A) and glutamic

acid decarboxylase antibodies (GADA)<sup>4 8 12 13</sup> are known to influence the decline of residual beta-cell function. Even though the clinical importance of residual endogenous insulin secretion is well known,<sup>13–15</sup> further knowledge on the natural course of the disease with particular identification of influencing factors possible to modify is desirable. We have made a retrospective study of type 1 diabetes subjects, whose beta-cell functions from diagnosis and onwards, as well as clinical course, are known. The aim was to identify early characteristics associated with a rapid or slow decline of beta-cell function in these children and adolescent with newly diagnosed type 1 diabetes and to investigate how the decline of C-peptide affects the clinical course of the disease during the first 6 years.

## Research design and methods

### Study subjects

Inclusion criteria to participate in the study were children and adolescent with newly diagnosed type 1 diabetes (<18 years), with start of insulin treatment at admission, with informed consent to participate and followed regularly at Crown Princess Victoria Children's Hospital, Linköping, Sweden. Children with secondary diabetes and transition to non-insulin treatment during the years of follow-up were excluded from the study. All subjects were followed as part of clinical routine with measurements of residual beta-cell function at regular intervals from diagnosis until residual beta-cell function was undetectable. In order to study the difference between individuals with rapid or slow loss of residual beta-cell function, respectively, we included 50 subjects, some with rapid loss of C-peptide (n=20), defined as having undetectable C-peptide (<0.03 nmol/L) within 30 months after diagnosis and others who had residual function up to 6 years after diagnosis (n=30).

The study subjects were born 1989–2007 and were diagnosed with type 1 diabetes during the years 2004–2017. They were at diagnosis of the disease at an age of 10.6±2.5 years, and 44% (n=22) of them were male. Diagnosis of type 1 diabetes was based on the American Diabetes Association criteria for diagnosis and classification of type 1 diabetes. At diagnosis, all study subjects had been hospitalized at the Children's Hospital and started on multiple insulin injection therapy. Thereafter, they were followed by the diabetes team at regular visits. At the age of 18 years, the study subjects were transferred to a diabetes clinic for adults. In 14 of the study subjects some data were only available 4–6 years after diagnosis.

### Data collection

Descriptive data were registered and collected from medical records and from the Swedish Childhood Diabetes Registry (SWEDIABKIDS), a national incidence and quality control register.<sup>16</sup> Data included age, sex, HbA1c, blood glucose, blood pH and C-peptide at the time of diagnosis prior to start of insulin treatment. Weight, height, HbA1c and insulin dose (units/kg body

weight/24 hours) was registered at every follow-up visit, that is, at 10 days, 1, 3, 9, 18, 24 and 30 months and 3, 4, 5 and 6 years after diagnosis. BMI (kg/m<sup>2</sup>) and BMI SD scores (BMISDS), adjusted for age and sex, were generated automatically by the SWEDIABKIDS register.<sup>17</sup> The occurrence of episodes with ketoacidosis (defined as pH<7.30) or severe hypoglycemic events (SHs) were registered at every follow-up visit. SH was defined as an event of hypoglycemia (capillary blood glucose <3.5 mmol/L) with severe cognitive impairment (including coma and convulsions) requiring assistance of another person. Data of using continuous glucose monitoring (CGM) and flash glucose monitoring ((real time) CGM/ (intermittent scanning) CGM) were registered at every follow-up visit in SWEDIABKIDS. This technology was introduced to support the treatment regime during 2016–2018. In total, 13 study subjects (26%) obtained a (rt) CGM/ (is) CGM during the study period: 2016 (n=10 new users), 2017 (n=2 new users) and 2018 (n=1 new users). An annual HbA1c average was calculated for each individual year (mean four measurements per year). However, for the first year of disease, HbA1c measurements from the first 3 months after diagnosis were excluded.

### Procedures of study and biochemical analyses

A mixed meal tolerance test (MMTT) was performed under fasting conditions in the morning; C-peptide and glucose were sampled at baseline and at 30 min intervals during the 120 min test. Study subjects were instructed not to administer short-acting insulin within 6 hours prior to the test. An MMTT was performed at 3, 9, 18 and 30 months and at 3, 4, 5 and 6 years after diagnosis.<sup>3</sup> During the first 2 years of the study (2004–2005), a standardized breakfast with 50% carbohydrates, 33% lipids and 17% proteins was ingested as the mixed meal test. From 2006, the MMTT consisted of an ingestion of a standardized liquid meal of 6 mL Sustacal/kg body weight (maximum 360 mL, 1 calorie/mL; 55% carbohydrates, 21% lipids and 24% protein). The serum samples were stored at –20°C until analysis. C-peptide concentration was measured usually within 2 weeks from sampling at the research laboratory of the Division of Pediatrics, Linköping University, using a time-resolved fluoroimmunoassay (AutoDELFLIA C-peptide kit; Wallac) with a software program (1224 MultiCalc; Wallac) for automatic calculation of values. The level of detection of the assay was 0.03 nmol/L.

MMTTs were performed until the study subjects no longer had any detectable C-peptide (defined as <0.03 nmol/L). For the following time points during the study period, undetectable C-peptide levels (<0.03 nmol/L) were assigned a numeric value of 0.01 nmol/L for statistical analysis.

Since 2005, the Crown Princess Victoria Children's Hospital participates in the nationwide cohort study 'Better Diabetes Diagnosis', which was started to monitor newly diagnosed children and adolescents with diabetes for genetic predisposition and clinical phenotypes.<sup>6</sup>

Blood samples for C-peptide concentrations, autoantibodies and HLA-DQ genotypes were collected and analyzed at diagnosis. Fasting C-peptide concentrations were also collected and analyzed 10 and 30 days after diagnosis. C-peptide was analyzed in Linköping. Autoantibodies glutamic acid decarboxylase antibodies (GADA; detection limit 5 IU/mL) and islet antigen-2 antibodies (Islet antigen-2 antibodies; detection limit 7.5 kU/L) were analyzed using two-sided ELISA test. Samples negative for ELISA IA-2A were further analyzed with a high sensitivity IA-2A radio binding assay.<sup>6</sup> HLA DQA1-DQB1 genotypes were determined with PCR.<sup>18</sup> Analyses of autoantibodies and HLA genotypes were performed at the Department of Clinical Chemistry, Skåne University Hospital, Malmö, Sweden.

The study subjects were based on the characteristics of almost half of the study subjects having undetectable C-peptide concentrations at 30-month follow-up, divided into two groups based on rate of C-peptide decline, rapid progressors (n=20), which were then defined as having undetectable C-peptide within 30 months after diagnosis and slow progressors (n=30) for which C-peptide was still detectable 3–6 years after diagnosis.

Analyses of HbA1c, pH and blood glucose were performed at the Department of Clinical Chemistry, Linköping University Hospital. The laboratory is certified by a Swedish government authority (Swedac).

From October 2010, HbA1c was analyzed according to the International Federation of Clinical Chemistry and Laboratory Medicine reference method and expressed as mmol/mol. Prior to October 2010, analyses were according to the Mono S standard expressed in per cent. In SWEDIABKIDS, analyses performed with the Mono S standard were recalculated using the expression HbA1c (IFCC; mmol/mol) = 10.45 × HbA1c (Mono S; %) – 10.62 (<http://www.ngsp.org/convert1.asp>).


### Statistics

Statistical analyses were performed using SPSS V.28.0.0. Values are given as means±SD (range). Student's t-test for independent samples was used to compare differences between two groups of normal distributed continuous data, and  $\chi^2$  test and Fisher's exact test were used for analyses of categorical data. Fisher's test was used when expected cell count was less than 5. Predictors of residual C peptide secretion as a binary outcome were compared using univariate and multivariate logistic regression analyses and analysis of variance for the main effects, expressed with ORs and 95% CI. P values <0.05 were considered statistically significant.

**Table 1** Comparison of maximum C-peptide concentration (nmol/L) mean±SD (range) in 50 children and adolescent with type 1 diabetes after stimulation with a mixed meal test during the 6 year study period in slow and rapid progressor group

Time	All subjects	Slow progressor	Rapid progressor	P value
	n=50	n=30	n=20	
Day 0	0.37±0.36 (0.07–1.86)	0.47±0.43 (0.12–1.86)	0.22±0.12 (0.07–0.60)	0.008
3 months	0.29±0.18 (0.04–0.78)	0.32±0.18 (0.08–0.75)	0.25±0.17 (0.04–0.78)	0.163
9 months	0.30±0.23 (0.04–0.95)	0.40±0.25 (0.07–0.95)	0.15±0.07 (0.04–0.29)	<0.001
18 months	0.15±0.14 (0.01–0.63)	0.22±0.15 (0.04–0.63)	0.07±0.05 (0.01–0.20)	<0.001
24 months	0.18±0.20 (0.01–0.73)	0.27±0.20 (0.07–0.73)	0.01±0.00 (0.01–0.01)	0.003
30 months	0.12±0.16 (0.01–0.71)	0.19±0.17 (0.03–0.71)	0.01±0.00 (0.01–0.01)	<0.001
3 years	0.05±0.11 (0.01–0.56)	0.13±0.16 (0.01–0.56)	0.01±0.00 (0.01–0.01)	0.048
4 years	0.07±0.10 (0.01–0.37)	0.11±0.11 (0.01–0.37)	0.01±0.00 (0.01–0.01)	<0.001
5 years	0.04±0.10 (0.01–0.45)	0.09±0.14 (0.01–0.45)	0.01±0.00 (0.01–0.01)	0.045
6 years	0.04±0.07 (0.01–0.28)	0.07±0.09 (0.01–0.28)	0.01±0.00 (0.01–0.01)	0.007

P values <0.05 were considered statistically significant.



**Figure 1** Comparison of maximum C-peptide concentration (nmol/L) (A), HbA1c (mmol/mol) (B), BMISDS (C) and daily insulin dose (U/kg/24 hours) (D) in 50 children and adolescents with type 1 diabetes from diabetes diagnosis and during 6 years of follow-up, (mean±SD), in slow (n=30) versus rapid (n=20) progressor group. BMISDS, body mass index SD score.

**RESULTS**

The C-peptide concentration was highest at diagnosis (0.37±0.36nmol/L (mean±SD) (n=42)) (table 1).


Thirty months after diagnosis, C-peptide was not measurable in the rapid progressor group (n=20)

(C-peptide <0.03 nmol/L) (table 1). There was no difference between sexes but a difference in C-peptide at every time point, except for 3 months after diagnosis, between the slow and rapid progressor groups (figure 1A) (table 1). In a linear regression analysis, C-peptide was

**Table 2** Characteristics of 50 children and adolescent at diagnosis of type 1 diagnosis (mean±SD)

	All subjects mean±SD n=50	Slow progressor mean±SD n=30	Rapid progressor mean±SD n=20	P value
Male/female	22/28	15/15	7/13	0.295
Year of diagnosis	2009±3	2008±3	2010±3	0.108
Age at diagnosis (years)	10.6±2.5	11.1±2.6	10.0±2.3	0.132
HbA1c (mmol/mol)	99±26	96±29	104±19	0.250
HbA1c (%)	11.2±4.5	10.9±4.8	11.7±3.9	0.250
Blood glucose (mmol/L)	28.8±14.9	25.4±9.4	33.6±19.7	0.068
BMISDS	-0.07±1.26	0.08±1.40	-0.29±1.02	0.339
pH	7.35±0.09	7.36±0.07	7.32±0.11	0.129
Ketoacidosis (pH<7.0)	7/48	2/28	5/20	0.084
GADA detectable	33/42	18/25	15/17	0.208
IA-2A detectable	32/42	17/25	15/17	0.131
GADA and IA-2A	29/42	14/25	15/17	0.027
HLA-risk profile	29/39	16/24	13/15	0.164

Differences in the slow versus rapid progressor group. P value <0.05 was considered as statistically significant. BMISDS, body mass index SD scores; GADA, glutamic acid decarboxylase antibodies; IA-2A, islet antigen-2 antibodies.



**Figure 2** Comparison of maximum C-peptide concentration (nmol/L) after stimulation with a mixed meal test in 50 children with versus no severe hypoglycemia events during the 6-year study period (mean±SD): at 18 months ( $0.09\pm 0.07$  nmol/L) versus ( $0.17\pm 0.15$  nmol/L) ( $p=0.025$ ), at 30 months ( $0.04\pm 0.07$  nmol/L) versus ( $0.14\pm 0.18$  nmol/L) ( $p=0.008$ ) and at 6 years ( $0.02\pm 0.03$  nmol/L) versus ( $0.10\pm 0.10$  nmol/L) ( $p=0.018$ );  $p$  values  $<0.05$  were considered statistically significant.

predicted positively by age at diagnosis at 18 months ( $r=0.123$ ,  $p=0.017$ ) and 30 months ( $r=0.087$ ,  $p=0.038$ ) and also positively predicted by BMISDS at diagnosis ( $r=0.195$ ,  $p=0.006$ ), 3 months ( $r=0.203$ ,  $p=0.002$ ) (online supplemental figure S1), 9 months ( $r=0.174$ ,  $p=0.005$ ), 30 months ( $r=0.118$ ,  $p=0.022$ ), 3 years ( $r=0.252$ ,  $p=0.009$ ), 4 years ( $r=0.136$ ,  $p=0.016$ ) and 6 years ( $r=0.134$ ,  $p=0.026$ ). High HbA1c at diagnosis predicted a lower C-peptide at diagnosis ( $r=-0.300$ ,  $p<0.001$ ) and at 10 days ( $r=-0.126$ ,  $p=0.015$ ). Blood glucose, but not sex, pH or ketoacidosis, at diagnosis was also negatively associated with C-peptide at diagnosis ( $r=-0.301$ ,  $p<0.001$ ). The average calculated HbA1c during the first and second year was negatively associated with C-peptide at 30 months ( $r=-0.088$ ,  $p=0.039$ ) and ( $r=-0.094$ ,  $p=0.030$ ) for first and second year, respectively). Detectable GADA at diagnosis was a negative predictor for C-peptide, with lower C-peptide at 18 months ( $p=0.025$ ) and 30 months ( $p=0.042$ ). Detectable IA-2A was also a negative predictor for C-peptide at 9 months ( $p=0.005$ ), 18 months ( $p=0.003$ ) and 30 months ( $p=0.001$ ). Both GADA and IA-2A were negative predictors of C-peptide at 9 months ( $p=0.011$ ), 18 months ( $p=0.008$ ) and 30 months ( $p<0.001$ ).

When comparing clinical features in the slow versus rapid progressor groups during the study period (figure 1B–D), there was no difference at diabetes diagnosis in sex, age, HbA1c, blood glucose, BMISDS, pH, ketoacidosis or HLA risk profile (HLADQ genotypes DRB1\*04-DQA103-DQB10302) but a higher frequency of both detectable GADA and IA-2A in rapid progressors ( $p=0.027$ ) (table 2).

We observed no difference between slow and rapid progressors in mean HbA1c for each individual year or in BMISDS (figure 1B,C). Daily insulin doses were lower in slow progressors at 24-month follow-up ( $0.82\pm 0.32$  vs  $1.01\pm 0.31$  U/kg/24 hours,  $p=0.044$ ) (figure 1D). IDA1C was lower in the slow progressors at 9-month follow-up ( $8.8\pm 1.2$  vs  $9.9\pm 1.7$ ,  $p=0.019$ ), but there was no difference

in number of study subjects in partial remission. There were no episodes of diabetic ketoacidosis during the study period.

Ten children (six girls and four boys) experienced 22 events of SH during the study period: 3/30 (10%) from the slow progressor group and 7/20 (35%) from the rapid progressor group ( $p=0.030$ ). During the first 3 years after diagnosis, six study subjects experienced an event of SH, and all of these subjects were rapid progressors ( $p=0.002$ ). The study subjects in the rapid progressor group had altogether 18 events of SH during the study period. There was a lower mean C-peptide concentration in those with SH ( $n=10$ ) versus not any SH ( $n=40$ ) at 18 months ( $p=0.025$ ), 30 months ( $p=0.008$ ) and 6 years ( $p=0.018$ ) after the diagnosis (figure 2). When comparing slow and rapid progressors, low C-peptide concentrations at 18 months ( $0.22\pm 0.15$  nmol/L) versus ( $0.07\pm 0.05$  nmol/L) ( $p<0.001$ ) and 30 months ( $0.19\pm 0.17$  nmol/L) versus ( $0.01\pm 0.00$  nmol/L) ( $p<0.001$ ) (table 1) were associated with SH.

When applying a binary logistic regression analysis, expressed with ORs, there was no difference for any of clinical characteristics between the progressor groups at diagnosis. However, there was a difference in OR in the rapid progressor group for detectable GADA and IA-2A (OR: 5.893,  $p=0.038$ ) and for lower C-peptide at diagnosis (OR=0.001,  $p=0.037$ ). The study subjects with rapid progress had also nearly five times higher odds to experience a SH during the study period (OR=4.846,  $p=0.04$ ).

## DISCUSSION

The present study depicts the natural course of beta-cell function and clinical outcome in 50 children and adolescents newly diagnosed with type 1 diabetes during the first 6 years of follow-up. By comparing early clinical characteristics associated with a rapid or slow decline of stimulated C-peptide, we have investigated the impact of beta-cell function for the clinical course of type 1 diabetes.

Based on stimulated C-peptide levels and clinical descriptive data, we have identified that low age at diagnosis and presence of multiple autoantibodies can predict a rapid decline of beta-cell function, which affects the clinical course by increasing the risk and frequency of severe hypoglycemia. A high BMISDS at diagnosis is the most important predictor for maintaining beta-cell function over time. Rapid progressors, that is, C-peptide was undetectable ( $<0.03$  nmol/L) within 30 months after diagnosis, had a higher frequency of detectable autoantibodies GADA and IA-2A at diagnosis, but there was no difference in other clinical characteristics. Occurrence of multiple autoantibodies indicate an autoimmune progressive destruction of beta-cells and is a well-known predictor of rapid decline in C-peptide secretion.<sup>4 8 12 19</sup> The present results show that autoantibodies at diagnosis also are a predictor of rapid beta-cell functional loss in all subjects from the ninth month after diagnosis.

HbA1c did not differ between rapid and slow progressors, but study subjects with less C-peptide had higher average calculated HbA1c first 2 years and slow progressors needed less insulin. Thus, even if it seems to be possible to achieve good glycemic control with use of intensive insulin treatment in spite of low residual C-peptide,<sup>20</sup> even low own residual beta cell function improves metabolic control.

Furthermore, loss of residual C-peptide is known to be an important predictor for short-term complications such as severe hypoglycemia, which we confirm in this study, and ketoacidosis<sup>7 11 14</sup> as well as for long-term diabetes complications.<sup>14 21–23</sup>

We observed no episodes of ketoacidosis, but 22 events of SH during the 6 years of follow-up. SH was only experienced by study subjects with rapid loss of C-peptide during the first 3 years after diagnosis. These findings are well in line with previous evidence of a protective effect even of low levels of residual C-peptide on the frequency of SH.<sup>14 24</sup> We found that the slow progressors even with low C-peptide (>0.03 nmol/L) still had reduced risk of SH during the follow-up. The study subjects that experienced an SH during the 6 years of follow-up had lower levels of C-peptide at 18 and 30 months after diagnosis.

Subjects with a rapid decline of C-peptide presented with lower levels of C-peptide already at diagnosis and for every time point during the follow-up period, except during the remission period 3 months after diagnosis. These results are indicative of a lower beta-cell function in rapid progressors already at diagnosis, which could be related to a more aggressive immunological processes or a delayed diagnosis. At 3 months, the glycemic control was improved in both groups with the lowest average HbA1c and insulin requirements during the whole study period. The remaining beta-cells might temporarily recover, a clinical partial remission phase (honeymoon) starting after exogenous insulin treatment is initiated at diagnosis with improvement of the metabolic disturbance and peripheral insulin sensitivity.<sup>25</sup> The duration of the remission period depends at least partly on the recovery of the beta-cell function, with a shorter period of C-peptide secretion in the rapid progressor, with a maximum stimulated C-peptide 3 months after diagnosis versus 9 months in slow progressor.

In this study, BMISDS was an important predictor of higher C-peptide levels at diagnosis and during the 6 years of follow-up. BMISDS at baseline has previously in several, but not all, studies<sup>26</sup> been reported as an important predictor of residual C-peptide and to be associated with a decline of C-peptide during the first years after diagnosis.<sup>4 9 10 27 28</sup> Increased linear growth and overweight, as well as progression of insulin resistance, before diagnosis<sup>29–32</sup> may be an accelerator for symptoms at an early stage of type 1 diabetes, with higher BMISDS and lower HbA1c then associated with greater remaining beta-cell function.

Lower average calculated HbA1c in study subjects during the first and second year predicted higher residual

C-peptide 30 months after diagnosis. The residual C-peptide had a clinically beneficial effect on the frequency of SH, reflecting remaining endogenous insulin production during the first 30 months following diagnosis.

Sex did not emerge as predictor for residual C-peptide as previously shown.<sup>10 28 33</sup>

There was no age-dependent differences in C-peptide at diagnosis, but younger age at diagnosis predicted lower C-peptide levels 18 and 30 months after diagnosis. This is likely explained by a more aggressive form of the disease or an overall lower beta-cell mass in young children. Age at diagnosis was one of the most important predictors of residual C-peptide, as also shown previously by most<sup>1 4 5 21 24 27 34</sup> but not all studies.<sup>12</sup> HLA risk profile was not associated with the progression of the C-peptide decline during the study period, which suggests that the progression of beta-cell destruction are less affected by genetic factors. The HLA risk profile may be more important as a risk for developing type 1 diabetes.<sup>18</sup>

A strength of this study is the use of an MMTT analyzed with the same C-peptide assay during a longitudinal follow-up in children and adolescent, treated at a single pediatric diabetes center with a uniform treatment program and guidelines for pediatric diabetes with an ambitious treatment target.<sup>35</sup> There is a limitation in that some analyses were not available when the study subjects were transferred to a diabetic clinic for adults at the age of 18 years. The registered report of the frequency of hypoglycemia and ketoacidosis might have been under-reported.

## CONCLUSION

We conclude that low age at diagnosis and the presence of multiple autoantibodies predict a rapid decline of beta-cell function. Loss of C-peptide is associated with an increased risk and higher frequency of severe hypoglycemic events and less good blood glucose control. A higher BMISDS at diagnosis is predictive of a slow decline of beta-cell function.

**Acknowledgements** Thanks to the children and adolescents who participated in the study, and also the nurses who took all the samples are gratefully acknowledged.

**Contributors** JL initiated and designed the study. JL coordinated and recruited the study subjects. JL, AG and DE collected the data, and AG created the database, analyzed the data and contributed with statistical support and is also responsible for the overall content as the guarantor. AG drafted the first manuscript, and all authors edited and approved the final version of the manuscript for publication.

**Funding** This work was supported by the Swedish Child Diabetes Foundation (Barndiabetesfonden), Gillberggska stiftelsen, Sven Jerrings Foundation, Stiftelsen Samariten, Ingemar Swenne Foundation, Vetenskapsrådet, Diabetesfonden and Uppsala University.

**Competing interests** None declared.

**Patient consent for publication** Not applicable.

**Ethics approval** The Research Ethics Committee of the Faculty of Health Sciences, Linköping University approved the study with reference number 03-244. Participants gave informed consent to participate in the study before taking part.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data are available on reasonable request.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.


#### ORCID iDs

Annika Grönberg <http://orcid.org/0000-0002-8502-8568>

Daniel Espes <http://orcid.org/0000-0001-8843-7941>

#### REFERENCES

- Ludvigsson J, Heding LG. C-Peptide in children with juvenile diabetes. *Diabetologia* 1976;12:627–30.
- Ludvigsson J, Heding LG. Beta-cell function in children with diabetes. *Diabetes* 1978;27 Suppl 1:230–4.
- Nordwall M, Ludvigsson J. Clinical manifestations and beta cell function in Swedish diabetic children have remained unchanged during the last 25 years. *Diabetes Metab Res Rev* 2008;24:472–9.
- Ludvigsson J, Carlsson A, Deli A, *et al*. Decline of C-peptide during the first year after diagnosis of type 1 diabetes in children and adolescents. *Diabetes Res Clin Pract* 2013;100:203–9.
- Barker A, Lauria A, Schloot N, *et al*. Age-dependent decline of  $\beta$ -cell function in type 1 diabetes after diagnosis: a multi-centre longitudinal study. *Diabetes Obes Metab* 2014;16:262–7.
- Persson M, Becker C, Elding Larsson H, *et al*. The Better Diabetes Diagnosis (BDD) study - A review of a nationwide prospective cohort study in Sweden. *Diabetes Res Clin Pract* 2018;140:236–44.
- Madsbad S, Alberti KG, Binder C, *et al*. Role of residual insulin secretion in protecting against ketoacidosis in insulin-dependent diabetes. *Br Med J* 1979;2:1257–9.
- Mortensen HB, Swift PGF, Holl RW, *et al*. Multinational study in children and adolescents with newly diagnosed type 1 diabetes: association of age, ketoacidosis, HLA status, and autoantibodies on residual beta-cell function and glycemic control 12 months after diagnosis. *Pediatr Diabetes* 2010;11:218–26.
- Lauria A, Barker A, Schloot N, *et al*. BMI is an important driver of  $\beta$ -cell loss in type 1 diabetes upon diagnosis in 10 to 18-year-old children. *Eur J Endocrinol* 2015;172:107–13.
- Szypowska A, Groele L, Wysocka-Mincewicz M, *et al*. Factors associated with preservation of C-peptide levels at the diagnosis of type 1 diabetes. *J Diabetes Complications* 2018;32:570–4.
- Komulainen J, Lounamaa R, Knip M, *et al*. Ketoacidosis at the diagnosis of type 1 (insulin dependent) diabetes mellitus is related to poor residual beta cell function. Childhood diabetes in Finland Study Group. *Arch Dis Child* 1996;75:410–5.
- Ludvigsson J, Hellström S. Autoantibodies in relation to residual insulin secretion in children with IDDM. *Diabetes Res Clin Pract* 1997;35:81–9.
- Ludvigsson J. The clinical potential of low-level C-peptide secretion. *Expert Rev Mol Diagn* 2016;16:933–40.
- Steffes MW, Sibley S, Jackson M, *et al*. Beta-cell function and the development of diabetes-related complications in the diabetes control and complications trial. *Diabetes Care* 2003;26:832–6.
- Lachin JM, McGee P, Palmer JP, *et al*. Impact of C-peptide preservation on metabolic and clinical outcomes in the diabetes control and complications trial. *Diabetes* 2014;63:739–48.
- Hanberger L, Samuelsson U, Lindblad B, *et al*. A1C in children and adolescents with diabetes in relation to certain clinical parameters: the Swedish childhood diabetes registry SWEDIABKIDS. *Diabetes Care* 2008;31:927–9.
- Karlberg J, Luo ZC, Albertsson-Wikland K. Body mass index reference values (mean and SD) for Swedish children. *Acta Paediatr* 2001;90:1427–34.
- Ilonen J, Kiviniemi M, Lempainen J, *et al*. Genetic susceptibility to type 1 diabetes in childhood - estimation of HLA class II associated disease risk and class II effect in various phases of islet autoimmunity. *Pediatr Diabetes* 2016;17 Suppl 22:8–16.
- Steck AK, Vehik K, Bonifacio E, *et al*. Predictors of progression from the appearance of islet autoantibodies to early childhood diabetes: the environmental determinants of diabetes in the young (TEDDY). *Diabetes Care* 2015;38:808–13.
- Anderzén J, Hermann JM, Samuelsson U, *et al*. International benchmarking in type 1 diabetes: large difference in childhood HbA1c between eight high-income countries but similar rise during adolescence-A quality registry study. *Pediatr Diabetes* 2020;21:621–7.
- Kuhreiter WM, Washer SLL, Hsu E, *et al*. Low levels of C-peptide have clinical significance for established type 1 diabetes. *Diabet Med* 2015;32:1346–53.
- Effect of intensive therapy on residual beta-cell function in patients with type 1 diabetes in the diabetes control and complications trial. A randomized, controlled trial. The diabetes control and complications trial research group. *Ann Intern Med* 1998;128:517–23.
- Luppi P, Cifarelli V, Wahren J. C-Peptide and long-term complications of diabetes. *Pediatr Diabetes* 2011;12:276–92.
- Sørensen JS, Johannesen J, Pociot F, *et al*. Residual  $\beta$ -cell function 3–6 years after onset of type 1 diabetes reduces risk of severe hypoglycemia in children and adolescents. *Diabetes Care* 2013;36:3454–9.
- DiMeglio LA, Cheng P, Beck RW, *et al*. Changes in beta cell function during the proximate post-diagnosis period in persons with type 1 diabetes. *Pediatr Diabetes* 2016;17:237–43.
- Dabelea D, Mayer-Davis EJ, Andrews JS, *et al*. Clinical evolution of beta cell function in youth with diabetes: the search for diabetes in youth study. *Diabetologia* 2012;55:3359–68.
- Greenbaum CJ, Beam CA, Boulware D, *et al*. Fall in C-peptide during first 2 years from diagnosis: evidence of at least two distinct phases from composite type 1 diabetes TrialNet data. *Diabetes* 2012;61:2066–73.
- Grönberg A, Espes D, Carlsson P-O. Better HbA1c during the first years after diagnosis of type 1 diabetes is associated with residual C peptide 10 years later. *BMJ Open Diabetes Res Care* 2020;8:e000819.
- Hyppönen E, Virtanen SM, Kenward MG, *et al*. Obesity, increased linear growth, and risk of type 1 diabetes in children. *Diabetes Care* 2000;23:1755–60.
- Ljungkrantz M, Ludvigsson J, Samuelsson U. Type 1 diabetes: increased height and weight gains in early childhood. *Pediatr Diabetes* 2008;9:50–6.
- Bonfig W, Kapellen T, Dost A, *et al*. Growth in children and adolescents with type 1 diabetes. *J Pediatr* 2012;160:900–3.
- Knerr I, Wolf J, Reinehr T, *et al*. The 'accelerator hypothesis': relationship between weight, height, body mass index and age at diagnosis in a large cohort of 9,248 German and Austrian children with type 1 diabetes mellitus. *Diabetologia* 2005;48:2501–4.
- Samuelsson U, Lindblad B, Carlsson A, *et al*. Residual beta cell function at diagnosis of type 1 diabetes in children and adolescents varies with gender and season. *Diabetes Metab Res Rev* 2013;29:85–9.
- Steck A, Liu X, Krischer J, *et al*. 212-OR: factors associated with decline of C-peptide in a cohort of young children diagnosed with type 1 diabetes. *Diabetes* 2019;68.
- Swift PGF, Skinner TC, de Beaufort CE, *et al*. Target setting in intensive insulin management is associated with metabolic control: the Hvidoere childhood diabetes Study Group centre differences study 2005. *Pediatr Diabetes* 2010;11:271–8.



### Supplemental figure 1

Correlations between baseline BMI SDS and maximum C-peptide concentration (nmol/L) 3 months after diagnosis ( $p=0.002$ ) in 50 children and adolescents with type 1 diabetes. Body mass index standard deviation score (BMISDS).