




Higher burden of cardiometabolic and socioeconomic risk factors in women with type 2 diabetes: an analysis of the Glycemic Reduction Approaches in Diabetes (GRADE) baseline cohort

Barbara I Gulanski ^{1,2}, Nicole M Butera,³ Heidi Krause-Steinrauf,³ Judith H Lichtman,⁴ Tasma Harindhanavudhi,⁵ Jennifer B Green,⁶ Colleen E Suratt,³ Hiba AbouAssi,⁶ Cyrus Desouza ⁷, Andrew J Ahmann,⁸ Deborah J Wexler,⁹ Vanita R Aroda ^{10,11} GRADE Research Group³

To cite: Gulanski BI, Butera NM, Krause-Steinrauf H, *et al*. Higher burden of cardiometabolic and socioeconomic risk factors in women with type 2 diabetes: an analysis of the Glycemic Reduction Approaches in Diabetes (GRADE) baseline cohort. *BMJ Open Diab Res Care* 2023;**11**:e003159. doi:10.1136/bmjdr-2022-003159

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjdr-2022-003159>).

DJW and VRA are senior co-authors.

Received 3 October 2022
Accepted 18 February 2023



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Barbara I Gulanski;
Barbara.Gulanski@yale.edu

ABSTRACT

Introduction Type 2 diabetes mellitus (T2DM) is a powerful risk factor for cardiovascular disease (CVD), conferring a greater relative risk in women than men. We sought to examine sex differences in cardiometabolic risk factors and management in the contemporary cohort represented by the Glycemia Reduction Approaches in Diabetes: A Comparative Effectiveness Study (GRADE).

Research design and methods GRADE enrolled 5047 participants (1837 women, 3210 men) with T2DM on metformin monotherapy at baseline. The current report is a cross-sectional analysis of baseline data collected July 2013 to August 2017.

Results Compared with men, women had a higher mean body mass index (BMI), greater prevalence of severe obesity (BMI \geq 40 kg/m²), higher mean LDL cholesterol, greater prevalence of low HDL cholesterol, and were less likely to receive statin treatment and achieve target LDL, with a generally greater prevalence of these risk factors in younger women. Women with hypertension were equally likely to achieve blood pressure targets as men; however, women were less likely to receive ACE inhibitors or angiotensin receptor blockers. Women were more likely to be divorced, separated or widowed, and had fewer years of education and lower incomes.

Conclusions This contemporary cohort demonstrates that women with T2DM continue to have a greater burden of cardiometabolic and socioeconomic risk factors than men, particularly younger women. Attention to these persisting disparities is needed to reduce the burden of CVD in women.

Trial registration number ClinicalTrials.gov (NCT01794143)

INTRODUCTION

Cardiovascular disease (CVD) is the primary cause of mortality among women, and women with type 2 diabetes mellitus (T2DM) have a threefold higher relative risk for cardiovascular (CV) death than women without

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Decades of research have shown that women with type 2 diabetes mellitus (T2DM) are less likely to receive evidence-based care for cardiometabolic risk factor management

WHAT THIS STUDY ADDS

⇒ In this contemporary cohort, disparities persist in the evidence-based management of traditional cardiovascular risk factor of dyslipidemia, and substantial differences in non-traditional adverse socioeconomic factors are evident, which may be more relevant for addressing cardiac risk in this population.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE, OR POLICY

⇒ There is a need to optimize care strategies to reduce the heightened risk factor burden and treatment gaps for women with T2DM.

diabetes, even after controlling for traditional cardiac risk factors including blood pressure (BP), cholesterol, body mass index (BMI) and smoking.^{1–3} In comparison, diabetes doubles the CV mortality risk among men.¹

The higher relative risk of CVD for women with diabetes has been attributed to multiple factors, including a higher baseline burden of traditional cardiometabolic risk factors (eg, hypertension, obesity, diabetes), novel risk factors (eg, inflammation, prothrombotic phenotype), non-traditional risk factors (eg, depression, lower socioeconomic status (SES)),^{4–10} and less aggressive management of traditional cardiac risk factors for primary prevention.^{11–14} Factors that contribute to worse outcomes compared with men include sex differences in cardiac symptom

presentation,^{15 16} lower use of cardiac procedures and evidence-based treatment for secondary prevention,^{17–22} and more postprocedural complications.^{23–25}

The Glycemia Reduction Approaches in Diabetes: A Comparative Effectiveness Study (GRADE) is a prospective, randomized controlled trial comparing four glucose-lowering medications on metabolic and patient-centered outcomes in adults with T2DM.²⁶ Unlike earlier studies which examined diabetes as a simple binary variable (yes/no) without accounting for duration of disease, degree of glycemic control, or specific glucose-lowering treatment, GRADE offers a unique opportunity to examine sex differences in a large contemporary group of individuals with an early clinical stage of T2DM who have undergone detailed clinical and metabolic phenotyping at baseline.

Our aim was to examine sex differences in baseline traditional and socioeconomic risk factors within this cohort, and to determine whether there were sex differences in the provision of evidence-based treatment for prevention of CV complications. Identified differences in this representative population may provide meaningful insight on current gaps that contribute to persistent disparities in CVD burden for women with T2DM.

RESEARCH DESIGN AND METHODS

Study design, setting, and participants

The design of the GRADE Study has been previously described (GRADE protocol).²⁶ In brief, GRADE is a parallel-group, randomized clinical trial that compares the effectiveness of four major classes of glucose-lowering medications in individuals with T2DM treated with metformin on glycemic, metabolic, and patient-centered outcomes. Participants provided written informed consent.

Recruitment occurred between 2013 and 2017 and follow-up continued through spring 2021. The present analysis is restricted to data collected during the screening/baseline period of the GRADE Study. The study was conducted at 36 funded clinical centers, including nine additional subsites, across the USA. Sites included academic hospitals, integrated health systems, and Veterans Affairs (VA) medical centers.

Eligibility criteria included individuals with a history of T2DM for <10 years in duration and ≥30 years of age at diagnosis (≥20 years of age at diagnosis for Native Americans) who were treated with metformin, but no other glucose-lowering medications, and who had a glycated hemoglobin (HbA1c) level of 6.8%–8.5% (51–69 mmol/mol) at randomization, measured at a central laboratory, after metformin therapy was maximized to a dose of at least 1000 mg daily and up to 2000 mg daily, as tolerated, during the run-in period. Exclusion criteria included individuals with type 1 diabetes mellitus, chronic kidney disease with an estimated glomerular filtration rate (eGFR) <30 mL/min/1.73 m², congestive heart failure (NY State Heart Association Functional Classification ≥III), history of a major CV event (myocardial infarction (MI), stroke, or vascular procedure such as coronary artery or peripheral bypass grafting, stent

placements (peripheral or coronary) or angioplasty) within the preceding year, new diagnosis or treatment for cancer within the prior 5 years, planned pregnancy in women of childbearing potential, planned major surgery, and history of pancreatitis.

Demographic and clinical variables were collected by self-report including age, race, ethnicity, sex, diabetes duration (years), smoking history, marital status, years of education, employment, and income. Physical measurements including height (m), weight (kg), waist circumference (cm), hip circumference (cm), and BP (mm Hg) were obtained by trained clinical research staff; per protocol, each physical measurement was taken in duplicate and averaged. BMI (kg/m²) and waist-to-hip ratio were calculated based on these physical measurements. Concomitant medications known to affect glycemia or CV risk were collected by self-report for all participants with the exception of querying for the use of medications for depression and anxiety; this question was added after study initiation and collected at baseline for 2494 participants only.

Hypertension was considered present if it was reported by the participant, the participant reported taking an antihypertensive medication, measured systolic BP was ≥130 mm Hg, or measured diastolic BP was ≥80 mm Hg. Hyperlipidemia was present if it was reported by the participant, the participant reported taking a lipid-lowering medication, or measured low-density lipoprotein cholesterol (LDL-C) was ≥100 mg/dL. Sex-specific thresholds were used to define increased waist circumference, ≥88 cm for women and ≥102 cm for men (for Asian Americans, ≥80 cm for women and ≥90 cm for men). Sex-specific thresholds were also used to define abnormal high-density lipoprotein cholesterol (HDL-C) levels, specifically HDL-C <50 mg/dL for women and <40 mg/dL for men. Metabolic syndrome was defined by the presence of at least three of the following criteria: abdominal obesity, high triglycerides (≥150 mg/dL, or lipid-lowering medication), low HDL-C; <50 mg/dL for women, <40 mg/dL for men, or lipid-lowering medication), hypertension (systolic BP ≥130 mm Hg, diastolic BP ≥80 mm Hg, or antihypertensive medication), or high fasting glucose (≥100 mg/dL, or glycemia-lowering medication);²⁷ since metformin was a criterion for enrollment in GRADE, all participants met at least 1 of the metabolic syndrome criteria (ie, using glycemia-lowering medication).

Laboratory data were analyzed at the Advanced Research and Diagnostic Laboratory at the University of Minnesota, using standardized laboratory procedures, including HbA1c (standardized per National Glycohemoglobin Standardization Program protocol). LDL-C was calculated using the Friedewald equation if the triglyceride concentration was <400 mg/dL (LDL-C=TC [total cholesterol]–HDL-C–triglyceride/5.0).²⁸ HDL-C was measured using the Roche HDL-Cholesterol direct method (Roche Diagnostics, Indianapolis, Indiana, USA). Triglycerides were measured using Triglyceride GB (glycerol-blanked) reagent (Roche Diagnostics, Indianapolis, Indiana, USA).

Statistical analyses

Analyses were presented by sex (female/male). Descriptive statistics included means \pm SD and nominal *t* tests to explore differences between men and women for all continuous variables, and proportions and nominal Pearson's χ^2 tests to explore differences between men and women for all categorical variables.

The proportion of participants receiving clinically indicated treatment for hypertension (ie, BP medications among those with hypertension), hyperlipidemia (ie, statin treatment among those with hyperlipidemia) and those achieving target levels of LDL-C $<$ 100mg/dL ($<$ 2.59mmol/L) and BP $<$ 140/90 mm Hg and $<$ 130/80 mm Hg were presented by sex. Differences between men and women for each of these variables were tested using Pearson's χ^2 tests, and a Holm adjustment was applied to the *p* values from these tests to control the type I error rate in this set of analyses.

Exploratory analyses were conducted to assess whether sex differences in cardiometabolic risk factors differed by sociodemographic factors, such as age ($<$ 45 years, 45–54 years, 55–64 years, 65+years), income ($<$ US\$10 000, US\$10 000 to $<$ US\$50 000, US\$50 000 to $<$ US\$75 000, \geq US\$75 000), education ($<$ high school, high school graduate/GED, some college, college graduate or more), race (white, black or African-American, other or more than one race), and ethnicity (Hispanic, non-Hispanic). Means or proportions for each cardiometabolic risk factor were calculated for men and women within each category of the sociodemographic factor. Exploratory analyses to assess sex differences in the cardiometabolic factor as a function of the sociodemographic factor and the interaction with sex were conducted using the following described procedures. Linear (for continuous risk factors) or logistic (for binary risk factors) regression models were fit for each risk factor with covariates for sex, the sociodemographic factor, and the interaction term. A nominal joint Wald test of the interaction term was then conducted to explore whether sex differences in that risk factor varied by that sociodemographic factor (ie, to test for homogeneity of sex differences in the cardiometabolic risk factor across the sociodemographic factor).

Medical centers affiliated with VA accounted for 10 of the 45 total sites across the 36 funded clinical centers in GRADE. Since the patient population within VA Medical Centers was expected to differ from a more general patient population,²⁹ a sensitivity analysis was conducted that repeated the previous analyses stratified by whether participants were enrolled at a VA study site, allowing for comparison of sex differences in sociodemographic factors and cardiometabolic risk factors between VA study sites and non-VA study sites.

RESULTS

A total of 5047 individuals with T2DM participated in the GRADE Study, including 1837 women and 3210 men, ranging in age from 22 years to 85 years. The average ages for women and men were 55.4 years and 58.2 years, respectively. The percentage of participants age 65 years

or older was almost twice as high in men as in women (men 28.8% vs women 15.6%). Compared with men, women in this cohort were more likely to self-report as black, Native American, or Hispanic. The average duration of diabetes was approximately 4 years for both sexes (table 1).

Sex differences in cardiometabolic risk factors

At baseline, the average HbA1c for both women and men was 7.5% (58.3 mmol/mol). Compared with men, women had higher BMI across all demographic and socioeconomic groups (table 1; online supplemental table 1), and more women met criteria for class III obesity (BMI \geq 40 kg/m², women 23.6% vs men 14.5%). Women were more likely to meet sex-specific criteria for increased waist circumference than men (women 92.9% vs men 79%; table 1). Overall, women had lower systolic BP and diastolic BP, but the absolute differences were small (table 1). The overwhelming majority of all study participants ($>$ 90%) met criteria for metabolic syndrome.

Lipid levels varied by sex and race. Women had higher mean LDL-C levels than men (table 1). The sex differences in LDL-C levels were greater for white participants (white men 75.6mg/dL (1.96mmol/L) vs white women 84.5mg/dL (2.19mmol/L)); however, black participants had higher LDL-C levels overall (black men 91.3mg/dL (2.36mmol/L) vs black women 94.9mg/dL (2.46mmol/L)) (online supplemental table 2). Women consistently had a greater prevalence of low HDL-C than men across all demographic and socioeconomic groups, based on sex-specific criteria. Sex differences in low HDL-C levels varied across racial groups; black participants had a lower prevalence of low HDL-C (men 32.8% vs women 50.8%) than white participants (men 51.4% vs women 69%) (online supplemental tables 3 and 4). Men, on the other hand, had consistently higher triglyceride levels than women.

Sex differences in evidence-based therapies and treatment targets

Women with dyslipidemia were less likely to receive statin treatment than men overall (table 2). Stratified analyses showed that women treated with a statin had higher LDL-C levels and were less likely to achieve target LDL-C $<$ 100mg/dL ($<$ 2.59mmol/L) than men across demographic and socioeconomic groups (figure 1; online supplemental tables 2 and 5). Overall, younger women, black and Hispanic women, and women with lower education and income had higher LDL-C levels and lower prevalence of LDL-C $<$ 100mg/dL ($<$ 2.59mmol/L) while being treated with statin therapy (online supplemental table 2).

In contrast, there were no sex differences in the percentage of individuals with hypertension receiving BP-lowering medications and achieving BP targets; however, women with hypertension were less likely to receive angiotensin converting enzyme inhibitors (ACE

Table 1 Baseline characteristics by sex

	Overall (n=5047)	Men (n=3210)	Women (n=1837)	P value
Age at baseline visit (years)	57.2±10.0	58.2±10.1	55.4±9.6	<0.001
Categorized age				<0.001
Age <45 years	623 (12.3%)	346 (10.8%)	277 (15.1%)	
Age 45–54 years	1436 (28.5%)	856 (26.7%)	580 (31.6%)	
Age 55–64 years	1778 (35.2%)	1085 (33.8%)	693 (37.7%)	
Age ≥65 years	1210 (24.0%)	923 (28.8%)	287 (15.6%)	
Diabetes duration (years), mean	4.0±2.8	4.1±2.8	4.0±2.8	0.258
<i>Demographics</i>				
Race				<0.001
American Indian/Alaska Native	137 (2.7%)	56 (1.7%)	81 (4.4%)	
Asian	182 (3.6%)	140 (4.4%)	42 (2.3%)	
Hawaiian/Pacific Islander	28 (0.6%)	20 (0.6%)	8 (0.4%)	
Black or African-American	1000 (19.8%)	491 (15.3%)	509 (27.7%)	
White	3314 (65.7%)	2288 (71.3%)	1026 (55.9%)	
Other/multiple	319 (6.3%)	175 (5.5%)	144 (7.8%)	
Unknown/not reported	67 (1.3%)	40 (1.2%)	27 (1.5%)	
Ethnicity				<0.001
Hispanic/Latino	929 (18.4%)	502 (15.6%)	427 (23.2%)	
Not Hispanic/Latino	4077 (80.8%)	2685 (83.6%)	1392 (75.8%)	
Unknown	41 (0.8%)	23 (0.7%)	18 (1.0%)	
Marital status				<0.001
Married	2918 (57.8%)	2061 (64.2%)	857 (46.7%)	
Living with partner	288 (5.7%)	179 (5.6%)	109 (5.9%)	
Divorced/separated/widowed	1233 (24.4%)	631 (19.7%)	602 (32.8%)	
Single	606 (12.0%)	339 (10.6%)	267 (14.6%)	
Years of education				<0.001
Some high school	364 (7.2%)	174 (5.4%)	190 (10.3%)	
High school graduate/GED	1039 (20.6%)	632 (19.7%)	407 (22.2%)	
Some college	1463 (29.0%)	951 (29.6%)	512 (27.9%)	
College graduate or more	2180 (43.2%)	1453 (45.3%)	727 (39.6%)	
Employment				<0.001
Employed full-time or part-time, or seasonally	3004 (59.6%)	1875 (58.4%)	1129 (61.5%)	
Retired	1203 (23.9%)	902 (28.1%)	301 (16.4%)	
Full-time homemaker	152 (3.0%)	7 (0.2%)	145 (7.9%)	
Not employed	380 (7.5%)	229 (7.1%)	151 (8.2%)	
Never worked or disabled	145 (2.9%)	97 (3.0%)	48 (2.6%)	
Student or other	160 (3.2%)	98 (3.1%)	62 (3.4%)	
Income				<0.001
<US\$10 000	296 (5.9%)	146 (4.5%)	150 (8.2%)	
US\$10 000 to <US\$50 000	1942 (38.5%)	1162 (36.2%)	780 (42.5%)	
US\$50 000 to <US\$75 000	788 (15.6%)	529 (16.5%)	259 (14.1%)	
US\$75 000+	1401 (27.8%)	1009 (31.4%)	392 (21.3%)	
Refused	620 (12.3%)	364 (11.3%)	256 (13.9%)	
Current smoking	695 (13.8%)	514 (16.0%)	181 (9.9%)	<0.001
<i>Medications</i>				
Depression/anxiety medications	472* (18.9%)	283 (17.1%)	189 (22.5%)	0.001

Continued

Table 1 Continued

	Overall (n=5047)	Men (n=3210)	Women (n=1837)	P value
Exogenous estrogen and/or progesterone	156 (8.5%)	0	156 (8.5%)	n/a
Exogenous androgens	84 (2.6%)	84 (2.6%)	0	n/a
<i>Anthropometrics</i>				
Weight (kg)	100.0±22.3	104.0±21.7	92.9±21.6	<0.001
BMI (kg/m ²)	34.3±6.8	33.7±6.3	35.4±7.5	<0.001
BMI categories				<0.001
Underweight/normal weight (<25 kg/m ²)	233 (4.6%)	148 (4.6%)	85 (4.6%)	
Overweight (25 to <30 kg/m ²)	1213 (24.1%)	837 (26.1%)	376 (20.5%)	
Class I obesity (30 kg/m ² to <35 kg/m ²)	1603 (31.8%)	1086 (33.9%)	517 (28.2%)	
Class II obesity (35 kg/m ² to <40 kg/m ²)	1093 (21.7%)	668 (20.9%)	425 (23.1%)	
Class III obesity (≥40 kg/m ²)	896 (17.8%)	463 (14.5%)	433 (23.6%)	
Waist circumference (cm)	112.3±15.8	114.1±15.6	109.3±15.6	<0.001
Waist circumference >sex and race-specific threshold†	4244 (84%)	2538 (79%)	1706 (92.9%)	<0.001
Systolic BP (mm Hg)	128.3±14.7	129.0±14.6	127.3±14.9	<0.001
Diastolic BP (mm Hg)	77.3±9.9	77.8±9.8	76.5±9.9	<0.001
<i>Laboratory tests</i>				
HbA1c (%) (mmol/mol)	7.5±0.5 58.3 (5.3)	7.5±0.5 58.4 (5.2)	7.5±0.5 58.3 (5.3)	0.678
HbA1c ≤7% (≤53 mmol/mol)	1101 (21.8%)	696 (21.7%)	405 (22.0%)	0.790
Total cholesterol (mg/dL) (mmol/L)	163.8±37.8 4.24±0.98	158.7±37.2 4.10±0.96	172.7±37.1 4.47±0.96	<0.001
Triglycerides (mg/dL) (mmol/L)	154.0±121.6 1.74±1.37	160.9±134.1 1.82±1.51	142.0±94.7 1.60±1.07	<0.001
HDL-cholesterol (mg/dL) (mmol/L)	43.4±10.6 1.12±0.27	41.0±9.3 1.06±0.24	47.7±11.3 1.23±0.29	<0.001
HDL-cholesterol ≤sex-specific threshold‡	2686 (53%)	1510 (47%)	1175 (64%)	<0.001
LDL-cholesterol (mg/dL) (mmol/L)	90.5±31.7 2.34±0.82	86.8±30.9 2.25±0.80	96.9±32.1 2.51±0.83	<0.001
LDL-cholesterol<100 mg/dL (<2.59 mmol/L)	3152 (65.0%)	2118 (69.2%)	1034 (57.7%)	<0.001
Fasting glucose (mg/dL) (mmol/L)	151.5±30.9 8.42±1.72	153.3±30.8 8.52±1.71	148.3±30.9 8.24±1.72	<0.001

All variables presented as means (SD) except as noted otherwise. Nominal p values are presented from exploratory comparisons of sex differences, and so should not be interpreted as testing hypotheses.
 *This question was added after study start and was answered by 2494 participants at baseline.
 †Waist circumference <88 cm for women, <102 cm for men (for Asian Americans, <80 cm for women, <90 cm for men).
 ‡HDL ≥50 mg/dL for women, ≥40 mg/dL for men.
 BMI, body mass index; BP, blood pressure; GED, Graduate Equivalency Degree; HbA1c, glycated hemoglobin; HDL, high-density lipoprotein; LDL, low-density lipoprotein.

inhibitors) or angiotensin receptor blockers (ARBs) (women 57.4% vs men 62.7%, p=0.003; [table 2](#)).

Age analysis of cardiometabolic risk factors and evidence-based treatment

Younger women had higher mean BMI ([figure 2](#)) and there was a trend for greater sex differences in mean BMI in the younger age group. Likewise, younger women had the highest prevalence of low HDL-C levels and the sex difference in low HDL-C was most evident in the younger population ([figure 2](#); online supplemental table 3).

Younger women treated with statins had modestly higher LDL-C levels and lower levels of achieving LDL-C<100 mg/dL ([figure 1](#)).

Socioeconomic factors

Compared with men, women in this cohort were more likely to be divorced, separated or widowed. Overall, women had fewer years of formal education and had lower incomes; the majority of women reported incomes below the median for the US population at that time ([table 1](#)). Women were more likely to take antidepressant and anxiolytic medications than men ([table 1](#)).

Sensitivity analysis by VA site

Sensitivity analyses did not indicate that the observed sex differences were due to differences in the cohort enrolled at VA study sites compared with non-VA study sites.

Table 2 Sex differences in treatment targets and evidence-based therapies at baseline

	Overall (n=5047)	Men (n=3210)	Women (n=1837)	P value
Lipid management				
Statin treatment (%) among those with elevated blood lipids	3210 (66.4%)	2181 (70.7%)	1029 (58.8%)	<0.001
LDL<100mg/dL (%) in entire cohort	3152 (65%)	2118 (69.2%)	1034 (57.7%)	<0.001
Hypertension management				
BP medications (%) among those with hypertension	3495 (72.4%)	2263 (73.1%)	1232 (71.2%)	0.454
BP<140/90 (%)	3800 (75.3%)	2401 (74.8%)	1399 (76.2%)	0.58
BP<130/80 (%)	2170 (43%)	1347 (42%)	823 (44.8%)	0.211
ACE inhibitor or ARB for those with hypertension (%)	2933 (60.8%)	1939 (62.7%)	994 (57.4%)	0.003

A Holm adjustment was applied to all p values presented in this table to control the type I error rate in this set of analyses. ACE, angiotensin converting enzyme; ARB, angiotensin receptor blocker; BP, blood pressure.


DISCUSSION

Compared with men with similar glycemic control and duration of diabetes, women with T2DM in the GRADE study were younger, and yet had a greater burden of traditional cardiometabolic risk factors (greater prevalence and severity of obesity, higher LDL-C and lower HDL-C), were less likely to receive evidenced-based treatment for dyslipidemia, and had a higher prevalence of adverse socioeconomic factors, all of which may ultimately contribute to greater burden of CV disease in women. Overall, the magnitude and severity of these risk factors appears to be more pronounced among younger women with T2DM. Unlike prior studies, women and men were equally likely to achieve target BP goals, although women with hypertension in the GRADE cohort were less likely to receive ACE inhibitors or ARB medications.

GRADE enrolled individuals under protocol-specified conditions of glycemic control, diabetes duration, and metformin monotherapy. Consequently, this cohort allows for a detailed assessment of sex differences in

cardiometabolic risk factors across a wide age range without the confounding factors of variable levels of glycemic control and diabetes duration, and the metabolic impact of different pharmacologic agents, as seen in earlier studies. The sex-stratified and age-stratified data highlight the disproportionate cardiometabolic risk factor burden among women with T2DM, especially younger women. Despite a decade or more difference in age, younger women, compared with older women and men, had a higher BMI, higher prevalence of increased waist circumference and low HDL-C, and were less likely to achieve target LDL-C levels despite statin treatment. Although CVD is generally thought to be a disease of older individuals, the disproportionate cardiometabolic burden among younger women with T2DM may contribute to a higher relative risk for premature macrovascular disease. This was demonstrated in a recent analysis of the Women's Health Study which showed that diabetes is a powerful predictor of premature coronary heart disease (CHD) in women younger than 55 years

LDL <100 mg/dL, on statins, by sex & age (prevalence)



Mean LDL (mg/dL), on statins, by sex and age




Figure 1 LDL-C<100mg/dL and mean LDL-C among statin users by sex and age at baseline in the Glycemia Reduction Approaches in Diabetes: A Comparative Effectiveness Study (GRADE) cohort. Red bars=women; blue bars=men. Error bars reflect the SE of means and proportions. The nominal p value for the interaction term between sex and age is provided to explore whether sex differences vary by age group, and so should not be interpreted as testing a hypothesis.




Figure 2 Baseline cardiometabolic risk factors by sex and age in the Glycemia Reduction Approaches in Diabetes: A Comparative Effectiveness Study (GRADE) cohort. Red bars=women; blue bars=men. Error bars reflect the SE of means and proportions. The nominal p value for the interaction term between sex and age is provided to explore whether sex differences vary by age group, and so should not be interpreted as testing hypotheses. BMI, body mass index.

(>10-fold higher adjusted relative risk),³⁰ and postacute MI studies that have shown higher mortality rates among younger women <50 years relative to men of comparable age.^{31 32} A worldwide meta-analysis demonstrated that the relative risk for CV mortality is highest among younger women with diabetes after adjustment for traditional cardiometabolic risk factors.¹

These results also demonstrate contemporary persistence of earlier reports of a high burden of cardiometabolic risk factors among women with T2DM and a lower likelihood of receiving evidence-based treatment for dyslipidemia.^{11 13 33} Even among women in GRADE treated with statins, LDL-C levels remained higher than men of comparable age. Younger women, black and Hispanic women, and women from lower SES groups (lower income and lower education) had higher LDL-C levels while on statin therapy. The reasons women are less likely to achieve target lipid control are complex and multifactorial, including being less likely than men to be offered statins by healthcare providers and less likely to believe that statins are safe and effective.¹³ Preferences for pharmacologic treatment and adherence to interventions differ by sex.³⁴ Women are more likely to decline statin treatment, be treated with less potent statins, have non-adherence due to cost, and to report discontinuing their statin because of a side effect.¹³ Although the US Food and Drug Administration (FDA) recently recommended removal of labeling of statins as contraindicated in pregnancy, they still recommend against taking statins

during pregnancy and breast feeding which impacts prescribing for women of childbearing potential.³⁵

Our data expand on findings from earlier studies of sex differences in cardiometabolic risk factors in several areas. Prior studies have shown that women with T2DM were less likely to receive evidence-based hypertension treatment compared with men.^{12 36} In contrast, women with hypertension in GRADE were as likely as men to receive antihypertensive medications and to achieve target BP goals. However, women with hypertension were less likely to receive ACE inhibitor or ARBs. While the reason for this finding in GRADE is unknown, earlier studies reported that women are more likely to experience side effects such as cough with ACE inhibitors and more likely to discontinue treatment.³⁷ The lower use of ACE inhibitors or ARBs in women has also been attributed to teratogenic risks of these agents in women of reproductive age.

In addition to traditional cardiometabolic risk factors, we identified sex differences in several adverse socioeconomic factors in GRADE. Women in GRADE were younger, had lower household income, lower levels of education, and were more likely to be single or divorced, separated or widowed than men. Several SES measures have been shown to have a strong inverse relationship with CHD including income, level of education, employment, and neighborhood socioeconomic factors,^{38 39} and women are disproportionately represented among those living in poverty. In a recent study examining the

relationship of low SES and premature CHD, individuals aged 35–64 years with lower SES experienced double the incidence of MI and CHD-related deaths compared with individuals of higher SES, with less than half of the excess events attributable to traditional risk factors (eg, cigarette smoking, hypertension, dyslipidemia, high BMI).⁴⁰ Based on computer simulation, 60% of the excess risk for premature MI and CHD mortality was tied to SES, and women comprised the majority of individuals in the low SES cohort.

Depression is another non-traditional risk factor associated with an increased risk for CHD and stroke.^{41–44} Women in GRADE were more likely to take antidepressants and anxiolytics than men across demographic and SES groups. A recent pooled analysis of prospective cohort studies showed that even mild depression (below the threshold for diagnosis of clinical depression) is associated with an increased incidence of CHD and stroke.⁴⁵

Studies that examined both traditional and non-traditional risk factors in populations with CHD found similar sex differences in the pattern of cardiometabolic and psychosocial risk factors. In the Variation in Recovery: Role of Gender on Outcomes of Young AMI (acute myocardial infarction) Patients Study, women younger than 55 years had significantly higher rates of cardiometabolic risk factors including diabetes (women 38.8% vs men 26.7%) and obesity (women 51% vs men 44.5%) than men. These younger women also had higher rates of depression and stress, poorer physical and mental health status, lower income, higher unemployment, and a lower quality of life. The authors concluded that young women with acute MI have a unique risk profile compared with men of comparable age, and that cardiometabolic factors and SES factors play a prominent role.^{6, 46} Taken together, these data suggest that women with T2DM, especially younger women, have a unique pattern of cardiometabolic and socioeconomic risk that may worsen lifetime CVD outcomes, and that adjustment for traditional cardiometabolic risk factors does not fully account for their higher risk.

The strengths of our study include detailed cardiometabolic data on a large sample size of individuals across a wide age range at a comparable early stage of diabetes duration, glycemic control, and glucose-lowering therapy. We accounted for sex-specific criteria for increased waist circumference and low HDL-C, which revealed relevant sex differences in these CV risk factors, unlike earlier studies which reported population means without adjustment. Limitations include the lack of data on the reproductive status of GRADE participants and reproductive risk factors for CVD including polycystic ovary syndrome or gestational diabetes. In addition, individuals who choose to volunteer in clinical trials may not be fully representative of the general population of those with T2DM, including the distribution and clinical characteristics of participants based on sex.

A large proportion of the GRADE cohort (n=1216, 24.1%) was enrolled at VA Medical Centers. Since the

patient population at VA-affiliated study sites was more likely to be male and older age, sensitivity analyses were conducted stratifying results by whether participants were enrolled in a VA study site or not and by age. Sensitivity analyses, along with age stratification, showed that results were similar among those enrolled at VA study sites compared with those at non-VA study sites.

There are different approaches for the statistical analysis of descriptive data. The statistical approach used in this descriptive analysis included the Holm procedure to control the type I error rate for hypothesis testing of the main analyses (sex differences in BP and lipid management) and nominal p values for the exploratory analyses. For outcome-based analyses, measures of discriminatory accuracy (such as multilevel analysis of individual heterogeneity and discriminatory accuracy)^{47, 48} might be appropriate; however, there were no outcomes in this analysis, being descriptive and cross-sectional in nature.

CONCLUSIONS

In conclusion, despite decades of research documenting that women with T2DM were less likely to receive evidence-based care for cardiometabolic risk factor management than men, data from the contemporary GRADE cohort demonstrates that women, particularly younger women, with T2DM continue to have a greater burden of adverse cardiometabolic and socioeconomic risk factors than men. Disparities persist in the evidence-based management of traditional cardiac risk factor of dyslipidemia, and substantial differences in non-traditional adverse socioeconomic factors remain evident, which may be more relevant for addressing cardiac risk in this population. Collectively, these data indicate the need to optimize care strategies to reduce the heightened risk factor burden and treatment gaps for women with T2DM.

Author affiliations

¹Department of Medicine, Section of Endocrinology, Yale School of Medicine, New Haven, Connecticut, USA

²Department of Medicine, Section of Endocrinology, VA Connecticut Healthcare System, West Haven, Connecticut, USA

³The Biostatistics Center, Department of Biostatistics and Bioinformatics, Milken Institute School of Public Health, The George Washington University, Rockville, Maryland, USA

⁴Department of Chronic Disease Epidemiology, Yale University School of Public Health, New Haven, Connecticut, USA

⁵Division of Diabetes, Endocrinology and Metabolism, University of Minnesota Health, Minneapolis, Minnesota, USA

⁶Division of Endocrinology, Metabolism, and Nutrition, Duke University Medical Center, Durham, North Carolina, USA

⁷Division of Diabetes, Endocrinology & Metabolism, University of Nebraska Medical Center, Omaha, Nebraska, USA

⁸Division of Endocrinology Diabetes and Clinical Nutrition, Oregon Health & Science University, Portland, Oregon, USA

⁹Diabetes Unit, Massachusetts General Hospital, Boston, Massachusetts, USA

¹⁰Division of Endocrinology, Diabetes and Hypertension, Brigham and Women's Hospital, Boston, Massachusetts, USA

¹¹MedStar Health Research Institute, Hyattsville, Maryland, USA

Collaborators The GRADE Research Group (April 30, 2021): Current Clinical Centers: Crandall, JP; McKee, MD; Behringer-Massera, S; Brown-Friday, J; Khori, E;

Ballentine-Cargill, K; Duran, S; Estrella, H; Gonzalez de la torre, S; Lukin, J (Albert Einstein College of Medicine). Phillips, LS; Burgess, E; Olson, D; Rhee, M; Wilson, P; Raines, TS; Boers, J; Costello, J; Maher-Albertelli, M; Mungara, R; Savoye, L; White, CA; Gullett, C; Holloway, L; Morehead, F; Person, S; Sibymon, M; Tanukonda, S; Adams, C; Ross, A (Atlanta VA Medical Center). Balasubramanyam, A; Gaba, R; Gonzalez Hattery, E; Ideozu, A; Jimenez, J; Montes, G; Wright, C (Baylor College of Medicine). Hollander, P; Roe, E; Jackson, A; Smiley, A; Burt, P; Estrada, L; Chionh, K (Baylor Research Institute). Ismail-Beigi, F; Falck-Ytter, C; Sayeed Kassem, L; Sood, A; Tiktin, M; Kulow, T; Newman, C; Stancil, KA; Cramer, B; Iacoboni, J; Kononets, MV; Sanders, C; Tucker, L; Werner, A; Maxwell, A; McPhee, G; Patel, C; Colosimo, L; Krol, A (Case Western Reserve University/Cleveland VA/MetroHealth Medical Center). Goland, R; Pring, J; Alfano, L; Kringas, P; Hausheer, C; Tejada, J; Gumpel, K; Kirpich, A; Schneider, H (Columbia University Medical Center). Green, JB; AbouAssi, H; Chatterjee, R; Feinglos, MN; English Jones, J; Khan, SA; Kimpel, JB; Zimmer, RP; Furst, M; Satterwhite, BM; Thacker, CR; Evans Kreider, K (Duke University Medical Center). Mariash, CN; Mather, KJ; Ismail, HM; Lteif, A; Mullen, M; Hamilton, T; Patel, N; Riera, G; Jackson, M; Pirics, V; Aguilar, D; Howard, D; Hurt, S (Indiana University). Bergenstal, R; Carlson, A; Martens, T; Johnson, M; Hill, R; Hyatt, J; Jensen, C; Madden, M; Martin, D; Willis, H; Konerza, W; Yang, S; Kleeberger, K; Passi, R (International Diabetes Center). Fortmann, S; Herson, M; Mularski, K; Glauber, H; Prihoda, J; Ash, B; Carlson, C; Ramey, PA; Schield, E; Torgirson-Ojerio, B; Arnold, K; Kauffman, M; Panos, E; Sahnow, S; Bays, K; Berame, K; Cook, J; Ghioni, D; Gluth, J; Schell, K; Criscola, J; Friason, C; Jones, S; Nazarov, S (Kaiser Permanente Northwest). Barzilay, J; Rassouli, N; Puttnam, R; Ojoawo, B; Nelson, R; Curtis, M; Hollis, B; Sanders-Jones, C; Stokes, K; El-Haq, Z; Kolli, A; Tran, T (Kaiser Permanente of Georgia). Wexler, D; Larkin, ME; Meigs, J; Chambers, B; Dushkin, A; Rocchio, G; Yepes, M; Steiner, B; Dulin, H; Cayford, M; Chu, K; DeManbey, A; Hillard, M; Martin, K; Thangthaeng, N; Gurry, L; Kochis, R; Raymond, E; Ripley, V; Stevens, C (Massachusetts General Hospital). Park, J; Aroda, V; Ghazi, A; Magee, M; Rensing, A; Loveland, A; Hamm, M; Hurtado, M; Kuhn, A; Leger, J; Manandhar, L; Mwicigi, F; Sanchez, O; Young, T (MedStar Health Research Institute/MedStar Baltimore). Garg, R, Lagari-Libhaber, V; Florez, HJ; Valencia, WM; Marks, J; Casula, S; Orpesa-Gonzalez, L; Hue, L; Cuadot, A; Nieto-Martinez, R; Riccio Veliz, AK; Gutt, M; Kendal, YJ; Veciana, B (Miami VA Healthcare System/University of Miami). Ahmann, A; Aby-Daniel, D; Joarder, F; Morimoto, V; Sprague, C; Yamashita, D; Cady, N; Rivera-Eschright, N; Kirchhoff, P; Morales Gomez, B; Adducci, J; Goncharova, A (Oregon Health & Science University). Hox, SH; Petrovitch, H; Matwichyna, M; Jenkins, V; Broadwater, L; Ishii, RR; Bermudez, NO (Pacific Health Research and Education Institute/VA Pacific Islands). Hsia, DS; Greenway, FL; Waguespack, C; King, E; Fry, G; Dragg, A; Gildersleeve, B; Arceneaux, J; Haynes, N; Thomassie, A; Pavlonis, M; Bourgeois, B; Hazlett, C (Pennington Biomedical Research Center). Mudaliar, S; Henry, R; Boeder, S; Pettus, J; Diaz, E; Garcia-Acosta, D; Maggs, S; DeLue, C; Stallings, A; Castro, E; Hernandez, S (San Diego VA Medical Center). Krakoff, J; Curtis, JM; Killean, T; Khalid, M; Joshevama, E; Diaz, E; Martin, D; Tsingine, K; Karshner, T (Southwestern American Indian Center). Albu, J; Pi-Sunyer, FX; Frances, S; Maggio, C; Ellis, E; Bastawrose, J; Gong, X (St. Luke's-Roosevelt Hospital). Banerji, MA; August, P; Lee, M; Lorber, D; Brown, NM; Josephson, DH; Thomas, LL; Tsovian, M; Cherian, A; Jacobson, MH; Mishko, MM (SUNY Downstate Medical Center/New York Hospital-Queens). Kirkman, MS; Buse, JB; Dostou, J; Machineni, S; Young, L; Bergamo, K; Goley, A; Kerr, J; Largay, JF; Guarda, S; Cuffee, J; Culmer, D; Fraser, R; Almeida, H; Coffey, S; Debnam, E; Kiker, L; Morton, S; Josey, K; Fuller, G (The University of North Carolina Diabetes Care Center). Garvey, WT; Cherrington, AL; Dyer, D; Lawson, MCR; Griffith, O; Agne, A; McCullars, S (University of Alabama Birmingham). Cohen, RM; Craig, J; Rogge, MC; Burton, K; Kersey, K; Wilson, C; Lipp, S; Vonder Meulen, MB; Adkins, C; Onadeko, T (University of Cincinnati/Cincinnati VA Medical Center). Rasouli, N; Baker, C; Schroeder, E; Razzaghi, M; Lyon, C; Penalzoza, R; Underkofler, C; Lorch, R; Douglass, S; Steiner, S (University of Colorado-Denver/VA). Sivitz, WI; Cline, E; Knosp, LK; McConnell, J; Lowe, T (University of Iowa). Herman, WH; Pop-Busui, R; Tan, MH; Martin, C; Waltje, A; Katona, A; Goodhall, L; Eggleston, R; Kuo, S; Bojescu, S; Bule, S; Kessler, N; LaSalle, E; Whitley, K (University of Michigan). Seaquist, ER; Bantle, A; Harindhanavudhi, T; Kumar, A; Redmon, B; Bantle, J; Coe, M; Mech, M; Taddese, A; Lesne, L; Smith, S (University of Minnesota). Desouza, C; Kuechenmeister, L; Shivaswamy, V; Burbach, S; Rodriguez, MG; Seipel, K; Alfred, A; Morales, AL; Eggert, J; Lord, G; Taylor, W; Tillson, R (University of Nebraska Medical Center/Omaha VA). Schade, DS; Adolphe, A; Burge, M; Duran-Valdez, E; Martinez, J; Bancroft, A; Kunkel, S; Ali Jamaledin Ahmad, F; Hernandez McGinnis, D; Pucchetti, B; Scripsick, E; Zamorano, A (University of New Mexico). DeFronzo, G; Cersosimo, E; Abdul-Ghani, M; Triplitt, C; Juarez, D; Garza, RI; Verastiqui, H; Wright, K; Puckett, C (UT Health San Antonio). Raskin, P; Rhee, C; Abraham, S; Jordan, LF; Sao, S; Morton, L; Smith, O; Osornio Walker, L; Schnurr-Breen, L; Ayala, R; Kreymer, RB; Sturgess, D (University of Texas-Southwestern Medical Center). Utzschneider, KM; Kahn, SE; Alarcon-Casas

Wright, L; Boyko, EJ; Tsai, EC; Trencle, DL; Trikudanathan, S; Fattaleh, BN; Montgomery, BK; Atkinson, KM; Kozedub, O; Concepcion, T; Moak, C; Prikhodko, N; Rhothisen, S (VA Puget Sound Health Care System/University of Washington). Elasy, TA; Martin, S; Shackelford, L; Goidel, R; Hinkle, N; Lovell, C; Myers, J; Lipps Hogan, J (Vanderbilt University). McGill, JB; Salam, M; Schweiger, T; Kissel, S; Recklein, C; Clifton, MJ (Washington University). Tamborlane, W; Camp, A; Gulanski, B; Inzucchi, SE; Pham, K; Alguar, M; Gatcomb, P; Lessard, K; Perez, M; Iannone, L; Magenheimer, E; Montosa, A (Yale University/Fair Haven Community Health Center/West Haven VA Medical Center). Study Units: Cefalu, WT; Fradkin, J; Burch, HB; Bremer, AA (NIH/NIDDK). Nathan, DM (Chairman's Office, Massachusetts General Hospital, Harvard Medical School). Lachin, JM; Krause-Steinrauf, H; Younes, N; Zebu, I; Butera, N; Buys, CJ; Fagan, A; Gao, Y; Ghosh, A; Gramzinski, MR; Hall, SD; Kazemi, E; Legowski, E; Liu, H; Suratt, CE; Tripputi, M; Arey, A; Backman, M; Bethupu, J; Lund, C; Mangat Dhaliwal, P; McGee, P; Mesimer, E; Ngo, L (Coordinating Center, The George Washington University Biostatistics Center). Steffes, M; Seegmiller, J; Saenger, A; Arends, V; Gabrielson, D (Central Biochemical Laboratory, University of Minnesota Advanced Research and Diagnostic Laboratory). Conner, T; Warren, S; Day, J; Huminik, J; Scrymgeour, A (Drug Distribution Center, VA Cooperative Studies Program Clinical Research Pharmacy Coordinating Center). Soliman, EZ; Pokharel, Y; Zhang, ZM; Campbell, C; Hu, J; Keasler, L; Hensley, S; Li, Y (ECG Reading Center, EPICARE, Wake Forest University). Mihalcea, R; Min, DJ; Perez-Rosas, V; Prosser, L; Resnicow, K; Ye, W (Economic Evaluation and Assessment Center, University of Michigan). Shao, H; Zhang, P (Economic Evaluation and Assessment Center, Centers for Disease Control and Prevention). Luchsinger, J; Sanchez, D; Assuras, S (Neurocognitive Coordinating Center, Columbia University Medical Center). Groessl, E; Sakha, F; Chong, H; Hillery, N (QWB Reading Center, University of California San Diego Health Services Research Center). Collaborators: Everett, BM (Cardiovascular Adjudication Advisor). Abdouch, I; Bahtiyar, G; Brantley, P; Broyles, FE; Canaris, G; Copeland, P; Craine, JJ; Fein, WL; Gliwa, A; Hope, L; Meiners, R; Meiners, V; O'Neal, H; Park, JE; Sacerdote, A; Sledge Jr., E; Soni, L; Steppel-Reznik, J; Turchin, A (Collaborating Investigators-Recruitment Sites). Brooks-Worrell, B; Hampe, CS; Palmer, JP; Shojaie, A (Beta Cell Ancillary Study). Higgins, J (Continuous Glucose Monitoring Sub-study). Fischer, L; Golden, S; Gonzalez, J; Naik, A; Walker, E (Emotional Distress Sub-study). Doner Lotenberg, L; Gallivan, JM; Lim, J; Tuncer, DM (National Diabetes Education Program Sub-study). Behringer-Massera, S (Recruitment Ancillary Study).

Contributors All authors affirm that authorship is merited based on the ICMJE authorship criteria. BIG, HK-S, TH, CD, DJW, and VRA contributed to the conception and design of research for the manuscript. BIG, HK-S, JBG, HA, CD, DJW, and VRA contributed to the acquisition of data for the manuscript. NMB and HK-S contributed to the statistical analysis for the manuscript. BIG, NMB, HK-S, JHL, TH, JBG, CES, HA, AJA, DJW and VRA contributed to the interpretation of data and results for the manuscript. HK-S, AJA, DJW, and VRA contributed to the supervision and management of research. BIG, NMB, CD, AJA, and VRA contributed to the drafting of the manuscript. BIG, NMB, HK-S, JHL, TH, JBG, CES, HA, CD, AJA, DJW, and VRA contributed to the critical review of the manuscript. BIG and NMB are the guarantors of this work and as such, had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

Funding The GRADE Study is supported by a grant from the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK) of the National Institutes of Health under Award Number U01DK098246. The planning of GRADE was supported by a U34 planning grant from the NIDDK (U34-DK-088043). The American Diabetes Association supported the initial planning meeting for the U34 proposal. The National Heart, Lung, and Blood Institute and the Centers for Disease Control and Prevention also provided funding support. The Department of Veterans Affairs provided resources and facilities. Additional support was provided by grant numbers P30 DK017047, P30 DK020541-44, P30 DK020572, P30 DK072476, P30 DK079626, P30 DK092926, U54 GM104940, UL1 TR000439, UL1 TR000445, UL1 TR001108, UL1 TR001409, UL1 TR001449, UL1 TR002243, UL1 TR002345, UL1 TR002378, UL1 TR002489, UL1 TR002529, UL1 TR002535, UL1 TR002537, UL1 TR001425 and UL1 TR002548. Educational materials have been provided by the National Diabetes Education Program. Material support in the form of donated medications and supplies has been provided by Becton, Dickinson and Company, Bristol-Myers Squibb, Merck, NovoNordisk, Roche Diagnostics, and Sanofi. The content of this manuscript is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Competing interests JBG reports grants from NIDDK during the conduct of the study; and grants and personal fees from Boehringer Ingelheim/Lilly, personal fees from NovoNordisk, grants from Roche, personal fees from Hawthorne Effect/Omada, grants and personal fees from Sanofi/Lexicon, personal fees from Pfizer,

grants from Glaxo SmithKline, personal fees from Bayer, grants from Merck, and grants and personal fees from AstraZeneca outside the submitted work. HA reports other from Novo Nordisk during the conduct of the study; and other from Novo Nordisk outside the submitted work. CD reports grants and personal fees from Novo Nordisk, personal fees from Astra Zeneca, grants from Sanofi, grants from the Department of Defense, and consultation for Bayer outside the submitted work. AJA reports personal fees from Novo Nordisk and personal fees from Lilly during the conduct of the study; and personal fees from Medtronic outside the submitted work. DJW reports other from Novo Nordisk outside the submitted work. VRA reports grants and other from Applied Therapeutics, grants and other from Fractyl, grants and other from Novo Nordisk, other from Pfizer, grants and other from Sanofi, other from Eli Lilly, and other from Janssen (Spouse) outside the submitted work. BIG, NMB, JHL, TH, HK-S and CES have nothing to disclose.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants. GRADE is a multicenter RCT, approved by over 30 institutional review boards (institutional review board approval was obtained at each participating institution). We have indicated the primary review board submission information from the most recent annual renewal below: The George Washington University Office of Human Research - Institutional Review Board (IRB Number: 071245; Last Approved: 8/15/2022; Expires: 8/23/2023.) Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement The Glycemia Reduction Approaches in Diabetes: A Comparative Effectiveness Study (GRADE) is being conducted with funding from the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK). This manuscript is based on the baseline (pretreatment) data from the 5047 participants enrolled into the study. This baseline data will be archived with the NIDDK data repository and will be available for sharing with other investigators in 2023.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Barbara I Gulanski <http://orcid.org/0000-0003-4056-3409>

Cyrus Desouza <http://orcid.org/0000-0001-6660-0568>

Vanita R Aroda <http://orcid.org/0000-0002-7706-4585>

REFERENCES

- Prospective Studies C, Asia Pacific Cohort Studies C. Sex-specific relevance of diabetes to occlusive vascular and other mortality: a collaborative meta-analysis of individual data from 980 793 adults from 68 prospective studies. *Lancet Diabetes Endocrinol* 2018;6:538–46.
- Huxley R, Barzi F, Woodward M. Excess risk of fatal coronary heart disease associated with diabetes in men and women: meta-analysis of 37 prospective cohort studies. *BMJ* 2006;332:73–8.
- Peters SAE, Huxley RR, Woodward M. Diabetes as a risk factor for stroke in women compared with men: a systematic review and meta-analysis of 64 cohorts, including 775,385 individuals and 12,539 strokes. *Lancet* 2014;383:1973–80.
- Benjamin EJ, Muntner P, Alonso A, et al. Heart disease and stroke statistics-2019 update: a report from the American heart association. *Circulation* 2019;139:e56–528.
- Regensteiner JG, Golden S, Huebschmann AG, et al. Sex differences in the cardiovascular consequences of diabetes mellitus: a scientific statement from the American heart association. *Circulation* 2015;132:2424–47.
- Bucholz EM, Strait KM, Dreyer RP, et al. Editor's choice—sex differences in young patients with acute myocardial infarction: a VIRGO study analysis. *Eur Heart J Acute Cardiovasc Care* 2017;6:610–22.
- Leifheit-Limson EC, D'Onofrio G, Daneshvar M, et al. Sex differences in cardiac risk factors, perceived risk, and health care provider discussion of risk and risk modification among young patients with acute myocardial infarction: the VIRGO study. *J Am Coll Cardiol* 2015;66:1949–57.
- Beckman AL, Bucholz EM, Zhang W, et al. Sex differences in financial barriers and the relationship to recovery after acute myocardial infarction. *J Am Heart Assoc* 2016;5:e003923.
- Smolderen KG, Strait KM, Dreyer RP, et al. Depressive symptoms in younger women and men with acute myocardial infarction: insights from the VIRGO study. *J Am Heart Assoc* 2015;4:e001424.
- Shah AJ, Ghasemzadeh N, Zaragoza-Macias E, et al. Sex and age differences in the association of depression with obstructive coronary artery disease and adverse cardiovascular events. *J Am Heart Assoc* 2014;3:e000741.
- Gu A, Kamat S, Argulian E. Trends and disparities in statin use and low-density lipoprotein cholesterol levels among US patients with diabetes, 1999–2014. *Diabetes Res Clin Pract* 2018;139:1–10.
- Wexler DJ, Grant RW, Meigs JB, et al. Sex disparities in treatment of cardiac risk factors in patients with type 2 diabetes. *Diabetes Care* 2005;28:514–20.
- Nanna MG, Wang TY, Xiang Q, et al. Sex differences in the use of statins in community practice. *Circ Cardiovasc Qual Outcomes* 2019;12:e005562.
- Vimalananda VG, Miller DR, Palnati M, et al. Gender disparities in lipid-lowering therapy among veterans with diabetes. *Womens Health Issues* 2011;21(4 Suppl):S176–81.
- Canto JG, Rogers WJ, Goldberg RJ, et al. Association of age and sex with myocardial infarction symptom presentation and in-hospital mortality. *JAMA* 2012;307:813–22.
- Lichtman JH, Leifheit EC, Safdar B, et al. Sex differences in the presentation and perception of symptoms among young patients with myocardial infarction: evidence from the VIRGO study (variation in recovery: role of gender on outcomes of young AMI patients). *Circulation* 2018;137:781–90.
- Gupta A, Barrabes JA, Strait K, et al. Sex differences in timeliness of reperfusion in young patients with ST-segment-elevation myocardial infarction by initial electrocardiographic characteristics. *J Am Heart Assoc* 2018;7:e007021.
- Leifheit-Limson EC, Spertus JA, Reid KJ, et al. Prevalence of traditional cardiac risk factors and secondary prevention among patients hospitalized for acute myocardial infarction (AMI): variation by age, sex, and race. *J Womens Health (Larchmt)* 2013;22:659–66.
- Eindhoven DC, Hilt AD, Zwaan TC, et al. Age and gender differences in medical adherence after myocardial infarction: women do not receive optimal treatment—the Netherlands claims database. *Eur J Prev Cardiol* 2018;25:181–9.
- Peters SAE, Colantonio LD, Zhao H, et al. Sex differences in high-intensity statin use following myocardial infarction in the united states. *J Am Coll Cardiol* 2018;71:1729–37.
- Ferrara A, Mangione CM, Kim C, et al. Sex disparities in control and treatment of modifiable cardiovascular disease risk factors among patients with diabetes: translating research into action for diabetes (triad) study. *Diabetes Care* 2008;31:69–74.
- Virani SS, Woodard LD, Ramsey DJ, et al. Gender disparities in evidence-based statin therapy in patients with cardiovascular disease. *Am J Cardiol* 2015;115:21–6.
- Lichtman JH, Wang Y, Jones SB, et al. Age and sex differences in in-hospital complication rates and mortality after percutaneous coronary intervention procedures: evidence from the NCDR (®). *Am Heart J* 2014;167:376–83.
- Dreyer RP, Dharmarajan K, Kennedy KF, et al. Sex differences in 1-year all-cause rehospitalization in patients after acute myocardial infarction: a prospective observational study. *Circulation* 2017;135:521–31.
- Gupta A, Wang Y, Spertus JA, et al. Trends in acute myocardial infarction in young patients and differences by sex and race, 2001 to 2010. *J Am Coll Cardiol* 2014;64:337–45.
- Nathan DM, Buse JB, Kahn SE, et al. Rationale and design of the glycemia reduction approaches in diabetes: a comparative effectiveness study (grade). *Diabetes Care* 2013;36:2254–61.
- Alberti KG, Eckel RH, Grundy SM, et al. Harmonizing the metabolic syndrome: a joint interim statement of the international diabetes federation task force on epidemiology and prevention; national heart, lung, and blood institute. *International Atherosclerosis Society; and International Association for the Study of Obesity Circulation* 2009;120:1640–5.
- Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma,

- without use of the preparative ultracentrifuge. *Clin Chem* 1972;18:499–502.
- 29 Florez HJ, Ghosh A, Pop-Busui R, *et al.* Differences in complications, cardiovascular risk factor, and diabetes management among participants enrolled at veterans affairs (VA) and non-VA medical centers in the glycemia reduction approaches in diabetes: a comparative effectiveness study (grade). *Diabetes Res Clin Pract* 2022;184:109188.
 - 30 Dugani SB, Moorthy MV, Li C, *et al.* Association of lipid, inflammatory, and metabolic biomarkers with age at onset for incident coronary heart disease in women. *JAMA Cardiol* 2021;6:437–47.
 - 31 Plakht Y, Elkis Hirsch Y, Shiyovich A, *et al.* Heterogeneity of diabetes as a risk factor for all-cause mortality after acute myocardial infarction: age and sex impact. *Diabetes Res Clin Pract* 2021;182:109117.
 - 32 Vaccarino V, Parsons L, Every NR, *et al.* Sex-based differences in early mortality after myocardial infarction. National Registry of myocardial infarction 2 participants. *N Engl J Med* 1999;341:217–25.
 - 33 Wright AK, Kontopantelis E, Emsley R, *et al.* Cardiovascular risk and risk factor management in type 2 diabetes mellitus. *Circulation* 2019;139:2742–53.
 - 34 Kautzky-Willer A, Harreiter J. Sex and gender differences in therapy of type 2 diabetes. *Diabetes Res Clin Pract* 2017;131:230–41.
 - 35 FDA. FDA requests removal of strongest warning against cholesterol-lowering statins during pregnancy. *Drug Safety Communication* July 20, 2021.
 - 36 Collier A, Ghosh S, Hair M, *et al.* Gender differences and patterns of cardiovascular risk factors in type 1 and type 2 diabetes: a population-based analysis from a Scottish region. *Diabet Med* 2015;32:42–6.
 - 37 Qiao Y, Shin J-I, Sang Y, *et al.* Discontinuation of angiotensin converting enzyme inhibitors and angiotensin receptor blockers in chronic kidney disease. *Mayo Clin Proc* 2019;94:2220–9.
 - 38 Schultz WM, Kelli HM, Lisko JC, *et al.* Socioeconomic status and cardiovascular outcomes: challenges and interventions. *Circulation* 2018;137:2166–78.
 - 39 Dray-Spira R, Gary-Webb TL, Brancati FL. Educational disparities in mortality among adults with diabetes in the U.S. *Diabetes Care* 2010;33:1200–5.
 - 40 Hamad R, Penko J, Kazi DS, *et al.* Association of low socioeconomic status with premature coronary heart disease in US adults. *JAMA Cardiol* 2020;5:899–908.
 - 41 Jha MK, Qamar A, Vaduganathan M, *et al.* Screening and management of depression in patients with cardiovascular disease: JACC state-of-the-art review. *J Am Coll Cardiol* 2019;73:1827–45.
 - 42 Daskalopoulou M, George J, Walters K, *et al.* Depression as a risk factor for the initial presentation of twelve cardiac, cerebrovascular, and peripheral arterial diseases: data linkage study of 1.9 million women and men. *PLoS One* 2016;11:e0153838.
 - 43 Carney RM, Freedland KE. Depression and coronary heart disease. *Nat Rev Cardiol* 2017;14:145–55.
 - 44 Lichtman JH, Froelicher ES, Blumenthal JA, *et al.* Depression as a risk factor for poor prognosis among patients with acute coronary syndrome: systematic review and recommendations: a scientific statement from the American heart association. *Circulation* 2014;129:1350–69.
 - 45 Harshfield EL, Pennells L, Schwartz JE, *et al.* Association between depressive symptoms and incident cardiovascular diseases. *JAMA* 2020;324:2396–405.
 - 46 Dreyer RP, Sciria C, Spatz ES, *et al.* Young women with acute myocardial infarction: current perspectives. *Circ Cardiovasc Qual Outcomes* 2017;10:e003480.
 - 47 Merlo J. Multilevel analysis of individual heterogeneity and discriminatory accuracy (MAIHDA) within an intersectional framework. *Soc Sci Med* 2018;203:74–80.
 - 48 Wemrell M, Bennet L, Merlo J. Understanding the complexity of socioeconomic disparities in type 2 diabetes risk: a study of 4.3 million people in Sweden. *BMJ Open Diabetes Res Care* 2019;7:e000749.

Table S.1. Sex differences in BMI and Waist Circumference, stratified by age, income, education, race, ethnicity.

	BMI (kg/m ²)			Increased Waist Circumference*		
	Men	Women	p-value	Men	Women	p-value
Age categories						
<45 years	35.2 ± 7.4	37.6 ± 8.5	0.051	258 (74.8%)	256 (93.1%)	0.133
45-54 years	34.6 ± 6.8	36 ± 7.7		677 (79.2%)	538 (93.1%)	
55-64 years	33.5 ± 5.9	35 ± 7.1		856 (79%)	647 (93.6%)	
≥65 years	32.4 ± 5.4	32.8 ± 5.9		741 (80.5%)	258 (90.2%)	
Income						
<\$10,000	33.1 ± 6.9	34.1 ± 7.4	0.193	102 (69.9%)	132 (88.6%)	0.486
\$10,000 - \$50,000	33.8 ± 6.5	36.2 ± 8		897 (77.3%)	723 (93.1%)	
\$50,000 - \$75,000	33.8 ± 5.8	35.3 ± 7.2		435 (82.5%)	239 (92.6%)	
\$75,000+	33.6 ± 6.2	35.3 ± 6.7		826 (82%)	373 (95.4%)	
Education						
Some high school	32.4 ± 5.8	33.7 ± 7.1	0.398	110 (63.2%)	162 (85.7%)	0.699
High school graduate	33.7 ± 6.1	35.1 ± 7.6		485 (76.7%)	370 (91.1%)	
Some college	34.3 ± 6.7	36.1 ± 7.4		776 (81.8%)	486 (95.1%)	
College graduate or more	33.3 ± 6.1	35.5 ± 7.5		1161 (80.1%)	680 (94.1%)	
Race						
White	33.9 ± 6.3	35.8 ± 7.3	0.671	1855 (81.2%)	953 (93.2%)	0.242
Black or African American	33.7 ± 6.2	35.3 ± 7.8		366 (74.8%)	466 (92.1%)	
Other or more than one race	32 ± 6.3	34.2 ± 6.9		284 (72.6%)	255 (93.1%)	
Ethnicity						
Hispanic/Latino	32.9 ± 6	33.7 ± 6.9	0.01	309 (61.7%)	374 (88.2%)	0.224
Non-Hispanic/Latino	33.8 ± 6.3	35.9 ± 7.5		2205 (82.3%)	1309 (94.3%)	

* Increased waist circumference ≥88 cm for females, ≥102 cm for males (for Asian Americans, ≥80 cm for females, ≥90 cm for males)

The nominal p-value for the interaction term between sex and the sociodemographic variable (i.e., age, income, education, race, or ethnicity) is provided to explore whether sex differences vary by the sociodemographic variable, and so should not be interpreted as testing hypotheses.

Table S.2. Sex differences in LDL levels, stratified by age, income, education, race, ethnicity.

	LDL (mg/dL), on statins			LDL <100 mg/dL, on statin		
	Men	Women	p-value	Men	Women	p-value
Age categories						
<45 years	84.8 ± 29.1	92.5 ± 32	0.941	106 (75.7%)	50 (65.8%)	0.134
45-54 years	83.5 ± 31	91.3 ± 31		347 (72%)	192 (65.8%)	
55-64 years	78.5 ± 26.8	87.2 ± 30.7		607 (79.9%)	316 (73.1%)	
≥65 years	74.6 ± 25.6	84.1 ± 27		624 (86.7%)	153 (73.6%)	
Income						
<\$10,000	84.9 ± 34.2	89.2 ± 34.7	0.388	47 (66.2%)	42 (61.8%)	0.719
\$10,000 - \$50,000	80.7 ± 29.2	88.4 ± 32.3		561 (77.5%)	286 (72%)	
\$50,000 - \$75,000	77.6 ± 28.9	89.8 ± 31.6		270 (79.6%)	110 (70.1%)	
> \$75,000	76.3 ± 24.5	83.6 ± 23.7		612 (84.2%)	188 (76.7%)	
Education						
Some high school	86.9 ± 29.3	94.2 ± 37.7	0.802	67 (71.3%)	55 (60.4%)	0.777
High school graduate	81.7 ± 28.8	91.3 ± 29.1		309 (76.9%)	148 (65.2%)	
Some college	78.5 ± 28.8	91.3 ± 29.1		478 (79%)	191 (72.9%)	
College graduate or more	76.9 ± 26.5	86.7 ± 29.3		830 (82.9%)	317 (74.1%)	
Race						
White	75.6 ± 25.8	84.5 ± 28.7	0.123	1299 (84%)	441 (75%)	0.023
Black or African American	91.3 ± 31.8	94.9 ± 32.6		189 (61.6%)	170 (61.4%)	
Other or more than one race	81.8 ± 29.6	91.3 ± 30.2		182 (79.8%)	89 (69%)	
Ethnicity						
Hispanic/Latino	85.6 ± 31.9	92.4 ± 36.2	0.367	177 (71.7%)	119 (63%)	0.573
Non-Hispanic/Latino	77.7 ± 27.1	87.3 ± 28.7		1498 (81.4%)	585 (72.1%)	

The nominal p-value for the interaction term between sex and the sociodemographic variable (i.e., age, income, education, race, or ethnicity) is provided to explore whether sex differences vary by the sociodemographic variable, and so should not be interpreted as testing hypotheses.

Table S.3. Sex differences in low HDL and Blood Pressure, stratified by age, income, education, race, ethnicity.

	Low HDL ^a			BP <140/90, on medication		
	Men	Women	p-value	Men	Women	p-value
Age categories						
<45 years	174 (50.3%)	231 (83.4%)	<0.001	106 (71.1%)	94 (74.6%)	0.835
45-54 years	429 (50.3%)	392 (67.6%)		395 (73.7%)	262 (72.6%)	
55-64 years	493 (45.5%)	402 (58.3%)		571 (70.1%)	374 (72.2%)	
≥65 years	409 (44.3%)	148 (51.6%)		547 (71.7%)	164 (72.6%)	
Income						
<\$10,000	66 (45.2%)	95 (63.3%)	0.349	56 (60.2%)	67 (73.6%)	0.366
\$10,000 - \$50,000	586 (50.5%)	508 (65.3%)		575 (70%)	366 (69.7%)	
\$50,000 - \$75,000	264 (50.1%)	169 (65.3%)		275 (72.8%)	136 (73.9%)	
>\$75,000	422 (41.9%)	249 (63.5%)		544 (75.1%)	209 (76.3%)	
Education						
Some high school	78 (44.8%)	106 (55.8%)	0.42	69 (72.6%)	82 (71.9%)	0.326
High school graduate	298 (47.3%)	273 (67.1%)		320 (70.6%)	199 (75.7%)	
Some college	483 (50.9%)	352 (69%)		471 (69.1%)	259 (72.1%)	
College graduate or more	646 (44.5%)	441 (60.7%)				
Race						
White	1173 (51.4%)	707 (69%)	0.008	1184 (72.5%)	500 (72.9%)	0.13
Black or African American	161 (32.8%)	258 (50.8%)		224 (62.9%)	271 (70.9%)	
Other or more than one race	156 (39.9%)	195 (71.2%)		191 (77.3%)	112 (75.7%)	
Ethnicity						
Hispanic/Latino	221 (44%)	298 (69.8%)	0.002	178 (67.7%)	163 (71.5%)	0.544
Non-Hispanic/Latino	1273 (47.5%)	868 (62.5%)		1429 (72%)	725 (73%)	

^a Low HDL <50 mg/dL for females, <40 mg/dL for males

The nominal p-value for the interaction term between sex and the sociodemographic variable (i.e., age, income, education, race, or ethnicity) is provided to explore whether sex differences vary by the sociodemographic variable, and so should not be interpreted as testing hypotheses.

Table S.4. Exploratory analysis results for prevalence of cardiometabolic risk factors by sex, stratified by age, education, race, ethnicity, and income.

CVD Risk Factor	P-value for interaction between sex and				
	Age	Education	Race	Ethnicity	Income
BMI (kg/m ²)	0.051	0.398	0.671	0.01	0.193
Waist circumference \geq gender-specific threshold ^a	0.133	0.699	0.242	0.224	0.486
Insulin sensitivity (HOMA2-S%)	0.212	0.677	0.03	0.65	0.647
Insulin secretion (Insulinogenic index)	0.756	0.789	0.041	0.81	0.179
HDL $<$ gender-specific threshold ^b	<0.001	0.42	0.008	0.002	0.349
Triglycerides (mg/dL)	0.323	0.828	0.239	0.777	0.129
Metabolic Syndrome	0.292	0.338	0.503	0.161	0.706
Depression/anxiety medications (%) ^c	0.922	0.113	0.57	0.085	0.14
ASCVD risk score (%) ^d	<0.001	0.038	<0.001	<0.02	0.16
Heart attack or stroke (%)	0.338	0.121	0.832	0.671	0.546

^a Normal waist circumference <88 cm for females, <102 cm for males (for Asian Americans, <80 cm for females, <90 cm for males).

^b Normal HDL ≥ 50 mg/dL for females, ≥ 40 mg/dL for males.

^c This question was added after the study started and was answered by 2494 participants at baseline.

^d Estimated 10-year ASCVD risk. ASCVD risk scores are intended for primary prevention of CVD only, and so this variable is analyzed only for those without history of heart attack or stroke.

The nominal p-value for the interaction term between sex and the sociodemographic variable (i.e., age, income, education, race, or ethnicity) is provided to explore whether sex differences vary by the sociodemographic variable, and so should not be interpreted as testing hypotheses.

Table S.5. Exploratory analysis results for evidence-based treatments for lipid and blood pressure management by sex, stratified by age, education, race, ethnicity, and income.

CVD Risk Factor	P-value for interaction between sex and				
	Age	Education	Race	Ethnicity	Income
LDL (mg/dL) in entire cohort	0.006	0.181	0.025	0.02	0.173
LDL (mg/dL) in those treated with statins	0.941	0.802	0.123	0.367	0.388
LDL < 100 mg/dL in those treated with statins	0.134	0.777	0.023	0.573	0.719
LDL (mg/dL) in those NOT treated with statins	0.003	0.373	0.631	0.359	0.374
LDL < 100 mg/dL in those NOT treated with statins	0.345	0.97	0.253	0.408	0.536
Systolic BP (mmHg) in entire cohort	0.001	0.003	0.152	0.015	0.255
Diastolic BP (mmHg) in entire cohort	0.326	0.025	0.017	<0.001	0.003
Systolic BP (mmHg) in those treated with blood pressure medications	0.063	0.176	0.054	0.656	0.499
Diastolic BP (mmHg) in those treated with blood pressure medications	0.344	0.498	<0.001	0.237	0.083
BP <140/90 mmHg in those treated with blood pressure medications	0.835	0.326	0.13	0.544	0.336
BP <130/80 mmHg in those treated with blood pressure medications	0.17	0.179	<0.001	0.854	0.575

The nominal p-value for the interaction term between sex and the sociodemographic variable (i.e., age, income, education, race, or ethnicity) is provided to explore whether sex differences vary by the sociodemographic variable, and so should not be interpreted as testing hypotheses.