


# Association of rurality and identifying as black with receipt of specialty care among patients hospitalized with a diabetic foot ulcer: a Medicare cohort study

Lindsay Taylor,<sup>1</sup> Ronald Gangnon,<sup>2</sup> W Ryan Powell,<sup>1,3</sup> Joseph Kramer,<sup>1,3</sup> Amy J H Kind,<sup>1,3</sup> Christie M Bartels,<sup>1</sup> Meghan B Brennan <sup>1</sup>

**To cite:** Taylor L, Gangnon R, Powell WR, *et al.* Association of rurality and identifying as black with receipt of specialty care among patients hospitalized with a diabetic foot ulcer: a Medicare cohort study. *BMJ Open Diab Res Care* 2023;**11**:e003185. doi:10.1136/bmjdr-2022-003185

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjdr-2022-003185>).

Received 20 October 2022  
Accepted 28 March 2023



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

<sup>1</sup>Medicine, University of Wisconsin-Madison, Madison, Wisconsin, USA  
<sup>2</sup>Population Health, University of Wisconsin-Madison, Madison, Wisconsin, USA  
<sup>3</sup>University of Wisconsin Center for Health Disparities Research, Madison, Wisconsin, USA

**Correspondence to**  
Dr Meghan B Brennan;  
mbbrennan@medicine.wisc.edu

## ABSTRACT

**Introduction** Rural patients with diabetic foot ulcers, especially those identifying as black, face increased risk of major amputation. Specialty care can reduce this risk. However, care disparities might beget outcome disparities. We aimed to determine whether a smaller proportion of rural patients, particularly those identifying as black, receive specialty care compared with the national proportion.

**Research design and methods** This 100% national retrospective cohort examined Medicare beneficiaries hospitalized with diabetic foot ulcers (2013–2014). We report observed differences in specialty care, including: endocrinology, infectious disease, orthopedic surgery, plastic surgery, podiatry, or vascular surgery. We used logistic regression to examine possible intersectionality between rurality and race, controlling for sociodemographics, comorbidities, and ulcer severity and including an interaction term between rurality and identifying as black.

**Results** Overall, 32.15% (n=124 487) of patients hospitalized with a diabetic foot ulcer received specialty care. Among rural patients (n=13 100), the proportion decreased to 29.57%. For patients identifying as black (n=21 649), the proportion was 33.08%. Among rural patients identifying as black (n=1239), 26.23% received specialty care. This was >5 absolute percentage points less than the overall cohort. The adjusted OR for receiving specialty care among rural versus urban patients identifying as black was 0.61 (95% CI 0.53 to 0.71), which was lower than that for rural versus urban patients identifying as white (aOR 0.85, 95% CI 0.80 to 0.89). This metric supported a role for intersectionality between rurality and identifying as black.

**Conclusions** A smaller proportion of rural patients, particularly those identifying as black, received specialty care when hospitalized with a diabetic foot ulcer compared with the overall cohort. This might contribute to known disparities in major amputations. Future studies are needed to determine causality.

## INTRODUCTION

Rural patients face an estimated 35% higher odds of major amputation following diabetic foot ulcers compared with patients living in urban areas.<sup>1 2</sup> Patients identifying as black

## WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Rural patients, particularly those who identify as black, face increased risk of major amputation following hospitalization with a diabetic foot ulcer.

## WHAT THIS STUDY ADDS

⇒ A smaller proportion of these same patients receive specialty care for their foot ulcer.

## HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ Differences in specialty care might be contributing to disparities in major amputations among patients with diabetic foot ulcers. Further research on the impact of specialty care on disparities in diabetic foot ulcer outcomes is needed.

are twice as likely to undergo a major amputation compared with those identifying as non-Hispanic white.<sup>3 4</sup> The intersection of these social identities amplifies the risk of poor outcomes: rural Medicare patients identifying as black face a greater than 10% absolute increased risk of major amputation or death.<sup>3</sup> Both social determinants of health and health system factors may contribute to these disparities, although we know little about disease-specific root causes or ultimate solutions.

To investigate the associations between social determinants of health and health system factors with disparities in major amputation, we adopted an intersectionality lens.<sup>5–8</sup> Intersectionality emerged from black feminism as a way to understand and address social injustices faced by multiply marginalized people.<sup>9</sup> Its goal is to improve social justice for multiply marginalized people, such as rural Americans identifying as black. Core tenets include: overlapping identities, historically oppressed populations, and

social determinants of health.<sup>10</sup> These overlaps have the capacity to amplify disparities, including healthcare disparities.

Our study's objective was to identify healthcare system factors that may be contributing to rural disparities in major amputations, particularly among rural patients identifying as black. The main factor we investigated was inpatient specialty care, as advocated by multiple professional societies as best practice for those with diabetic foot ulcers.<sup>11–13</sup> We hypothesize that rural patients hospitalized with diabetic foot ulcers receive less specialty care when compared with the national overall cohort of patients with similar comorbidities and ulcer severities. We further hypothesize that differences in specialty care particularly impact rural patients identifying as black, consistent with our use of an intersectionality lens. We restricted our analysis to patients identifying as black because our data set contained few rural patients identifying as Hispanic and concerns for misclassification of other racial and ethnic identities.<sup>14</sup> Our hypothesis is based on (1) The construct of intersectionality, (2) Previous Medicare findings that rural Medicare patients identifying as black face a 1.24-fold amplified risk of major amputation or death following hospitalization with a diabetic foot ulcer, and (3) Prior work demonstrating that rural patients lack access to specialists.<sup>3,9,15</sup> Our hope is that this initial study will identify potential differences in healthcare that may be contributing to disparities in major amputations. Further studies defining a more causal relationship will be needed, with the ultimate goal of designing interventions to provide equitable healthcare and alleviate disparities in major amputations.

## MATERIALS AND METHODS

### Data sources

We used a 100% national sample of adult Medicare beneficiaries hospitalized between January 1, 2013 and December 31, 2014, obtained through the Centers for Medicare and Medicaid Services. We required patients to be continuously enrolled in Medicare parts A and B for 12 months preceding index hospitalization and have a geolinkable address. We used five-digit ZIP codes to link Medicare data to categorize rurality (Rural-Urban Commuting Area, or RUCA, codes) and nine-digit ZIP codes to categorize neighborhood disadvantage (Neighborhood Atlas Area Deprivation Index, or ADI).<sup>16,17</sup>

### Study design

We constructed a retrospective, national cohort of Medicare beneficiaries hospitalized with diabetic foot ulcers by first identifying patients with diabetes based on the Chronic Conditions Warehouse (CCW) flag, which is based on diabetes-related diagnostic billing codes (eg, 1 code in inpatient, skilled nursing facility, home health claims or 2 codes in hospital outpatient or carrier claims within a 2-year window).<sup>18</sup> Next, we categorized diabetic foot ulcers as early stage (ie, ulcers not complicated by


osteomyelitis or gangrene), osteomyelitis, or gangrene. This was done using a validated algorithm generating severity categories that corresponded to hospital length of stay, amputation, and mortality.<sup>19</sup> We excluded patients with incomplete claims due to health maintenance organization or railroad benefits, as well as those admitted to psychiatric or long-term acute care hospitals, due to the potential for incomplete claims.<sup>20</sup> For patients hospitalized more than once, only the first index admission was included to maintain statistical independence. We used the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) cohort reporting guidelines to present our findings.<sup>21</sup>

### Outcome

Our outcome was receipt of specialty care during the index hospitalization. This was defined dichotomously as being seen by at least one of six specialists identified using National Provider and Plan Enumeration System Medical Specialty codes/Provider Taxonomy codes: endocrinology, infectious disease, orthopedic surgery, plastic surgery, podiatry, or vascular surgery (online supplemental table 1).<sup>22,23</sup> We chose these six specialties because they are the most frequent disciplines represented in published descriptions of multidisciplinary teams caring for patients with diabetic foot ulcers as identified through systematic review.<sup>11</sup> To ensure that the specialist provided care for the ulcer, we required them to code for it during billing. Endocrinologists were the exception. We required endocrinologists to code for diabetes during billing instead of the ulcer, since this most directly reflected the physiologic aspect addressed by their specialty.<sup>11</sup>

### Primary explanatory variables

We investigated two primary explanatory variables: rurality and identifying as black. Rurality was captured using RUCA codes: urban (RUCA 1, metropolitan area core, referent), suburban (RUCA 2–6, metropolitan area with commuting and micropolitan areas), and rural (RUCA 7–10, small town and rural areas) which is based on population density, urbanization, and daily commuting factors (figure 1).<sup>17,24</sup> We operationalized race using



**Figure 1** Map of the contiguous USA identifying ZIP codes by the rurality categorization used in our primary analysis.

Research Triangle Institute codes, where patients identify as: American Indian/Alaskan Native, Asian/Pacific Islander, black/African American (hereafter black), Hispanic, other, unknown, and non-Hispanic white (hereafter white).<sup>25</sup> We focused analysis on patients identifying as black because of low numbers of rural patients identifying as Hispanic and concerns for misclassification of other racial and ethnic identities.<sup>14</sup>

### Covariates

Sociodemographic variables included age, sex, receiving Medicaid coverage the year prior to hospitalization, and neighborhood disadvantage. Neighborhood disadvantage was measured using the ADI, a social determinant of health composite construct consisting of 17 census block group measures that describe housing, poverty, employment and education characteristics.<sup>16</sup> More disadvantaged neighborhoods have higher national ADI rankings. Ulcer severity was operationalized using the validated algorithm described above.<sup>14</sup> Comorbidities were adjusted for using Medicare CCW flags, Elixhauser Comorbidity Index variables, and baseline-year Hierarchical Condition Category (HCC) scores.<sup>26 27</sup> Higher HCC scores correlate with higher healthcare costs and utilization and is commonly used as a proxy for illness burden and patient complexity.<sup>27</sup> The following comorbidities were identified using validated CCW conditional categories: myocardial infarction, ischemic heart disease, hyperlipidemia, hypertension, and stroke.<sup>18</sup> Nine Elixhauser comorbidities with <5% prevalence in our cohort were consolidated into a single indicator (positive if at least one was present).<sup>26</sup>

### Statistical analysis

We described patient characteristics, overall and stratified by both rurality and racial and ethnic identity. Capitalizing on the strength of a full US Medicare population, we used observed differences in specialty care as well as predicted probabilities to explore the intersection of rurality and identifying as black on an additive scale (testing whether interactions were greater than the sum of each, individual effect).<sup>28</sup> We used ORs to describe the potential role of intersectionality on a multiplicative scale (testing whether interactions were greater than the product of each, individual effect).<sup>28</sup> Because we used a full, 100% hospitalized Medicare patient data set, observations are actual differences between subgroups, not estimates; therefore, no tests of statistical differences were run.

Our main analysis focused on the primary outcome of receiving care from at least one specialist. We performed logistic regression, sequentially building models to assess how covariates and interactions influence the associations between rurality and receipt of specialty care. We started with univariable analysis (Model 1), then added age and sex (Model 2). Model 3 included all sociodemographic factors. When examining race, we only report regression results for patients identifying as black or

white because (1) Our study's hypothesis was focused on this difference, (2) Our data set contained limited numbers of rural patients identifying as Hispanic (n=431) and (3) The Research Triangle Institute race variable may miscategorize patients identifying as other races.<sup>14</sup> Our fourth model added ulcer severity and comorbidities. Model 5 included an interaction term between rural residence and identifying as black if it met our *a priori* level of significance with a two-sided value of  $p < 0.001$ . For the interaction model (Model 5), ORs and adjusted predicted probabilities were calculated by racial identity and rurality.<sup>28</sup> When reporting predicted probabilities, we used a hypothetical patient with the common characteristics of the cohort. We used R software to perform our statistical analysis.<sup>29</sup>

We performed supportive, stratified analyses: observed rates of specialty care stratified by ulcer severity to account for confounding by indication (eg, patients with more advanced ulcers may be more likely to receive specialty care). Stratified analysis may begin to identify ulcer severity windows or targets for potential future interventions. We also performed secondary analysis, categorizing rurality differently using RUCA codes and Census data, to ensure our findings with respect to rurality were robust.<sup>30</sup> Specifically, in the secondary analysis rurality was categorized as urban (RUCA codes 1.0, 1.1, 2.0, 2.1, 3.0, 4.1, 5.1, 7.1, 8.1, 10.1), large rural (4.0, 4.2, 5.0, 5.2, 6.0, 6.1), and small rural/isolate (7.0, 7.2, 7.3, 7.4, 8.0, 8.2, 8.3, 8.4, 9.0, 9.1, 9.2, 10.0, 10.2, 10.3, 10.4, 10.5, 10.6; online supplemental table 2).

### RESULTS

During the 2-year study period, 124 487 Medicare beneficiaries were hospitalized with diabetic foot ulcers. Over half of the cohort was male with a mean age of 72 years, and 28.51% identified as a racial or ethnic minority. Of the total patients, 13 100 (10.52%) lived in a rural setting. Most rural patients identified as white, with a higher proportion living in disadvantaged neighborhoods and previously receiving Medicaid compared with the overall cohort. Prior diagnoses of comorbid conditions were slightly lower among rural patients compared with the overall population (table 1). Relative to the overall cohort, the group of patients identifying as black—regardless of where they lived—had higher proportions of those who: lived in disadvantaged neighborhoods, previously received Medicaid, experienced a higher prevalence of comorbidities, and received more severe ulcer diagnoses (table 1).

### Receipt of specialty care

Overall, 40 027 (32.2%) patients saw at least one specialist (table 1). A smaller proportion (29.57%) of rural patients received specialty care. A higher proportion of patients identifying as black (33.08%) received specialty care; however, among rural patients also identifying as black, the observed proportion receiving specialty care



**Table 1** Characteristics of patients hospitalized with diabetic foot ulcer by rurality and racial and ethnic identity

Characteristic	Rurality			Racial and ethnic identity			
	Full cohort* n=124 487 (100%)	Urban patients n=84 590 (67.95%)	Suburban patients n=26 262 (21.10%)	Rural patients n=13 100 (10.52%)	Non-Hispanic white n=88 525 (71.11%)	Black n=21 649 (17.39%)	Hispanic n=10 158 (8.16%)
Specialty care							
Endocrinology	377 (0.30)	290 (0.34)	61 (0.23)	26 (0.20)	287 (0.32)	52 (0.24)	32 (0.32)
Infectious disease	2020 (1.62)	1463 (1.73)	369 (1.41)	187 (1.43)	1477 (1.67)	337 (1.56)	143 (1.41)
Orthopedic surgery	6435 (5.17)	3882 (4.59)	1641 (6.25)	900 (6.87)	4848 (5.48)	858 (3.96)	496 (4.88)
Plastic surgery	1018 (0.82)	799 (0.94)	150 (0.57)	63 (0.48)	655 (0.74)	222 (1.03)	95 (0.94)
Podiatry	16 267 (13.07)	11 973 (14.15)	2980 (11.35)	1285 (9.81)	11 360 (12.83)	2609 (12.05)	1682 (16.56)
Vascular surgery	15 668 (12.59)	10 934 (12.93)	3139 (11.95)	1556 (11.88)	10 444 (11.80)	3405 (15.73)	1297 (12.77)
Received specialty care	40 027 (32.15)	28 021 (33.13)	8049 (30.65)	3874 (29.57)	27 868 (31.48)	7162 (33.08)	3577 (35.21)
Sociodemographics							
Age, mean (SD)	71.54 (13.02)	71.94 (13.15)	70.54 (12.71)	70.99 (12.64)	72.86 (12.43)	68.19 (14.14)	68.38 (13.50)
Male	71 286 (57.26)	48 019 (56.77)	15 287 (58.21)	7663 (58.50)	51 447 (58.12)	11 263 (52.03)	6062 (59.68)
Race and ethnicity							
Black	21 649 (17.39)	17 346 (20.51)	3003 (11.43)	1239 (9.56)	--	--	--
Hispanic	10 158 (8.16)	8075 (9.55)	1283 (4.89)	431 (3.29)	--	--	--
Non-Hispanic white	88 525 (71.11)	56 585 (66.89)	21 175 (80.63)	10 707 (81.73)	--	--	--
Other/unknown†	4155 (3.34)	2584 (3.05)	801 (3.05)	723 (5.52)	--	--	--
Reside in ≥ 80th percentile ADI	26 430 (21.23)	16 044 (18.97)	6198 (23.60)	4050 (30.92)	14 111 (15.94)	8252 (38.12)	3219 (31.69)
Mean ADI percentile (SD)	53.29 (27.56)	48.34 (28.65)	61.68 (22.07)	69.09 (18.37)	50.09 (26.51)	64.31 (27.48)	59.51 (28.50)
Medicaid receipt	46 199 (37.11)	30 731 (36.33)	10 033 (38.20)	5365 (40.95)	25 551 (28.86)	11 929 (55.10)	6570 (64.68)
Ulcer severity and comorbidities							
Ulcer severity							
Early stage	81 277 (65.29)	55 144 (65.19)	17 189 (65.45)	8603 (65.67)	60 227 (68.03)	12 735 (58.82)	5823 (57.32)
Osteomyelitis	26 892 (21.60)	18 188 (21.50)	5705 (21.72)	2890 (22.06)	18 967 (21.43)	4486 (20.72)	2432 (23.94)
Gangrene	16 318 (13.11)	11 258 (13.31)	3368 (12.82)	1607 (12.27)	9331 (10.54)	4428 (20.45)	1903 (18.73)
History of myocardial infarction	22 768 (18.29)	15 097 (17.85)	5052 (19.24)	2521 (19.24)	16 635 (18.79)	3510 (16.21)	1898 (18.68)
History of ischemic heart disease	104 863 (84.24)	72 016 (85.14)	21 672 (82.52)	10 742 (82.00)	74 645 (84.32)	18 209 (84.11)	8710 (85.75)
History of stroke or TIA	40 772 (32.75)	28 754 (33.99)	8061 (30.69)	3767 (28.76)	27 557 (31.13)	8571 (39.59)	3452 (33.98)
History of hyperlipidemia	116 144 (93.30)	79 315 (93.76)	24 331 (92.65)	12 028 (91.82)	82 855 (93.60)	19 895 (91.90)	9593 (94.44)
History of hypertension	123 368 (99.10)	83 867 (99.15)	26 012 (99.05)	12 957 (98.91)	87 612 (98.97)	21 575 (99.66)	10 079 (99.22)

Continued

**Table 1** Continued

	Rurality				Racial and ethnic identity			
History of obesity	29 370 (23.59)	19 826 (23.44)	6468 (24.23)	3030 (23.13)	21 578 (24.38)	4999 (23.09)	1998 (19.67)	
History of peripheral vascular disease	48 077 (38.62)	32 929 (38.93)	10 130 (38.57)	4864 (37.13)	32 813 (37.07)	9313 (43.02)	4436 (43.67)	
History of renal failure	51 841 (41.64)	36 231 (42.83)	10 401 (39.60)	4864 (37.13)	32 941 (37.21)	11 786 (54.44)	5182 (51.01)	
HCC Community Score, mean (SD)	2.68 (1.90)	2.68 (1.94)	2.68 (1.85)	2.68 (1.77)	2.59 (1.85)	2.98 (2.06)	2.74 (1.97)	

<sup>a</sup>We were unable to categorize the rurality of 535 patients.

ADI, Area Deprivation Index; HCC, hierarchical conditional category; TIA, transient ischemic attack.

\*We were unable to categorize the rurality of 535 patients.

ADI, Area Deprivation Index; HCC, hierarchical conditional category; TIA, transient ischemic attack.

decreased to 26.23%. This was 5.92 absolute percentage points less than the overall cohort, more than twice the decrease experienced by the general rural population, and a 6.85 absolute percentage point decrease from the overall population identifying as black.

After controlling for sociodemographic factors, comorbidity and ulcer severity, rural patients remained less likely to receive specialty care than their urban counterparts (aOR 0.83, 95% CI 0.79 to 0.86; [table 2](#), Model 4). In the same model, patients identifying as black were less likely to receive specialty care than those identifying as white (aOR 0.92, 95% CI 0.89 to 0.97). Applying an intersectionality perspective, we found a significant interaction between identifying as black and rural residence, indicating that the intersection of these two social identities diminished the likelihood of receiving specialty care ([table 2](#), Model 5). On the multiplicative scale, the OR for rural versus urban patients identifying as black was 0.61 (95% CI 0.53 to 0.71), which was lower than the OR for rural versus urban patients identifying as white (aOR 0.85, 95% CI 0.80 to 0.89). On the additive scale, the predicted probability of receiving specialty care when diagnosed with an early stage ulcer was 8.89 absolute percentage points less among rural patients identifying as black compared with urban patients identifying as black ( $p<0.05$ ). This rural-urban difference was more pronounced than the difference in predicted probabilities of receiving specialty care for rural compared with urban patients identifying as white presenting with early ulcer, which was only 3.25 absolute percentage points ( $p<0.05$ ; [figure 2](#)). In a secondary analysis when categorizing rurality four ways, these results were not substantially changed (online supplemental file 1).

When stratifying by ulcer severity, we observed that patients with diabetic foot ulcers complicated by osteomyelitis or gangrene were more than twice as likely to be seen by a specialist, compared with those with early stage disease ([figure 3](#)). Rural differences in specialty care also became more pronounced; among those with osteomyelitis, there was an absolute percentage point decrease of 4.83 among rural patients compared with the overall cohort. Among those with osteomyelitis, the difference for those who identified as black compared with the overall cohort was 3.5 absolute percentage points. Observed differences were particularly pronounced for rural patients identifying as black, who experienced a 16.66 absolute percentage point decrease in specialty care when diagnosed with osteomyelitis and a 14.57 absolute percentage point decrease when diagnosed with gangrene, compared with the overall cohort. Given specialty care has been associated with reduced risk of major amputation, this difference is likely to be clinically and statistically significant.<sup>11 31</sup>

## DISCUSSION


We observed lower proportions of specialty care among rural patients compared with the overall, national cohort

**Table 2** Adjusted ORs (aORs) of receiving specialty care for rural patients and patients identifying as black.

		Specialty care,* aOR (95% CI)	
		Patient characteristic	
Model	Model variables	Rural patients	Patients identifying as black
1	Patient characteristic	0.85 (0.81 to 0.88)	1.08 (1.04 to 1.11)
2	Patient characteristic + age + sex	0.83 (0.80 to 0.87)	1.03 (0.99 to 1.07)
3	Rural residence + identifying as black+age + sex + sociodemographic factors <sup>‡</sup>	0.87 (0.84 to 0.91)	1.04 (1.01 to 1.08)
4	Rural residence + identifying as black + age + sex + sociodemographic factors <sup>‡</sup> + comorbidities + ulcer severity	0.83 (0.79 to 0.86)	0.92 (0.89 to 0.97)
5	Rural residence + age + sex identifying as black + sociodemographic factors + comorbidities + ulcer severity + interaction (rural residence × identifying as black)	White rural	Black rural
		0.85 (0.80 to 0.89)	0.61 (0.53 to 0.71)

of patients hospitalized with diabetic foot ulcers. The most striking disparity was revealed by using an intersectionality approach: only 26.23% of rural patients identifying as black were seen by a specialist. This was >5 absolute percentage points lower than the overall national cohort and the overall group identifying as black. It was nearly twice the difference experienced by rural patients in general. Given the importance of specialty care on limb salvage and our conservative comparator group, we think a >5% difference is clinically and socially meaningful.


We identified this difference in the hospitalized setting, however, the intersection of rurality and identifying as black has been associated with decreased access to medical treatment across the healthcare continuum.<sup>32–34</sup>



**Figure 2** Predicted probabilities of receiving specialty care by social identity were calculated from Model 5 (interaction model), specifying the most common patient characteristics from this cohort (eg, a hypothetical 72-year-old patient with diabetes, hyperlipidemia, and hypertension hospitalized with an early stage ulcer).

Improving access to specialty care for rural patients in general might yield profound reductions in major amputation disparities currently faced by rural patients identifying as black.<sup>11 35</sup> Doing so may require policy shifts and programs to address structural inequities within the current healthcare system to either attract and retain specialists in rural, predominantly black communities or facilitate access for rural patients who identify as black to urban-based specialists. We also recognize that healthcare system inequities themselves are likely to be rooted in more long-standing social injustices. Rural educational disadvantages, which are pronounced in school systems serving predominantly black communities, are further exacerbated by migration patterns.<sup>5</sup> Those who remain in rural communities often face limited educational opportunities that translate into restricted employment and health insurance opportunities.<sup>5</sup> Local poverty and lack of insurance make the financial solvency of community hospitals challenging; while 65% of rural American counties are whole or partial health professional shortage areas, this percentage rises to 83% among rural counties with a majority of residents identifying as black.<sup>5</sup> Addressing disparities at the level of the health system may be a partial solution to a much more socially rooted problem.

Barriers to specialty care for rural patients hospitalized with diabetic foot ulcers regardless of race might include: limited access to providers in general and specialists in particular—especially infectious disease physicians and vascular surgeons. Compounding provider shortages are facility closures and an understaffed healthcare workforce.<sup>1 6 8 36</sup> The majority of multidisciplinary limb salvage teams operate in urban, tertiary care centers.<sup>11</sup> Rural patients are likely to have limited access to such teams.<sup>37</sup> More research is needed to identify the exact health system factors driving barriers to care. In addition, research on healthcare policy would be useful to help curtail rural disparities.<sup>6 8 37–40</sup> Currently, our healthcare policies favor large populations due to, among other things: (1) Financial marketing systems that rely on large populations of insured beneficiaries in order to provide



**Figure 3** Observed proportion of patients who received specialty care based on social identities and stratified by ulcer severity. Dashed lines indicate the percentage of the overall cohort who received specialty care stratified by ulcer severity. Vertical bars indicate the decrease from the observed proportions of each social identity group compared with the overall cohort, with the exact percentage noted above the bar.

services, (2) Prioritization of outcomes on the national level, which shifts resource allocation to large population centers, and (3) No method of compensating for innate efficiencies in remote healthcare settings.<sup>6</sup> Policies that recognize and respond to these challenges facing rural communities have the potential to significantly and broadly impact rural disparities, beyond those germane to patients with diabetic foot ulcers.

Rural disparities in specialty care were larger for patients with advanced ulcers, compared with early stage ulcers. This deficit is particularly concerning because patients with osteomyelitis or gangrene are (1) At highest risk of undergoing major amputation, and (2) May be most likely to benefit from specialty care.<sup>41</sup> Arguably, this is the tipping point at which specialty care is critical for limb salvage. Referral systems that prioritize access to specialty care based on ulcer severity may be particularly useful.<sup>35 36</sup> Such a system hinges on frontline providers—primary care providers, urgent care clinicians, emergency medicine specialists, and general medicine admitting teams—accurately staging ulcers and diagnosing peripheral vascular disease.

Despite the strengths of using a complete Medicare population, which significantly reduces internal validity concerns, some limitations should be noted. First, the generalizability of our findings may be limited because our data set was largely composed of older adults and a Medicare-insured population. Second, our data set spanned 2013–2014 and may not reflect current trends. Contemporary referral patterns and telemedicine practices may have shifted access to specialty care, especially amid the COVID-19 pandemic.<sup>42</sup> However, there is evidence that the rural mortality gap, both overall and related to diabetes, is increasing.<sup>6 43</sup> Third, claims data

may introduce misclassification bias or underestimate comorbidities, especially vascular disease. Vascular disease is underdiagnosed and undercoded since less than 25% of patients presenting with gangrenous ulcers previously carried a diagnosis of peripheral vascular disease.<sup>44</sup> These gaps may be larger for marginalized patients, including rural patients, who interact less with the healthcare system and simultaneously have higher rates of smoking that serves as a precursor to vascular disease.<sup>45–47</sup> It may also underestimate or misclassify clinician specialties as physicians may register with Medicare after completing a general residency but before completing their final, specialty fellowships.<sup>22 23</sup> Finally, our study investigated inpatient healthcare factors that might contribute to disparities in major amputations. However, it is unlikely that differences in healthcare delivery alone (whether delivered in the ambulatory or hospitalized setting) fully account for rural disparities in major amputations.<sup>48–50</sup> Future studies should explore additional social determinants of health that may underpin poor outcomes among rural patients with diabetic foot ulcers, such as educational and employment opportunities.

In conclusion, our findings indicate that rural disparities exist in healthcare delivery for hospitalized patients with diabetic foot ulcers. Specifically, a smaller proportion of rural patients receive specialty care. Racial and rural social identities overlap to amplify disparities in specialty care for rural patients identifying as black. Further research is needed to identify specific barriers to care and investigate whether addressing them will help reduce disparities in major amputations.

**Acknowledgements** The authors thank Dr. Chris Crnich, MD, PhD for serving as a sounding board to their findings. The authors also thank Jamie N. LaMantia,



BS, Senior Research Specialist at the University of Wisconsin–Madison School of Medicine and Public Health, Department of Medicine, for editorial assistance.

**Contributors** All authors made substantial contributions to the conception or design of the work; acquisition, analysis, or interpretation of data for the work; drafting the work or revising it critically for important intellectual content; final approval of the version to be published; and agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. The principal investigator (MB) is the guarantor who accepts full responsibility for the finished work and/or the conduct of the study, had access to the data, and controlled the decision to publish.

**Funding** This work was supported by the Agency for Healthcare Research and Quality (grant number 5K08 HS026279), the National Institute of Diabetes and Digestive and Kidney Diseases (grant number 1R01 DK132569-01), and the Diabetes Research Center at Washington University (grant number P30 DK020579). Additional funding was from the National Institute on Minority Health and Health Disparities Research grant R01MD010243 and the Wisconsin Clinical and Translational Science Award (NIH CTSA grant 1UL1TR002373; Bartels and Kind). This material is the result of work also supported with the resources and the use of facilities at the University of Wisconsin School of Medicine and Public Health Center for Health Disparities Research.

**Map disclaimer** The inclusion of any map (including the depiction of any boundaries therein), or of any geographic or locational reference, does not imply the expression of any opinion whatsoever on the part of BMJ concerning the legal status of any country, territory, jurisdiction or area or of its authorities. Any such expression remains solely that of the relevant source and is not endorsed by BMJ. Maps are provided without any warranty of any kind, either express or implied.

**Competing interests** None declared.

**Patient consent for publication** Not applicable.

**Ethics approval** This study involves human participants. The University of Wisconsin Health Sciences Institutional Review Board approved this study and waived written informed consent (2021–0013). Written informed consent was waived because: (1) The study was minimal risk, (2) The study did not adversely affect study participants' rights and welfare because use of the data is not expected to affect patients from whom the data are derived, and (3) It was impractical to obtain informed consent given the national cohort size was quite large and a number of the study participants were likely to be deceased.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data are available upon reasonable request. Due to data use agreements with CMS, summary data will be made available upon reasonable request to the corresponding author.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

#### ORCID iD

Meghan B Brennan <http://orcid.org/0000-0002-2316-4150>

#### REFERENCES

- Brennan MB, Allen GO, Ferguson PD, *et al.* The association between geographic density of infectious disease physicians and limb preservation in patients with diabetic foot ulcers. *Open Forum Infect Dis* 2017;4:ofx015. 10.1093/ofid/ofx015 Available: <https://doi.org/10.1093/ofid/ofx015>
- Skrepnek GH, Mills JL, Armstrong DG. A diabetic emergency one million feet long: disparities and burdens of illness among diabetic foot ulcer cases within emergency departments in the united states, 2006–2010. *PLoS One* 2015;10:e0134914.
- Brennan MB, Powell WR, Kaikow F, *et al.* Association of race, ethnicity, and rurality with major leg amputation or death among medicare beneficiaries hospitalized with diabetic foot ulcers. *JAMA Netw Open* 2022;5:e228399.
- Traven SA, Synovec JD, Walton ZJ, *et al.* Notable racial and ethnic disparities persist in lower extremity amputations for critical limb ischemia and infection. *J Am Acad Orthop Surg* 2020;28:885–92. 10.5435/JAAOS-D-19-00630 Available: <https://doi.org/10.5435/JAAOS-D-19-00630>
- Probst JC, Moore CG, Glover SH, *et al.* Person and place: the compounding effects of race/ethnicity and rurality on health. *Am J Public Health* 2004;94:1695–703. 10.2105/ajph.94.10.1695 Available: <https://doi.org/10.2105/ajph.94.10.1695>
- Probst J, Eberth JM, Crouch E. Structural urbanism contributes to poorer health outcomes for rural America. *Health Aff (Millwood)* 2019;38:1976–84. 10.1377/hlthaff.2019.00914 Available: <https://doi.org/10.1377/hlthaff.2019.00914>
- Fayfman M, Schechter MC, Amobi CN, *et al.* Barriers to diabetic foot care in a disadvantaged population: a qualitative assessment. *Journal of Diabetes and Its Complications* 2020;34:107688. 10.1016/j.jdiacomp.2020.107688 Available: <https://doi.org/10.1016/j.jdiacomp.2020.107688>
- Gong G, Phillips SG, Hudson C, *et al.* Higher US rural mortality rates linked to socioeconomic status, physician shortages, and lack of health insurance. *Health Aff (Millwood)* 2019;38:2003–10. 10.1377/hlthaff.2019.00722 Available: <https://doi.org/10.1377/hlthaff.2019.00722>
- Crenshaw K. Demarginalizing the intersection of race and sex: a black feminist critique of antidiscrimination doctrine, feminist theory and antiracist politics. *Univ Chic Leg Forum* 1989;1989:139–67.
- Bauer GR, Churchill SM, Mahendran M, *et al.* Intersectionality in quantitative research: A systematic review of its emergence and applications of theory and methods. *SSM Popul Health* 2021;14:100798. 10.1016/j.ssmph.2021.100798 Available: <https://doi.org/10.1016/j.ssmph.2021.100798>
- Musuza J, Sutherland BL, Kurter S, *et al.* A systematic review of multidisciplinary teams to reduce major amputations for patients with diabetic foot ulcers. *J Vasc Surg* 2020;71:1433–46.
- Mills JL, Conte MS, Armstrong DG, *et al.* The society for vascular surgery lower extremity threatened limb classification system: risk stratification based on wound, ischemia, and foot infection (wif). *J Vasc Surg* 2014;59:220–34.
- Schaper NC, Netten JJ, Apelqvist J, *et al.* Practical guidelines on the prevention and management of diabetic foot disease (IWGDF 2019 update). *Diabetes Metab Res Rev* 2020;36:e3266. 10.1002/dmrr.3266 Available: <https://onlinelibrary.wiley.com/toc/15207560/36/S1>
- Jarrin OF, Nyandege AN, Grafova IB, *et al.* Validity of race and ethnicity codes in medicare administrative data compared with gold-standard self-reported race collected during routine home health care visits. *Med Care* 2020;58:e1–8. 10.1097/MLR.0000000000001216 Available: <https://doi.org/10.1097/MLR.0000000000001216>
- Johnston KJ, Wen H, Joynt Maddox KE. Lack of access to specialists associated with mortality and preventable hospitalizations of rural Medicare beneficiaries. *Health Affairs* 2019;38:1993–2002. 10.1377/hlthaff.2019.00838 Available: <https://doi.org/10.1377/hlthaff.2019.00838>
- Kind AJH, Buckingham WR. Making neighborhood-disadvantage metrics accessible - the neighborhood atlas. *N Engl J Med* 2018;378:2456–8. 10.1056/NEJMp1802313 Available: <https://doi.org/10.1056/NEJMp1802313>
- WWAMI Rural Health Research Center. Rural urban commuting area codes maps classifications. Available: <https://depts.washington.edu/uwruca/ruca-maps.php> [Accessed 23 Mar 2022].
- Centers for Medicare & Medicaid Services. *Chronic conditions data warehouse: conditions categories*. 2021. Available: <https://www2.cdwdata.org/web/guest/condition-categories> [accessed 3 Mar 2021].
- Fincke BG, Miller DR, Turpin R. A classification of diabetic foot infections using ICD-9-CM codes: application to a large computerized medical database. *BMC Health Serv Res* 2010;10:192. 10.1186/1472-6963-10-192 Available: <https://doi.org/10.1186/1472-6963-10-192>
- Bartels CM, Kind AJH, Everett C, *et al.* Low frequency of primary lipid screening among medicare patients with rheumatoid arthritis. *Arthritis Rheum* 2011;63:1221–30. 10.1002/art.30239 Available: <https://doi.org/10.1002/art.30239>



- 21 von Elm E, Altman DG, Egger M, *et al.* The strengthening the reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *International Journal of Surgery* 2014;12:1495–9. 10.1016/j.ijsu.2014.07.013 Available: <https://doi.org/10.1016/j.ijsu.2014.07.013>
- 22 Baldwin L-M, Adamache W, Klabunde CN, *et al.* Linking physician characteristics and medicare claims data: issues in data availability, quality, and measurement. *Med Care* 2002;40(8 Suppl):IV–82. 10.1097/00005650-200208001-00012 Available: <https://doi.org/10.1097/00005650-200208001-00012>
- 23 DesRoches CM, Barrett KA, Harvey BE, *et al.* The results are only as good as the sample: assessing three national physician sampling frames. *J GEN INTERN MED* 2015;30:595–601. 10.1007/s11606-015-3380-9 Available: <https://doi.org/10.1007/s11606-015-3380-9>
- 24 Washington State Department of Health. Guidelines for using rural-urban classification systems for community health assessment. In: *Health WSDo*. 2016: 26.
- 25 Eicheldinger C, Bonito A. More accurate racial and ethnic codes for medicare administrative data. *Health Care Financ Rev* 2008;29:27–42.
- 26 Elixhauser A, Steiner C, Harris DR, *et al.* Comorbidity measures for use with administrative data. *Medical Care* 1998;36:8–27. 10.1097/00005650-199801000-00004 Available: <https://doi.org/10.1097/00005650-199801000-00004>
- 27 Li P, Kim MM, Doshi JA. Comparison of the performance of the CMS hierarchical condition category (CMS-HCC) risk adjuster with the Charlson and Elixhauser comorbidity measures in predicting mortality. *BMC Health Serv Res* 2010;10:245. 10.1186/1472-6963-10-245 Available: <https://doi.org/10.1186/1472-6963-10-245>
- 28 Guan A, Thomas M, Vittinghoff E, *et al.* An investigation of quantitative methods for assessing intersectionality in health research: a systematic review. *SSM Popul Health* 2021;16:100977.
- 29 R Core Team. *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing, 2021.
- 30 The Rural Health Research Center. *RUCA maps*. 2006. Available: <https://depts.washington.edu/uwruca/ruca-maps.php>
- 31 Brennan M, Liu Y, LaMantia J, *et al.* Associations between specialty care, limb salvage, and survival among patients with diabetic foot ulcers. In: *Diabetic Foot Conference (DFCon)*. Los Angeles, CA, 2022.
- 32 Goodney PP, Travis LL, Nallamothu BK, *et al.* Variation in the use of lower extremity vascular procedures for critical limb ischemia. *Circ Cardiovasc Qual Outcomes* 2012;5:94–102. 10.1161/CIRCOUTCOMES.111.962233 Available: <https://doi.org/10.1161/CIRCOUTCOMES.111.962233>
- 33 Humphries MD, Brunson A, Li CS, *et al.* Amputation trends for patients with lower extremity ulcers due to diabetes and peripheral artery disease using statewide data. *J Vasc Surg* 2016;64:S0741-5214(16)30830-8:1747–55. 10.1016/j.jvs.2016.06.096 Available: <https://doi.org/10.1016/j.jvs.2016.06.096>
- 34 Prompers L, Huijberts M, Apelqvist J, *et al.* High prevalence of ischaemia, infection and serious comorbidity in patients with diabetic foot disease in europe. baseline results from the eurodiale study. *Diabetologia* 2007;50:18–25. 10.1007/s00125-006-0491-1 Available: <https://doi.org/10.1007/s00125-006-0491-1>
- 35 McGilliv, Constantino M, Bolton T, *et al.* Diabetes amputation program: a structured systematic approach. *Diabetic Foot Journal* 2003;6:172–6.
- 36 Newhall K, Spangler E, Dzebisashvili N, *et al.* Amputation rates for patients with diabetes and peripheral arterial disease: the effects of race and region. *Annals of Vascular Surgery* 2016;30:292–298. 10.1016/j.avsg.2015.07.040 Available: <https://doi.org/10.1016/j.avsg.2015.07.040>
- 37 Sutherland BL, Pecanac K, Bartels CM, *et al.* Expect delays: poor connections between rural and urban health systems challenge multidisciplinary care for rural americans with diabetic foot ulcers. *J Foot Ankle Res* 2020;13.
- 38 Adcock AK, Choi J, Alvi M, *et al.* Expanding acute stroke care in rural America: a model for statewide success. *Telemed J E Health* 2020;26:865–71. 10.1089/tmj.2019.0087 Available: <https://doi.org/10.1089/tmj.2019.0087>
- 39 Bauer GR, Scheim AI. Advancing quantitative intersectionality research methods: intracategorical and intercategorical approaches to shared and differential constructs. *Social Science & Medicine* 2019;226:260–2. 10.1016/j.socscimed.2019.03.018 Available: <https://doi.org/10.1016/j.socscimed.2019.03.018>
- 40 Swayze EJ, Strzyzewski L, Avula P, *et al.* The impact of expanding gynecologic oncology care to ovarian cancer patients in small cities and rural communities. *Gynecol Oncol* 2021;161:S0090-8258(21)00331-0:852–7. .
- 41 Zahnd WE, Murphy C, Knoll M, *et al.* The intersection of rural residence and minority race/ethnicity in cancer disparities in the United States. *Int J Environ Res Public Health* 2021;18:1384. 10.3390/ijerph18041384 Available: <https://doi.org/10.3390/ijerph18041384>
- 42 Anichini R, Cosentino C, Papanas N. Diabetic foot syndrome in the COVID-19 era: how to move from classical to new approaches. *The International Journal of Lower Extremity Wounds* 2022;21:107–10. 10.1177/15347346221081572 Available: <https://doi.org/10.1177/15347346221081572>
- 43 Callaghan T, Ferdinand AO, Akinlotan MA, *et al.* The changing landscape of diabetes mortality in the United States across region and rurality, 1999–2016. *J Rural Health* 2020;36:410–5. 10.1111/jrh.12354 Available: <https://doi.org/10.1111/jrh.12354>
- 44 Brennan MB, Hess TM, Bartle B, *et al.* Diabetic foot ulcer severity predicts mortality among veterans with type 2 diabetes. *Journal of Diabetes and Its Complications* 2017;31:556–61. 10.1016/j.jdiacomp.2016.11.020 Available: <https://doi.org/10.1016/j.jdiacomp.2016.11.020>
- 45 Coughlin LN, Bonar EE, Bohnert KM, *et al.* Changes in urban and rural cigarette smoking and cannabis use from 2007 to 2017 in adults in the united states. *Drug Alcohol Depend* 2019;205:107699.
- 46 Dansky KH, Dirani R. The use of health care services by people with diabetes in rural areas. *J Rural Health* 1998;14:129–37. 10.1111/j.1748-0361.1998.tb00614.x Available: <https://doi.org/10.1111/j.1748-0361.1998.tb00614.x>
- 47 Doogan NJ, Roberts ME, Wewers ME, *et al.* A growing geographic disparity: rural and urban cigarette smoking trends in the United States. *Preventive Medicine* 2017;104:79–85. 10.1016/j.ypmed.2017.03.011 Available: <https://doi.org/10.1016/j.ypmed.2017.03.011>
- 48 Anderson SG, Shoo H, Saluja S, *et al.* Social deprivation modifies the association between incident foot ulceration and mortality in type 1 and type 2 diabetes: a longitudinal study of a primary-care cohort. *Diabetologia* 2018;61:959–67. 10.1007/s00125-017-4522-x Available: <https://doi.org/10.1007/s00125-017-4522-x>
- 49 Ferguson HJM, Nightingale P, Pathak R, *et al.* The influence of socio-economic deprivation on rates of major lower limb amputation secondary to peripheral arterial disease. *Eur J Vasc Endovasc Surg* 2010;40:76–80. 10.1016/j.ejvs.2010.03.008 Available: <https://doi.org/10.1016/j.ejvs.2010.03.008>
- 50 Riley J, Antza C, Kempegowda P, *et al.* Social deprivation and incident diabetes-related foot disease in patients with type 2 diabetes: A population-based cohort study. *Diabetes Care* 2021;44:731–9. 10.2337/dc20-1027 Available: <https://doi.org/10.2337/dc20-1027>

**Appendix A.****Supplemental 1.****Table S1.** Medical Specialty Codes/Provider Taxonomy Codes Used to Identify Specialists in the Medicare Dataset

Specialty	Medical Specialty codes	Provider Taxonomy codes
Endocrinology	11	207RE0101X
Infectious disease	11, 44	207RI0200X
Orthopedic surgery	2, 20	207X00000X, 207XS0114X, 207XX0004X, 207XS0117X, 207XX0801X, 207XP3100X, 207XX005X, 207XS0106X
Plastic surgery	2, 24	2086S0122X, 2082S0099X, 2082S0105X, 208200000X
Podiatry	48	213E00000X, 213ES0103X, 213ES0131X, 213EG0000X, 213EP1101X, 213EP0504X, 213ER0200X, 213ES0000X
Vascular surgery	2, 77	2086S0129X

**Supplemental 2. Sensitivity Analysis of alternative rurality coding scheme.****Table S2.** Alternative RUCA categorization scheme

Category	RUCA codes	Number of patients per code	Number discharged from critical access hospital (%)
Urban	1.0, 1.1, 2.0, 2.1, 3.0, 4.1, 5.1, 7.1, 8.1, 10.1	105 245	1040 (0.99)
Large Rural	4.0, 4.2, 5.0, 5.2, 6.0, 6.1	9596	359 (3.74)
Small Rural/isolated	7.0, 7.2, 7.3, 7.4, 8.0, 8.2, 8.3, 8.4, 9.0, 9.1, 9.2, 10.0, 10.2, 10.3, 10.4, 10.5, 10.6	9112	1334 (14.64)

**Table S3.** Observed primary and secondary outcomes by alternative RUCA categorization scheme

Characteristic	Full cohort* (n=124 487) (100%)	Rurality		
		Urban patients (n=105 245) (84.54%)	Large Rural (n=9596) (7.71%)	Small Rural/isolated (n=9112) (7.32%)
Outcome				
Seen by at least 1 specialist (primary outcome)	40 027 (32.15)	34 574 (32.85)	2671 (27.83)	2699 (29.62)



**Table S4.** Odds ratios for specialty care using alternative RUCA scheme

Outcome		At least one specialist, * aOR (95% CI)	
		Patient characteristic	
Model	Model variables	Small rural/isolated residence	Identifying as Black
1	Patient characteristic	0.86 (0.82-0.90)	1.08 (1.04-1.11)
2	Patient characteristic+ age + sex	0.85 (0.81-0.89)	1.03 (0.99-1.07)
3	Rural residence + identifying as Black + age + sex + sociodemographic factors <sup>†</sup>	0.89 (0.85-0.94)	1.04 (1.01-1.08)
4	Rural residence + identifying as Black + age + sex + sociodemographic factors <sup>†</sup> + comorbidities + ulcer severity	0.85 (0.81-0.90)	0.92 (0.89-0.97)
5	Rural residence + age + sex identifying as Black + sociodemographic factors + comorbidities + ulcer severity + interaction (rural residence x identifying as Black)	White	Black
		0.87 (0.80-0.89)	0.61 (0.51-0.74)