

Presentation and characteristics of children with screen-detected type 1 diabetes: learnings from the ELSA general population pediatric screening study

Lauren M Quinn ¹, Renuka P Dias,^{2,3} Christopher Bidder,⁴ Sudeshna Bhowmik,⁵ Kerstin Bumke,⁶ Jaikumar Ganapathi,⁷ Shaun Gorman,⁸ Edward Hind,⁹ Swati Karandikar,¹⁰ Kiran Kumar,¹¹ Nicholas Lipscomb,¹² Sheila McGovern,¹³ Vijith R Puthi,¹⁴ Tabitha Randell,¹⁵ Gemma Watts,¹⁶ Parth Narendran ^{1,17}

To cite: Quinn LM, Dias RP, Bidder C, *et al*. Presentation and characteristics of children with screen-detected type 1 diabetes: learnings from the ELSA general population pediatric screening study. *BMJ Open Diab Res Care* 2024;**12**:e004480. doi:10.1136/bmjdr-2024-004480

► Additional supplemental material is published online only. To view, please visit the journal online (<https://doi.org/10.1136/bmjdr-2024-004480>).

Received 22 July 2024
Accepted 5 September 2024



© Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to
Dr Lauren M Quinn;
l.quinn.1@bham.ac.uk

ABSTRACT

Introduction We describe the identification and management of general population screen-detected type 1 diabetes (T1D) and share learnings for best practice.

Research design and methods Children diagnosed with T1D through a general population screening initiative, the EarLy Surveillance for Autoimmune diabetes (ELSA) study, were reviewed and described.

Parents provided written, informed consent for inclusion in the case series.

Results 14 children with insulin requiring (stage 3) T1D are described. These cases offer unique insights into the features of screen-detected T1D. T1D is identified sooner through screening programs, characterized by absent/short symptom duration, median presenting glycated hemoglobin 6.6% (49 mmol/mol) and insulin requirements <0.5 units/kg/day. ELSA identified four children at stage 3 and another 4 progressed within 4 months of ELSA completion, including two single seropositive children. Six children developed stage 3 T1D prior to ELSA completion, including two children (14%, n=2/14) with diabetic ketoacidosis prior to confirmed antibody status.

Conclusions There are three main learnings from this case series. First, T1D identified through screening is at an earlier stage of its natural history and requires personalized insulin regimens with lower total daily insulin doses. Second, single autoantibody seropositivity can rapidly progress to stage 3. Finally, insulin requirement can manifest at any stage of the T1D screening pathway, and therefore early education around symptom recognition is essential for families participating in screening programs.

INTRODUCTION

Type 1 diabetes (T1D) is an autoimmune condition caused by destruction of the insulin-producing pancreatic islet beta cells.¹ Screening for islet-specific autoantibodies (Aab) in otherwise healthy children enables early identification of T1D up to 15 years before symptomatic onset of disease.² The

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ As increasing numbers of children are screened for type 1 diabetes (T1D), understanding their transition from research pathway into clinical care is important.

WHAT THIS STUDY ADDS

⇒ We describe the identification and management of the first 14 stage 3 T1D cases from the UK's Early Surveillance for Autoimmune diabetes (ELSA) study to share learnings for best practice. This paper adds three key messages to the previous literature reporting clinically milder phenotype of screen-detected T1D.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ These learnings from the ELSA study are important for healthcare professionals and policymakers to raise awareness and increase confidence in managing presymptomatic T1D.

presence of a single Aab associates with a 14.5% progression to insulin requiring T1D over a 10-year period.³ However, the presence of two or more Aab indicates a lifetime certainty of requiring insulin and has been termed T1D even before the onset of hyperglycemia.^{4 5} Children identified with two or more Aab can be subsequently staged with oral glucose tolerance testing (OGTT): normoglycemia (stage 1), dysglycemia (stage 2), non-insulin requiring hyperglycemia without osmotic symptoms (stage 3a) and insulin requiring with symptoms (stage 3b).⁵

Pediatric T1D screening studies are being set up internationally, including the EarLy Surveillance for Autoimmune diabetes




Figure 1 Summary of stage 3 cases identified in the ELSA study. Figure outlining where stage 3 cases were identified during the ELSA study. Four children were diagnosed after study enrollment and prior to DBS results notification (cases 10–13). One child was diagnosed following DBS results notification but prior to venous autoantibody (Aab) confirmation (case 14). Of these, two children had mild DKA. Two children progressed to stage 3 within 2 months of single Aab results notification (cases 7–8) and one child progressed to stage 3 prior to metabolic staging (case 9). Four children had stage 3 T1D from the ELSA OGTT (cases 1–4). Two children progressed from presymptomatic to symptomatic T1D within 6 months of ELSA staging (cases 5–6). DBS, dried blood spot; DKA, diabetic ketoacidosis; ELSA, EarLy Surveillance for Autoimmune diabetes; mAab, multiple autoantibody positive; OGTT, oral glucose tolerance testing; sAab, single autoantibody positive; T1D, type 1 diabetes.

(ELSA) study in the UK.⁶ Screening achieves a fivefold reduction in diabetic ketoacidosis (DKA),⁷ in addition to shorter duration or absence of symptoms, lower glycosylated hemoglobin (HbA1c) and lower initial insulin requirements.⁸ In contrast, up to 58% of children identified in usual care present in DKA,⁹ frequently due to late or missed diagnoses.^{10–12}

As increasing numbers of children are screened (>200,000 internationally),⁶ the transition of screen-detected cases from research into clinical care becomes relevant to healthcare providers. We wanted to describe children diagnosed at stage 3 T1D from the UK ELSA study and share key learnings for best practice management of screen-detected T1D.

RESEARCH DESIGN AND METHODS

The ELSA study aims to explore the feasibility and acceptability of screening children aged 3–13 years for T1D (www.elsadiabetes.nhs.uk) (figure 1). Recruitment approaches include home-testing, community clinics (schools, primary care, community centers) and hospitals and commenced in November 2022. The screening test is the RSR Limited 3-screen ELISA for anti-islet antigen 2 (IA-2A), anti-glutamic acid decarboxylase (anti-GAD) and anti-Zinc transporter 8 (ZnT8) Aab¹³ undertaken on a dried blood spot (DBS) collected on filter paper. Those screening positive on the DBS test are offered a confirmation on a venous blood sample measuring IA-2A, insulin, GAD and ZnT8 Aab. Children with a single Aab are followed up yearly with Aab testing in the follow-on Innovative approach towards understanding and arresting type 1 diabetes (INNODIA) study.¹⁴ Those with two or more Aab are staged with OGTT. Those who are stage 3

are referred directly into clinical care for insulin initiation, and those with stages 1 and 2 are signposted to the follow-up INNODIA program¹⁴ for 6 monthly monitoring until they progress to stage 3 T1D.

A patient and public involvement group, comprising both parents and children, contributed to the design of the ELSA study and development of participant information including leaflets, videos and educational resources.

We obtained written informed consent from parents for inclusion in the case series. Stage 3 cases identified between recruitment onset and April 2024 are included. Demographic data, including child's ethnicity, postcode for Index of Multiple Deprivation (IMD)¹⁵ and family history of diabetes, were collected at time of entry to the ELSA study. We contacted the child's lead consultant pediatrician who provided discharge summaries, clinic letters, laboratory investigations and ambulatory glucose profiles reports since T1D diagnosis. Parents were contacted to clarify the case histories. T1D and DKA diagnoses met International Society for Pediatric and Adolescent Diabetes (ISPAD) guidelines.¹⁶ HbA1c \geq 6.5% (\geq 48 mmol/mol) indicates stage 3 T1D. Children diagnosed with T1D are managed by a multidisciplinary care team, including pediatric consultant, diabetes specialist nurse, dietician and access to clinical psychology services, as recommended by National Institute for Health and Care Excellence (NICE) guidelines.¹⁷

RESULTS

Over 10,000 children have completed the ELSA study at the time of preparing this case series in June 2024. We report on the first 14 cases of stage 3 T1D identified during or following ELSA study participation. Of these,

71% were boys, median age 8 years (range 3–12 years), 86% were White European ethnicity (British, Irish or other) and 86% had a family history of T1D (first-degree or second-degree relative). Of the 10 children living in England, the median IMD level was 7 (range 1–9) (2 children lived in Northern Ireland, 1 in Scotland and 1 in Wales).

Of the 14 stage 3 T1D cases, 6 were diagnosed prior to ELSA study completion, 4 were screen detected as stage 3 at the time of ELSA metabolic staging and 4 progressed within 4 months of ELSA staging (see [figure 1](#) and online supplemental table 1 for case summaries).

Three key messages emerged: (1) screening identifies stage 3 T1D early, (2) single Aab can progress to stage 3 and (3) insulin-requiring stage 3 T1D emerges at any point of the screening pathway.

SCREENING IDENTIFIES CHILDREN WITH EARLY STAGE 3 T1D

Screen-detected T1D identifies children at an earlier stage of its natural history. Twelve children (cases 1–12, 86%) had absent or short (typically 1–2 weeks) duration of osmotic symptoms, a median HbA1c of 6.6% (range 5.5%–15.1%) (49 mmol/mol (range 37–142 mmol/mol)) and avoided DKA.

Of the ELSA stage 3 screen-detected cases (n=4), two were stage 3a (asymptomatic) and two stage 3b (symptomatic),⁵ all with HbA1c <6.3% (< 45 mmol/mol). The stage 3a cases (1 and 2) avoided hospital admission and commenced continuous glucose monitoring (CGM) under paediatric diabetes specialists. Case 1 progressed to stage 3b after 6 months and started multiple daily injections (MDI) specified dose with evening meal (total daily dose (TDD) <0.5 units/kg/day). Case 2 never experienced osmotic symptoms; instead, following increased CGM time above range, commenced insulin aspart set-dose with breakfast and split set-dose with evening meal (TDD <0.5 units/kg/day).

ELSA stage 3b cases had intermittent osmotic symptoms prior to ELSA OGTT, but home capillary blood glucose (CBG) readings were <11.1 mmol/L. Case 3 was admitted for one night and started MDI (TDD=0.3 units/kg/day). Following discharge, frequent hypoglycemia led to dose reduction and split dosing with evening meal. Case 4 had a weekend admission (three nights) following stage 3b identification. Insulin glargine commencement (1 unit) in hospital led to hypoglycemia and no further insulin was given during admission. Insulin aspart 0.5–1 unit with evening meal (1:20) (TDD <0.5 units/kg/day) was prescribed on discharge.

Follow-up of those identified at risk facilitates earlier stage 3 identification

One stage 2 (case 5) and one stage 1 child (case 6) progressed to stage 3b within 6 months of ELSA completion and prior to research follow-up. One parent (case 5) declined INNODIA follow-up and was referred to the local diabetes team, with remote support including joint

consultation with the ELSA clinical team. Recommendations for stage 2 management in clinical care followed the European guidelines, included CBG meter provision and training, 3 monthly HbA1c and intermittent CGM.^{18 19}

There were frequent telephone contacts and one paediatric assessment unit attendance for hyperglycemia which resolved within 2 hours without insulin. A few months later, after 2 days of osmotic symptoms, stage 3b T1D was diagnosed, avoiding DKA and overnight hospital admission, and ultra-rapid acting insulin aspart specified dose commenced with evening meals (<0.5 units/kg/day).

SINGLE AAB CAN PROGRESS TO STAGE 3

Two single Aab-positive children developed stage 3b T1D within 2 months of completing ELSA. Case 7 had intermittent osmotic symptoms which progressed during the ELSA study. Following antibody results notification, the ELSA paediatrician referred for urgent same day assessment and insulin was commenced. Case 8 developed a viral illness 1 month after ELSA and following a 2-day history of osmotic symptoms and high home CBG readings, parent contacted the local paediatric diabetes team for insulin initiation on an outpatient basis.

Insulin-requiring stage 3 T1D emerges at any stage of the screening pathway

Having entered the ELSA study, two cases presented with stage 3b T1D in DKA prior to completion. Case 13 developed osmotic symptoms prior to ELSA DBS results notification (screened positive) leading to emergency department attendance. This was mild DKA (pH 7.21, ketones 6.4 mmol/L) based on ISPAD 2022 criteria (mild DKA pH 7.2–7.3)¹⁶ with overnight admission. MDI regimen was commenced but due to frequent hypoglycemia, automated insulin delivery was started within 4 months of diagnosis, leading to an HbA1c of 6.5% (47 mmol/mol).

Case 14 screened positive in an ELSA community clinic. Language and travel barriers caused delays in confirmatory Aab testing and two appointment non-attendances. Parental concerns regarding polydipsia and increasing fatigue led to appropriate attendance at the emergency department. The child was diagnosed with mild DKA (pH 7.21, ketones 6.8 mmol/L),¹⁶ with HbA1c 12.9% (118 mmol/mol) and admitted for 8 days for insulin initiation and structured education due to significant language barriers. The child was discharged on TDD >0.5 units/kg/day and HbA1c improved to 5.5% (37 mmol/mol) within 3 months of diagnosis.

DISCUSSION

This case series illustrates how screening facilitates earlier identification of stage 3 disease, reminds us that children with single Aab rapidly progress to requiring insulin, and hence benefit from education and follow-up, and finally, that the requirement for insulin can manifest at any stage of the T1D screening pathway. These cases offer

unique insights into the characteristic features of screen-detected T1D.

The strength of this early cases series is the description of the first 14 stage 3 cases emerging from the ELSA study, which provide important learnings for clinicians managing early T1D and for organizations delivering population screening programs. While the clinically milder phenotype of screen-detected T1D is well described by Fr1da⁸ and others,^{20–22} our three messages build on this literature and offer specific examples for best practice management. As increasing numbers of children are screen detected and if screening programs are nationally adopted as in Italy,^{23,24} the research protocols will need to integrate into pediatric clinical services.²⁵ Hence, there is an urgent need to raise awareness and increase healthcare professionals' confidence managing these presymptomatic T1D cases.

Screening facilitates early stage 3 diagnosis

Screening, combined with education and follow-up, is the most effective way to prevent DKA.^{7,22,26} Fr1da (German general population T1D screening study)²⁷ has achieved the lowest DKA rate at 2.5% (n=3/128)^{7,8} and in this case series, all eight cases completing the ELSA study avoided DKA. Long-term benefits of screen-detected T1D include improved glucose control for at least the first 5 years after diagnosis^{28,29} and reduced parental stress at insulin commencement compared with diagnosis through usual care.³⁰

Further, screen-detected T1D leads to identification at an earlier stage of its natural history. ELSA and Fr1da's glucose metrics at stage 3 are comparable,⁸ including HbA1c (6.6% (49 mmol/mol) vs 6.8% (51 mmol/mol)), fasting glucose (4.7 vs 5.3 mmol/L) and proportion not requiring insulin immediately (86% vs 72%). Compared with unscreened populations (Europe/UK), HbA1c at diagnosis (without DKA) ranges from 10.1% to 10.5% (87–91 mmol/mol)^{8,31} and 98% require insulin immediately.⁸ Our series supports Fr1da's findings that screen-detected cases experience a clinically milder presentation at stage 3.⁸

ELSA study participation expedited stage 3 diagnosis in seven cases. When stage 3 T1D diagnosis occurred independently of ELSA participation (n=5), all children had a sibling or parent with T1D. Hence, osmotic symptom emergence combined with access to home CBG testing confirmed hyperglycemia and led to appropriate healthcare attendance. Four children avoided overnight admission, largely due to a first degree relative with T1D (sibling or parent) already educated in glucose monitoring and insulin administration.

Detection in presymptomatic stages identifies individuals for immunoprevention trials and treatments. The first agent licensed in the USA, teplizumab,³² which delays stage 3 onset,³³ is undergoing NICE consultation for UK licensing, with a decision expected in November 2024.³⁴

Single Aab progress to stage 3

The risk of progression to stage 3 T1D for single islet Aab is 14.5% (95% CI: 10.3% to 18.7%) in 10 years.³ Factors associated with increased rate of progression include IA-2A seropositivity (40.5% progress to T1D within 10 years (CI: 17.7% to 63.3%)),³ younger age at Aab detection^{35,36} and high T1D genetic risk score.^{35,37} In preschool children, a single seropositive Aab is most likely to progress to multiple within 2 years.^{35,38} Yet, single Aab results present uncertain significance and may provoke unnecessary health anxiety for a transiently occurring Aab. As such, single Aab results are not reported or followed-up in Fr1da or European action for the Diagnosis of Early Non-clinical Type 1 diabetes For disease Interception (EDENTIFI), the European population screening collaboration. In ELSA, our experience shows Aab families benefit from education similar to multiple seropositivity, including T1D symptom counselling and follow-up to risk stratify according to Aab status.³⁵ Acceptability assessments via qualitative interviews are ongoing to explore the benefits and harms of single Aab notification.³⁹ The recent Juvenile Diabetes Research Foundation (JDRF) guidance recommends monitoring for at least 2 years after initial seroconversion. For preschool children (<3 years), 6 monthly metabolic monitoring is recommended for 3 years, whereas for children ≥3 years, Aab status and metabolic monitoring are recommended annually for 3 years.⁴⁰

Follow-up of screen-detected T1D supports earlier stage 3 detection

Follow-up after screening is recommended for appropriate timing of insulin initiation, DKA prevention at onset and to facilitate entry to immunoprevention trials.^{5,7,41} European guidelines¹⁸ and an international consensus report⁴⁰ are now published for managing pre-stage 3 T1D and both recommend specialist follow-up according to T1D stage and age of child. To reduce invasive procedures and improve follow-up adherence,⁴² OGTTs are recommended less frequently than former research protocols (eg, INNODIA) and replaced with HbA1c and random plasma glucose 3–6 monthly. All multiple Aab-positive children should receive a CBG meter and training, with testing recommended for osmotic symptoms or febrile illness.¹⁸ This provides a rapid point-of-care test for families to inform appropriate health-seeking behaviors. In our experience, selected cases also benefit from clinical follow-up. As exemplified by case 5, some children at stage 2 who are symptomatic cannot wait for 12 monthly OGTT through INNODIA. There is need for clinical support for these families in between research visits. Our post-ELSA pathway recommends referral for children at stage 2 with intermittent osmotic symptoms, or if the family decline research follow-up, for example, due to geographical barriers. This facilitates safe management of presymptomatic T1D from experienced professionals and mitigates families' distress.

Arguably, a landmark advance is the release of an internationally authorized SNOMED code for 'presymptomatic T1D (disorder)' (1290118005), recommended for inclusion in the child's electronic health record. The International Classification of Diseases—10 similarly recognizes presymptomatic T1D with the following codes:

1. E10.A0 (T1D, presymptomatic, unspecified).
2. E10.A1 (T1D, presymptomatic, stage 1).
3. E10.A2 (T1D, presymptomatic, stage 2).

These codes aim to facilitate earlier diagnosis for high-risk children (Aab-positive stage 1–2) at future health-care contacts.

Managing early stage 3

Screen-detected T1D cases have lower insulin requirements than standard initiation protocols. The Environmental Determinants of Diabetes in the Young (TEDDY), the largest prospective study of children genetically predisposed to T1D⁴³ and Fr1da, showed this correlated with higher C-peptide at stage 3 detection and persists for up to 1 year after diagnosis, indicating partial remission phase.^{8 20 44} As such, screen-detected cases require personalized insulin initiation plans, with reduced TDD and less frequent MDI. Here, the priority is hypoglycemia avoidance. A set bolus given with the highest carbohydrate load of the day (usually evening meal) may suffice if basal insulin is not commenced. A recent Lancet comment recognized lack of empirical evidence for insulin initiation in early stage 3 and recommended a treat to target approach to match the child's physiology, including basal commencement for nocturnal hyperglycemia or prandial insulin for recurrent postprandial hyperglycemia >11.1 mmol/L (>200 mg/dL) for >2 hours or high carbohydrate portions.⁴⁵

Insulin-requiring stage 3 T1D can emerge at any time during the screening pathway

Two main learning points emerged from the DKA cases (14%, n=2/14). First, case 14 had known risk factors for DKA, including no family history of T1D, minority ethnicity and high deprivation^{46 47} demonstrating need for symptom vigilance from screening enrollment.¹¹ An important new finding from ELSA is that parents may enter a screening program due to T1D symptom concerns. In ELSA, from the first 10,202 children screened, 6 (0.06%) were diagnosed stage 3 (1 DKA) prior to study completion (Aab confirmation or OGTT), while in Fr1da, 19 out of 90,632 (0.02%) children were diagnosed stage 3 prior to Aab confirmation.⁷ Evidently, screening protocol timelines must be strictly adhered to and facilitate timely transition between follow-ups (screening test, confirmation and staging). Also, signposting and counselling must be incorporated early in any screening program to ensure medical services are appropriately accessed when symptoms arise. This ensures population screening is reserved for risk identification for the asymptomatic target population. However, screening

also offers important opportunities to educate about symptoms of T1D. In ELSA, this information is presented at screening enrollment and again by a healthcare provider at results notification to facilitate earlier stage 3 diagnosis. Families are advised not to wait for screening results if there are immediate concerns about emerging T1D. These essential safety measures must be integrated into population screening programs.

In conclusion, we report three key messages emerging from the first 14 stage 3 T1D cases detected during and following participation in the UK's pediatric screening study (ELSA). Screening, education and monitoring facilitate earlier stage 3 diagnosis, characterized by milder clinical presentation and personalized insulin regimens with lower insulin requirements. Single Aab can progress to stage 3, and therefore benefit from follow-up. Finally, education, including early symptom recognition, is essential to avoid late presentation for children enrolled in a screening program. We envisage that these learnings will support best practice dissemination at a time when screening studies are contributing to increasing numbers of T1D diagnoses.

Author affiliations

¹Institute of Immunology and Immunotherapy, University of Birmingham, Birmingham, UK

²Institute of Applied Health Research, University of Birmingham, Birmingham, UK

³Department of Paediatric Endocrinology, Birmingham Women's and Children's Hospitals NHS Foundation Trust, Birmingham, UK

⁴Department of Child Health, Swansea Bay University Health Board, Morriston Hospital, Swansea, UK

⁵Department of Paediatrics, Arrowse Park Hospital, Wirral, UK

⁶Paediatric Department, University Hospital Wishaw, Wishaw, UK

⁷Hillingdon Hospitals NHS Foundation Trust, Uxbridge, UK

⁸Bradford Teaching Hospitals NHS Foundation Trust, Bradford, UK

⁹North Hampshire Hospital, Basingstoke, UK

¹⁰Birmingham Heartlands Hospital, Birmingham, UK

¹¹Burton Hospitals NHS Foundation Trust, Derby, UK

¹²Department of Paediatrics, South West Acute Hospital, Enniskillen, UK

¹³South Eastern Health and Social Care Trust, Dundonald, UK

¹⁴Department of Paediatrics, Peterborough City Hospital, Peterborough, UK

¹⁵Nottingham Children's Hospital, Nottingham, UK

¹⁶Queen Alexandra Hospital, Portsmouth, UK

¹⁷Department of Diabetes, University Hospitals Birmingham NHS Foundation Trust, Birmingham, UK

Acknowledgements We would like to thank the parents, children and young people for their participation in the ELSA study and inclusion in the case series. We thank the ELSA study team at the University of Birmingham and the principal investigators at the ELSA follow-up testing sites for their contributions: Dr Rachel Besser, Oxford Children's Hospital, Oxford, UK; Dr Fiona Campbell, Leeds Teaching Hospitals NHS Trust, Leeds, UK; Dr Renuka Dias, Birmingham Children's Hospital, Birmingham, UK; Dr Chris Gardner, East Lancashire Hospitals NHS Trust, Burnley, UK; Dr Evelien Gevers, Barts Health, Royal London Children's Hospital, London, UK; Dr Loredana Marcovecchio, Cambridge Children's Hospital, Cambridge, UK; Dr Muriel Meso, Royal Free NHS Trust, London, UK; Dr Sarinda Miller, Southern Health and Social Care Trust, Northern Ireland, UK; Dr Chris Moudiotis, Royal Devon and Exeter NHS Foundation Trust, Exeter, UK; Dr Catherine Owen, Newcastle Hospitals NHS Foundation Trust, Newcastle, UK; Dr Tabitha Randell, Nottingham Children's Hospital, Derby Road, Nottingham, UK; Dr Rajeeb Rashid, Children's and Young People's Diabetes Service, Royal Hospital for Children, Glasgow, UK; Dr Shilpa Shah, Southern Health and Social Care Trust, Northern Ireland, UK; Dr Ambika Shetty, Noah's Ark Children's Hospital for Wales, UK; Dr Nicola Trevelyan,

Southampton Children's Hospital, Southampton, UK; Dr Ruben Willemson, Barts Health, Royal London Children's Hospital, London, UK

Contributors LMQ, RPD and PN researched data, contributed to discussion, wrote the first draft of the manuscript and reviewed and edited the final manuscript. CB, SB, KB, JG, SG, EH, SK, KK, NL, SM, VRP, TR and GW researched data and reviewed and edited the manuscript. All authors approved the final version of the manuscript. LMQ is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Funding The ELSA study was supported by Diabetes UK and Breakthrough T1D (formerly the Juvenile Diabetes Research Foundation), grant code: 20_0006315. The funders had no role in the study design; no role in data collection, analysis, and interpretation of data; no role in the writing of the protocol paper; and no role in the decision to submit the paper for publication.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants and was approved by Health and Research for Wales, IRAS: 309252. Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request. Not applicable.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Lauren M Quinn <http://orcid.org/0000-0003-0500-4417>

Parth Narendran <http://orcid.org/0000-0002-4583-8793>

REFERENCES

- DiMeglio LA, Evans-Molina C, Oram RA. Type 1 diabetes. *Lancet* 2018;391:2449–62.
- Insel RA, Dunne JL, Atkinson MA, *et al*. Staging presymptomatic type 1 diabetes: a scientific statement of JDRF, the Endocrine Society, and the American Diabetes Association. *Diabetes Care* 2015;38:1964–74.
- Ziegler AG, Rewers M, Simell O, *et al*. Seroconversion to multiple islet autoantibodies and risk of progression to diabetes in children. *JAMA* 2013;309:2473–9.
- American Diabetes Association. 2. Classification and Diagnosis of Diabetes: Standards of Medical Care in Diabetes—2021. *Diabetes Care* 2021;44:S15–33.
- Besser REJ, Bell KJ, Couper JJ, *et al*. ISPAD Clinical Practice Consensus Guidelines 2022: Stages of type 1 diabetes in children and adolescents. *Pediatr Diabetes* 2022;23:1175–87.
- Sims EK, Besser REJ, Dayan C, *et al*. Screening for Type 1 Diabetes in the General Population: A Status Report and Perspective. *Diabetes* 2022;71:610–23.
- Ziegler A-G, Kick K, Bonifacio E, *et al*. Yield of a Public Health Screening of Children for Islet Autoantibodies in Bavaria, Germany. *JAMA* 2020;323:339–51.
- Hummel S, Carl J, Friedl N, *et al*. Children diagnosed with presymptomatic type 1 diabetes through public health screening have milder diabetes at clinical manifestation. *Diabetologia* 2023;66:1633–42.
- Alonso GT, Coakley A, Pyle L, *et al*. Diabetic Ketoacidosis at Diagnosis of Type 1 Diabetes in Colorado Children, 2010–2017. *Diabetes Care* 2020;43:117–21.
- Usher-Smith JA, Thompson MJ, Zhu H, *et al*. The pathway to diagnosis of type 1 diabetes in children: a questionnaire study. *BMJ Open* 2015;5:e006470.
- Townson J, Cannings-John R, Francis N, *et al*. Presentation to primary care during the prodrome of type 1 diabetes in childhood: A case-control study using record data linkage. *Pediatr Diabetes* 2019;20:330–8.
- Cherubini V, Grimsmann JM, Åkesson K, *et al*. Temporal trends in diabetic ketoacidosis at diagnosis of paediatric type 1 diabetes between 2006 and 2016: results from 13 countries in three continents. *Diabetologia* 2020;63:1530–41.
- RSR. 3-screen islet cell antibody ELISA kit. 2020. Available: <https://www.rsrltd.com/pdf%20tech/3%20SCREEN%20ICA%20ELISA.pdf>
- Dunger DB, Bruggaber SFA, Mander AP, *et al*. INNODIA Master Protocol for the evaluation of investigational medicinal products in children, adolescents and adults with newly diagnosed type 1 diabetes. *Trials* 2022;23:414.
- Gov.UK. English indices of deprivation. 2019. Available: <https://www.gov.uk/government/statistics/english-indices-of-deprivation-2019>
- Glaser N, Fritsch M, Priyambada L, *et al*. ISPAD clinical practice consensus guidelines 2022: Diabetic ketoacidosis and hyperglycemic hyperosmolar state. *Pediatr Diabetes* 2022;23:835–56.
- NICE. National institute for health and care excellence (nice). diabetes (type 1 and type 2) in children and young people: diagnosis and management. 2023. Available: <https://www.nice.org.uk/guidance/ng18>
- Hendriks AEJ, Marcovecchio ML, Besser REJ, *et al*. Clinical care advice for monitoring of islet autoantibody positive individuals with presymptomatic type 1 diabetes. *Diabetes Metab Res Rev* 2024;40:e3777.
- Wilson DM, Pietropaolo SL, Acevedo-Calado M, *et al*. CGM Metrics Identify Dysglycemic States in Participants From the TrialNet Pathway to Prevention Study. *Diabetes Care* 2023;46:526–34.
- Steck AK, Larsson HE, Liu X, *et al*. Residual beta-cell function in diabetes children followed and diagnosed in the TEDDY study compared to community controls. *Pediatr Diabetes* 2017;18:794–802.
- Karges B, Prinz N, Placzek K, *et al*. A Comparison of Familial and Sporadic Type 1 Diabetes Among Young Patients. *Diabetes Care* 2021;44:1116–24.
- Winkler C, Schober E, Ziegler AG, *et al*. Markedly reduced rate of diabetic ketoacidosis at onset of type 1 diabetes in relatives screened for islet autoantibodies. *Pediatr Diabetes* 2012;13:308–13.
- Cherubini V, Chiarelli F. Autoantibody test for type 1 diabetes in children: are there reasons to implement a screening program in the general population? A statement endorsed by the Italian Society for Paediatric Endocrinology and Diabetes (SIEDP-ISPED) and the Italian Society of Paediatrics (SIP). *Ital J Pediatr* 2023;49:87.
- Bosi E, Catassi C. Screening type 1 diabetes and celiac disease by law. *Lancet Diabetes Endocrinol* 2024;12:12–4.
- Bonifacio E, Winkler C, Achenbach P, *et al*. Effect of population-wide screening for presymptomatic early-stage type 1 diabetes on paediatric clinical care. *Lancet Diabetes Endocrinol* 2024;12:376–8.
- Elding Larsson H, Vehik K, Gesualdo P, *et al*. Children followed in the TEDDY study are diagnosed with type 1 diabetes at an early stage of disease. *Pediatr Diabetes* 2014;15:118–26.
- Raab J, Haupt F, Scholz M, *et al*. Capillary blood islet autoantibody screening for identifying pre-type 1 diabetes in the general population: design and initial results of the Fr1da study. *BMJ Open* 2016;6:e011144.
- Yazidi M, Mahjoubi S, Oueslati I, *et al*. The remission phase in adolescents and young adults with newly diagnosed type 1 diabetes mellitus: prevalence, predicting factors and glycaemic control during follow-up. *Arch Endocrinol Metab* 2022;66:2359–399700000456:222–8.
- Lundgren M, Jonsdottir B, Elding Larsson H, *et al*. Effect of screening for type 1 diabetes on early metabolic control: the DiPiS study. *Diabetologia* 2019;62:53–7.
- Smith LB, Liu X, Johnson SB, *et al*. Family adjustment to diabetes diagnosis in children: Can participation in a study on type 1 diabetes genetic risk be helpful? *Pediatr Diabetes* 2018;19:1025–33.
- Khanolkar AR, Amin R, Taylor-Robinson D, *et al*. Diabetic Ketoacidosis Severity at Diagnosis and Glycaemic Control in the First Year of Childhood Onset Type 1 Diabetes-A Longitudinal Cohort Study. *Int J Environ Res Public Health* 2017;15:26.
- Mehta S, Ryabets-Lienhard A, Patel N. Pediatric Endocrine Society Statement on Considerations for Use of Teplizumab (Tzield) in Clinical Practice. *Horm Res Paediatr* 2024;2024:1–12.

- 33 Herold KC, Bundy BN, Long SA, *et al.* An Anti-CD3 Antibody, Teplizumab, in Relatives at Risk for Type 1 Diabetes. *N Engl J Med* 2019;381:603–13.
- 34 NICE. Teplizumab for delaying the onset of type 1 diabetes in people 8 years and over at risk of developing the condition TS ID 10753. 2023.
- 35 Anand V, Li Y, Liu B, *et al.* Islet Autoimmunity and HLA Markers of Presymptomatic and Clinical Type 1 Diabetes: Joint Analyses of Prospective Cohort Studies in Finland, Germany, Sweden, and the U.S. *Diabetes Care* 2021;44:dc201836:2269–76.
- 36 Bosi E, Boulware DC, Becker DJ, *et al.* Impact of Age and Antibody Type on Progression From Single to Multiple Autoantibodies in Type 1 Diabetes Relatives. *J Clin Endocrinol Metab* 2017;102:2881–6.
- 37 Beyerlein A, Bonifacio E, Vehik K, *et al.* Progression from islet autoimmunity to clinical type 1 diabetes is influenced by genetic factors: results from the prospective TEDDY study. *J Med Genet* 2019;56:602–5.
- 38 Krischer JP, Liu X, Lernmark Å, *et al.* Predictors of the Initiation of Islet Autoimmunity and Progression to Multiple Autoantibodies and Clinical Diabetes: The TEDDY Study. *Diabetes Care* 2022;45:2271–81.
- 39 Quinn LM, Shukla D, Greenfield SM, *et al.* EarLy Surveillance for Autoimmune diabetes: protocol for a qualitative study of general population and stakeholder perspectives on screening for type 1 diabetes in the UK (ELSA 1). *BMJ Open Diabetes Res Care* 2022;10:e002750.
- 40 Phillip M, Achenbach P, Addala A, *et al.* Consensus Guidance for Monitoring Individuals With Islet Autoantibody-Positive Pre-Stage 3 Type 1 Diabetes. *Diabetes Care* 2024;47:1276–98.
- 41 Elding Larsson H, Vehik K, Bell R, *et al.* Reduced prevalence of diabetic ketoacidosis at diagnosis of type 1 diabetes in young children participating in longitudinal follow-up. *Diabetes Care* 2011;34:2347–52.
- 42 Driscoll KA, Tamura R, Johnson SB, *et al.* Adherence to oral glucose tolerance testing in children in stage 1 of type 1 diabetes: The TEDDY study. *Pediatr Diabetes* 2021;22:360–8.
- 43 Group TS. The Environmental Determinants of Diabetes in the Young (TEDDY) study: study design. *Pediatr Diabetes* 2007;8:286–98.
- 44 Wentworth JM, Oakey H, Craig ME, *et al.* Decreased occurrence of ketoacidosis and preservation of beta cell function in relatives screened and monitored for type 1 diabetes in Australia and New Zealand. *Pediatr Diabetes* 2022;23:1594–601.
- 45 Besser REJ, Griffin KJ. Transitioning to stage 3 type 1 diabetes: when to start insulin. *The Lancet Diabetes & Endocrinology* 2024.
- 46 Maahs DM, Hermann JM, Holman N, *et al.* Rates of diabetic ketoacidosis: international comparison with 49,859 pediatric patients with type 1 diabetes from England, Wales, the U.S., Austria, and Germany. *Diabetes Care* 2015;38:1876–82.
- 47 Usher-Smith JA, Thompson MJ, Sharp SJ, *et al.* Factors associated with the presence of diabetic ketoacidosis at diagnosis of diabetes in children and young adults: a systematic review. *BMJ* 2011;343:d4092.

Supplementary Table 1 – Case summaries

Case	ELSA outcome	Age at ELSA entry (years)	Ethnicity	Gender	ELSA auto-antibodies	ELSA HbA1c % (mmol/mol)	Fasting glucose (mmol/L)	2-hr glucose (mmol/L)	Family History of type 1 diabetes	Co-morbidity	Time to stage 3b onset (symptomatic type 1 diabetes, insulin commenced)	DKA at diagnosis	Hospital admission at stage 3 identification	Details of hospital admission	HbA1c at treatment onset % (mmol/mol)	Treatment regimen
1	Stage 3a	7	White British	Female	IA-2A, ZnT8	6.0 (42)	4.6	12.6	Sibling		8 Months after stage 3a OGTT	No	No		6.6 (49)	Commenced 1 unit insulin detemir and then added 1 unit insulin aspart with evening meal added
2	Stage 3a	9	White British	Female	IAA, GAD	6.0 (42)	5.3	12.9	No		Remained asymptomatic, insulin commenced 4 months after stage 3a OGTT	No	No		6.0 (42)	Started 0.5 units insulin aspart with breakfast and split dose 0.5 and 0.5units pre and post evening meal
3	Stage 3b	9	White British	Male	IA-2A, ZnT8	6.3 (45)	4.3	12.1	Parent	Vestibular/abdominal migraines	Immediately	No	Yes, 1 night	Admitted for education	6.4 (46)	Commenced insulin immediately. Started insulin 19/09/23 at 0.3U/kg/day = insulin detemir 4 units at 2000 (detemir reduced to 3 units on 20/09/23), insulin aspart 2 units with meals three times daily (19/09/23)
4	Stage 3b	3	White British	Male	GAD, ZnT8	5.7 (39)	4.7	13.6	Sibling		Immediately, 10 days of thirst and hyperphagia prior to ELSA OGTT	No	Yes, 3 nights	Admitted for glucose monitoring	5.7 (39)	First night of admission, commenced on 1 unit insulin detemir leading to hypoglycaemia. Admitted for 3 nights for glucose

																monitoring. On discharge, advised insulin aspart 0.5-1 unit with evening meal (1:20). Mum also giving correction doses.
5	Stage 2	5	White British	Female	IA-2A, GAD, ZnT8	5.9 (41)	4.1	9.9	Parent	Constipation	3 months after stage 2 OGTT	No	No (2 paediatric assessment unit attendances pre-stage 3)		6.7 (50)	Commenced 0.5 units ultra-rapid acting insulin aspart with evening meal and increased to 0.5 units ultra-rapid acting insulin aspart with breakfast and 1 unit with evening meal
6	Stage 1	3	White British	Male	IA-2A, GAD, ZnT8	5.6 (38)	4.7	7	Sibling		4 months after stage 1 OGTT	No	Yes, 1 night			Now on omnipod pump with automated insulin delivery
7	Single Aab	8	White British	Male	IA-2A				Grandparent		1 month after completing ELSA	No	Yes		6.7 (50)	Commenced insulin degludec 6 units (initially reduced down to 2.5 units and then to 4.5 units) and insulin aspart 1:30 (reduced to 1:35)
8	Single Aab	3	Asian or British Pakistani	Male	GAD				Sibling		1-2 months after completing ELSA	No	No			
9	Multiple Aab	5	White British	Male	IA-2A, GAD, ZnT8				Parent and sibling		Whilst awaiting OGTT	No	Yes		6.6 (49)	
10	DBS positive	8	White British	Male					Sibling			No	Yes		5.5 (37)	Commenced 1 unit insulin glargine and 0.5 units fixed dose insulin aspart

11	DBS positive	9	White British	Male					Cousin	Juvenile Idiopathic Arthritis (on Methotrexate), Psoriasis, Vitiligo	4 months pre diagnosis	No	Yes, 3 nights	10.6 (92)	with meals. Commenced insulin detemir (BD) 3.5units and 2.5 units and insulin aspart 1:25-35, changed to ultra-rapid acting insulin aspart 1:25-35
12	DBS positive	7	White British	Female					Sibling		2 weeks pre-diagnosis	No	Yes, 3 nights	15.1 (142)	Commenced insulin degludec and insulin aspart
13	DBS positive	11	White British	Male					Parent	Coeliac disease	4 weeks pre-diagnosis	Yes, mild	Yes, 1 night	6.5 (47)	Commenced insulin aspart and insulin detemir and changed to ultra-fast acting insulin aspart, but experienced frequent hypoglycaemia. Commenced automated insulin delivery T-Slim within 4 months of diagnosis
14	DBS positive	12	Black or British Black African	Male					No		2 months pre-diagnosis/prior to ELSA venous confirmation	Yes, mild	Yes, 7 nights	12.9 (118)	Commenced insulin glargine 16 units at night and insulin aspart 1:10-20 with meals (reduced insulin doses)

Key: DBS – dried blood spot, Aab – autoantibody

