Supplemental Tables

Supp. Table 1: CKD Severity and Stage Definition

Severity	Stage	Evidence of Albuminuria	eGFR (ml/min/1.73m^2)
No CKD		Evidence of UACR levels <30 mg/g	≥90
Mild to Moderate	1	Evidence of UACR levels ≥30 mg/g	≥90
	2	Evidence of UACR levels ≥30 mg/g	60-89
	3a	N/A	45-59
Moderate to kidney failure	3b	N/A	30-44
	4	N/A	15-29
	5	N/A	<15 or dialysis

UACR: urinary albumin creatinine ratio; eGFR: estimated glomerular filtration rate ; N/A: not applicable

For Review Only

Supp. Table 2: Prevalence of CKD in T2D by Age and Gender

eGFR determined by CKD-EPI	Prevalence of CKD by Age				Prevalence of CKD by Gender			
	Age 18-64 Years		Age ≥65 Years		Male		Female	
	N	% (95% CI)	N	% (95% CI)	N	% (95% CI)	N	% (95% CI)
No CKD	725	74.3 (71.4-77.3)	397	41.3 (37.3-45.4)	551	60.0 (55.9-64.1)	571	61.3 (56.9-65.8)
CKD all stages	351	25.7 (22.7-28.6)	533	58.7 (54.6-62.7)	464	40.0 (35.9-44.1)	420	38.7 (34.2-43.1)
Stage 1 (eGFR ≥90 and UACR ≥30 mg/g)	170	12.4 (10.2-14.7)	31	2.8 (1.8-3.8)	116	10.5 (8.7-12.2)	85	6.5 (4.7-8.2)
Stage 2 (eGFR 60-89 and UACR ≥30 mg/g)	87	6.4 (4.6-8.1)	134	15.1 (12.0-18.2)	126	11.3 (8.7-13.9)	95	8.7 (6.3-11.1)
Stage 3a (eGFR 45-59)	48	4.2 (2.8-5.7)	209	22.7 (19.4-26.0)	136	12.1 (9.0-15.1)	121	11.7 (9.0-14.5)
Stage 3b (eGFR 30-44)	22	1.2 (0.7-1.8)	111	12.4 (9.6-15.1)	60	4.5 (3.2-5.8)	73	7.2 (5.2-9.2)
Stage 4 (eGFR 15-29)	9	0.6 (0.1-1.1)	39	5.2 (3.5-6.8)	14	1.2 (0.4-2.0)	34	3.8 (2.2-5.4)
Stage 5 (eGFR < 15)	15	0.7 (0.3-1.2)	9	0.5 (0.1-0.9)	12	0.5 (0.2-0.9)	12	0.8 (0.3-1.3)
Total	**1,076**	**100**	**930**	**100**	**1015**	**100**	**991**	**100**

CKD: chronic kidney disease; CKD-EPI: The Chronic Kidney Disease Epidemiology Collaboration equation; eGFR: estimated glomerular filtration rate (ml/min/1.72m^2); T2D: type 2 diabetes mellitus; UACR: urinary albumin creatinine ratio (mg/g).

Confi... ...iew Only