Decreased arterial distensibility and postmeal hyperinsulinemia in young Japanese women with family history of diabetes

Mika Takeuchi,¹ Bin Wu,^{2,3} Mari Honda,^{2,4} Ayaka Tsuboi,^{1,5} Kaori Kitaoka,^{1,6} Satomi Minato,^{1,7} Miki Kurata,^{1,8} Tsutomu Kazumi ^{1,2,9}, Keisuke Fukuo^{1,2,8}

To cite: Takeuchi M, Wu B, Honda M, *et al*. Decreased arterial distensibility and postmeal hyperinsulinemia in young Japanese women with family history of diabetes. *BMJ Open Diab Res Care* 2020;**8**:e001244. doi:10.1136/bmjdr-2020-001244

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/bmjdr-2020-001244>).

Received 4 February 2020
Revised 9 April 2020
Accepted 14 April 2020

ABSTRACT

Introduction To assess vascular function and characterize insulin secretion using a physiological approach in Japanese women with family history of type 2 diabetes (FHD).

Research design and methods Standardized mixed-meal tests were performed with multiple postprandial glucose, insulin and free fatty acids (FFA) measurements over a 30–120 min period in 31 Japanese women aged 21–24 years. Arterial distensibility was assessed as well.

Results Fasting glucose, triglyceride and insulin averaged <90 mg/dL, <60 mg/dL and <5 µU/mL, respectively, and did not differ cross-sectionally between 10 with (FHD+) and 21 without FHD (FHD–). FHD+ showed higher insulin responses not only during the first 30 min ($p=0.005$) but also during the second hour (60–120 min, $p<0.05$) in spite of identical postprandial suppression of FFA and identical fasting and postprandial glucose and FFA concentrations, except for higher 60 min FFA in FHD+. Further, FHD+ had decreased arterial distensibility ($p=0.003$). On multivariate regression analysis, arterial distensibility emerged as the only significant independent predictor of FHD+. Endurance training in FHD+ did not alter decreased arterial distensibility whereas it abolished postprandial hyperinsulinemia.

Conclusions FHD was associated with decreased arterial distensibility and postprandial hyperinsulinemia despite nearly identical postprandial glycemia and postprandial FFA suppression, suggesting that impaired vascular insulin sensitivity may precede glucose and lipid dysmetabolism in normal weight Japanese women aged 22 years.

INTRODUCTION

Both family history of type 2 diabetes (FHD) and obesity are two major independent risk factors for diabetes. Insulin resistance and β -cell failure are hallmarks of type 2 diabetes. Zhu *et al*¹ have recently demonstrated that among 4.9 million adults in the USA from 2012 to 2013, age-standardized prevalence estimates of both diabetes and pre-diabetes were higher among racial/ethnic minority groups including Asians than among whites, especially at lower body mass index (BMI)

Significance of this study

What is already known about this subject?

► Positive family history of type 2 diabetes (FHD) in Asians has been reported to be related to impaired insulin secretion to glucose and less muscle insulin resistance.

What are the new findings?

► Normoglycemic, normolipidemic Japanese women with positive FHD had postprandial hyperinsulinemia and decreased arterial distensibility, suggesting that impaired vascular insulin sensitivity may precede glucose and lipid dysmetabolism.

How might these results change the focus of research or clinical practice?

► The results may suggest clinical significance of standardized meal tests, a physiological approach, and highlight the importance of vascular insulin sensitivity/resistance in the context of FHD, which has been less studied.

values. Normal weight (BMI <25 kg/m²) type 2 diabetes is highly prevalent in Asia and Africa.² We have previously demonstrated that more than half of Japanese patients with type 2 diabetes are considered normal weight^{3,4} and that more than half of normal weight diabetics had FHD.⁴ These observations suggest that poor insulin secretion rather than obesity-driven insulin resistance may be at play in the natural history of diabetes in these populations. Indeed, decreased first-phase insulin secretion to glucose has been reported in young normal weight Japanese offspring of type 2 diabetics.^{5,6} Further, impaired β -cell function may be present earlier than visceral adiposity in some Japanese Americans who subsequently develop type 2 diabetes although both insulin resistance and impaired β -cell function contribute to the development of type 2 diabetes.⁷ Use of

© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Tsutomu Kazumi;
kazumi@mukogawa-u.ac.jp

the intravenous glucose tolerance test or hyperglycemic clamp may provide a more accurate assessment of insulin secretion compared with oral glucose tolerance test-derived measures.^{8,9} However, exclusion of the gastrointestinal tract limits the physiologic understanding of in vivo β -cell function. Although an oral glucose tolerance test is a standardized tool to evaluate glucose tolerance, meal tests provide us information on daily-life conditions. However, studies on meal testing in people with FHD are limited in general, in Asians in particular, as discussed later.

FHD also was proposed as a significant cardiovascular risk factor.¹⁰ Previous studies have found an association between FHD and cardiovascular disease as measured by arterial stiffness.^{11–13} Again, studies on arterial properties in people with FHD are limited in Asian populations. Therefore, we investigated glucose and lipid responses to a standardized breakfast and arterial distensibility in young Japanese women with FHD. Because our young women comprised collegiate female athletes and untrained women,^{14–17} we investigated whether chronic exercise training may affect results in women with FHD.

PATIENTS AND METHODS

Among 35 previously reported women (17 collegiate athletes and 18 untrained students),^{14–16} 31 women aged 21–24 years (16 collegiate athletes and 15 untrained students) provided data on family history of diabetes, which was considered positive if they reported that a parent or a grandparent was on anti-diabetic drugs. Unfortunately, information was not available on the extent of family history (ie, how many family members have the condition) and the nature of the family history (paternal or maternal). Ten women with FHD (FHD+) and 21 without FHD (FHD–) were compared cross-sectionally. The athletes were students of the Department of Health and Sports Sciences and had been training regularly for 2 years or longer prior to the study, 5 hours a day, 6 days a week. Non-athletes were students of the Department of Food Sciences and Nutrition and were not engaged in any regular sport activity. Subjects who reported that they were in treatment for acute or chronic inflammatory diseases, endocrine, cardiovascular, hepatic, renal diseases, hormonal contraception, unusual dietary habits were excluded. Nobody reported receiving any medications or having regular supplements. All subjects gave written consent after the experimental procedure had been explained.

At 8:30 after a 12-hours overnight fast, participants completed a standardized meal as previously reported.^{15,16} This meal was considered as a breakfast meal (total energy 450 kcal) and provided 33.3% of calories from fat (16.7g), 51.4% from carbohydrates (57.8g), and 15.3% from protein (17.2g). Participants were asked to consume the entire meal within 15 min. Venous blood was drawn at baseline (0 min), 30, 60, and 120 min after the start of the meal for the measurement of plasma

glucose (PG), serum insulin, triglyceride (TG) and free fatty acids (FFA) concentrations. PG was determined by the hexokinase/glucose-6-phosphate dehydrogenase method (inter-assay coefficient of variation (CV) <2%). Serum insulin was measured by an ELISA method with a narrow specificity excluding des-31, des-32, and intact proinsulin (inter-assay CV <6%). FFA was measured using enzymatic colorimetric method (Wako Chemicals, Tokyo, Japan). In fasted blood samples, the following were measured as previously reported¹⁷; serum cholesterol, TG, high-density lipoprotein (HDL) cholesterol, adiponectin, leptin and high-sensitivity C-reactive protein (hsCRP).

Early phase insulin responses were calculated as the increment in serum insulin during the first 30 min of the meal test (Δ insulin₃₀). The area under the concentration curve (AUC) of PG, serum insulin, TG and FFA was calculated with the trapezoidal method (glucose-AUC, insulin-AUC, TG-AUC, FFA-AUC, respectively). FFA suppression during the meal test was assessed by percentage changes in FFA (%FFA) calculated as FFA concentrations at 30, 60, and 120 min divided by the fasting FFA concentration in each participant. Adipose tissue insulin resistance (AT-IR) was calculated as a product of fasting insulin and FFA concentrations.¹⁸ Homeostasis model assessment-insulin resistance (HOMA-IR) was calculated as previously reported.¹⁹

Lean mass, fat mass, and bone mineral mass for arms, legs, trunk and the total body were quantified by whole-body dual-energy X-ray absorptiometry (DXA) (Hologic QDR-2000, software V.7.20D, Waltham, Massachusetts, USA) as previously reported.¹⁷ The leg region included the entire hip, thigh and leg. General adiposity was assessed using BMI and fat mass index (FMI), which was calculated as body fat mass in kilograms divided by height in meter squared. Abdominal fat accumulation was assessed by waist circumference and the ratio of trunk fat to leg fat.²⁰ Skeletal muscle mass was assessed by skeletal muscle mass index (SMI) calculated as lean mass in arms and legs in kilograms divided by height in meter squared.

Cardio-ankle vascular index (CAVI) was measured by VaSera device (VS-1000, Fukuda Denshi, Tokyo, Japan).²¹ CAVI is a new measure of arterial distensibility that reflects the distensibility from the ascending aorta to the ankle arteries and demonstrates little dependence on blood pressure during the evaluation. It has recently been validated to be a reliable screening tool for arterial distensibility.^{22,23}

Data were presented as mean $\pm$ SD. Due to deviation from normal distribution, HOMA-IR and hsCRP were logarithmically transformed for analyses. Differences between two groups were compared by t-test. Multivariate logistic regression analyses were done for FHD+ as a dependent variable. Independent variables included were those that displayed significant differences between FHD+ and FHD– women. A two-tailed $p < 0.05$ was considered statistically significant. All calculations were performed with SPSS system V.15.0.

Table 1 Anthropometric and cardiometabolic features of young Japanese women in the presence and absence of family history of diabetes

	Family history of diabetes						
	Present			Absent			P values
	n=10			n=21			
Age (years)	22.7	±	0.9	22.0	±	0.3	0.038
Body mass index (kg/m ²)	21.6	±	2.8	21.9	±	2.3	0.727
Fat mass index (kg/m ²)	5.8	±	2.5	5.8	±	1.8	0.996
Waist circumference (cm)	71.8	±	7.6	74.3	±	5.5	0.303
SMI (kg/m ²)	6.3	±	0.4	6.6	±	0.7	0.079
Trunk/leg fat ratio	1.24	±	0.23	1.19	±	0.25	0.595
HOMA-IR	0.9	±	0.5	0.7	±	0.4	0.178
AT-IR	2.1	±	1.2	1.5	±	0.8	0.159
HbA1c (%)	5.2	±	0.2	5.2	±	0.2	0.336
Fasting glucose (mg/dL)	86	±	6	83	±	4	0.159
30 min glucose (mg/dL)	107	±	15	106	±	15	0.847
60 min glucose (mg/dL)	84	±	15	85	±	20	0.852
120 min glucose (mg/dL)	72	±	12	77	±	12	0.308
Fasting insulin (μU/mL)	4.3	±	2.2	3.3	±	1.9	0.212
30 min insulin (μU/mL)	47	±	21	28	±	12	0.005
60 min insulin (μU/mL)	38	±	20	23	±	14	0.022
120 min insulin (μU/mL)	19	±	9	10	±	6	0.002
Δglucose30 (mg/dL)	21	±	16	23	±	15	0.813
Δinsulin30 (μU/mL)	43	±	20	25	±	12	0.005
Glucose-AUC (mg/dL/2 hour)	174	±	18	176	±	23	0.795
Insulin-AUC (μU/mL/2 hour)	63	±	28	38	±	17	0.004
Fasting FFA (mEq/L)	0.49	±	0.16	0.48	±	0.17	0.838
30 min FFA (mEq/L)	0.37	±	0.12	0.35	±	0.12	0.602
60 min FFA (mEq/L)	0.26	±	0.07	0.21	±	0.05	0.042
120 min FFA (mEq/L)	0.23	±	0.05	0.24	±	0.06	0.636
30 min %FFA (%)	78.0	±	27.6	75.5	±	21.6	0.779
60 min %FFA (%)	55.8	±	18.3	48.9	±	18.3	0.333
120 min %FFA (%)	51.0	±	23.1	56.3	±	26.2	0.590
Fasting TG (mg/dL)	58	±	23	51	±	17	0.283
30 min TG (mg/dL)	67	±	24	54	±	19	0.119
60 min TG (mg/dL)	79	±	23	66	±	23	0.185
120 min TG (mg/dL)	87	±	33	75	±	30	0.339
FFA-AUC (mEq/L/2 hour)	0.62	±	0.13	0.57	±	0.13	0.366
TG-AUC (mg/dL/2 hour)	151	±	49	127	±	44	0.197
HDL cholesterol (mg/dL)	72	±	13	75	±	12	0.546
Serum cholesterol (mg/dL)	179	±	26	176	±	22	0.750
Leptin (ng/mL)	8.8	±	6.1	6.4	±	3.4	0.157
Adiponectin (mg/L)	9.5	±	2.9	12.4	±	5.0	0.101
hsCRP (μg/dL)	31	±	52	45	±	109	0.715
Systolic BP (mm Hg)	112	±	8	114	±	7	0.456
Diastolic BP (mm Hg)	70	±	8	67	±	5	0.451
Resting pulse rate (bpm)	54	±	8	52	±	11	0.593
Athletes (n,%)	4	,	40.0	12	,	57.1	0.458

Mean±SD or n, %.

AT, adipose tissue; AUC, area under the concentration curve; BP, blood pressure; bpm, beats per minute; FFA, free fatty acids; %FFA, percent changes in FFA; Δglucose30, the increment in plasma glucose during the first 30 min; HDL, high-density lipoprotein; HOMA, homeostasis model assessment; hsCRP, high-sensitivity C-reactive protein; Δinsulin30, the increment in serum insulin during the first 30 min of the meal test; IR, insulin resistance; SMI, skeletal muscle mass index; TG, triglycerides.

Figure 1 Cardio-ankle vascular index (CAVI) scores in each Japanese woman in the presence (FHD+, circles) and absence (FHD-, squares) of family history of type 2 diabetes (FHD). Open and closed symbols denote athletes and non-athletes, respectively. A circle and a square with vertical bars; mean $\pm$ SD in the two groups. * $p=0.003$ versus FHD-.

RESULTS

As previously reported^{15–17} and shown in table 1, BMI averaged $<22\text{ kg/m}^2$, waist circumference $<75\text{ cm}$, HOMA-IR <1.0 and fasting TG $<60\text{ mg/dL}$, suggesting that young women were insulin sensitive and these surrogate markers of insulin resistance did not differ between 10 FHD+ and 21 FHD- (table 1).

Although there was no difference in the percentage of athletes between FHD+ and FHD-, SMI tended to be lower in FHD+ (table 1). FMI, trunk/leg fat ratio, serum adiponectin, leptin and hsCRP did not differ between two groups. There was also no difference in fasting glucose, HbA1c, AT-IR, HDL cholesterol, postprandial TG and TG-AUC. Although systolic and diastolic blood pressure did not differ, CAVI score was higher in FHD+ (figure 1).

Despite nearly identical postprandial glycemia and glucose-AUC, postprandial insulinemia and insulin-AUC were greater in FHD+ (table 1 and figure 2). In addition, FHD+ had greater $\Delta\text{insulin}_{30}$ despite nearly identical incremental glucose concentrations during the first 30 min. Further, postprandial suppression of FFA as assessed by %FFA and FFA concentrations did not differ

Figure 2 Mean ($\pm$ SD) plasma glucose, insulin and free fatty acids (FFA) concentration and percent changes in FFA (%FFA) in the presence (red lines) and absence (green lines) of family history of type 2 diabetes. * $p<0.05$; ** $p<0.01$.

between FHD+ and FHD- although 60 min FFA was higher in FHD+.

Results were essentially the same in the same analyses described above, in which four women who did not provide data on FHD were included as negative FHD (online supplementary table 1). We believe that it seems reasonable to assume that women who did not provide data on FHD might not have FHD.

Multivariate logistic regression analyses were done for FHD+ as a dependent variable. Independent variables included were $\Delta\text{insulin}_{30}$, insulin-AUC and arterial distensibility. Arterial distensibility emerged as the only significant independent predictor of FHD+ (OR; 12.0, 95% CI; 1.6 to 91.0, $p=0.016$).

FHD+ athletes compared with FHD+ non-athletes had higher SMI and lower resting pulse rate, the latter of which averaged $48\pm6\text{ bpm}$ (table 2). As shown in figure 3, postprandial insulinemia was increased in FHD+ non-athletes whereas postprandial insulinemia in FHD+ athletes was lower compared with FHD+ non-athletes but was comparable to FHD- women. Postprandial glycemia and postprandial suppression of FFA did not differ between FH+ athletes and FH+ non-athletes. However, the two FHD+ groups did not differ in CAVI score which was increased compared with FHD- (table 2 and figure 1).

DISCUSSION

In normal weight Japanese women aged 23 years who were not insulin resistant rather insulin sensitive, meal-induced insulin responses were increased rather than decreased not only during the first 30 min but also during the second hour (60–120 min) in FHD+ compared with FHD- in the presence of nearly identical postprandial glycemia and FFA suppression. Moreover, CAVI score was higher in FHD+, suggesting decreased arterial distensibility. On multivariate regression analysis, arterial distensibility emerged as the only significant independent predictor of FHD+. Furthermore, endurance training in FHD+ abolished increased postprandial insulin responses whereas it did not alter decreased arterial distensibility.

Type 2 diabetes in East Asians is well known to be characterized primarily by β -cell dysfunction rather than obesity-driven insulin resistance.²⁴ Japanese offspring of type 2 diabetics whose age and BMI were comparable to those of our young women showed decreased first-phase insulin secretion to intravenous glucose.^{5 6} Further, impaired β -cell function may be present earlier than visceral adiposity in some Japanese Americans who subsequently develop type 2 diabetes.⁷ In contrast, in the present study, young normal weight Japanese FHD+ women showed increased early-phase insulin responses after meal ingestion despite nearly identical incremental glucose concentrations, suggesting increased incretin effect. Nyholm *et al*²⁵ examined profiles of insulin in glucose tolerant but insulin resistant relatives of type 2 diabetics during a 24-hours period including three meals. They found that incremental insulin secretion after each

Table 2 Comparisons between female collegiate athletes and untrained students (non-athletes) with family history of diabetes

	Athletes			Non-athletes			P values
	n=4			n=6			
Age (years)	21.8	±	0.1	23.3	±	0.6	0.001
Body mass index (kg/m²)	20.9	±	0.7	22.0	±	3.6	0.490
Fat mass index (kg/m²)	4.4	±	1.0	6.7	±	2.8	0.162
Waist circumference (cm)	69.2	±	3.1	73.6	±	9.4	0.318
SMI (kg/m²)	6.6	±	0.3	6.0	±	0.3	0.018
Trunk/leg fat ratio	1.14	±	0.16	1.30	±	0.26	0.299
Fasting glucose (mg/dL)	88	±	5	84	±	6	0.317
30 min glucose (mg/dL)	114	±	15	103	±	14	0.264
60 min glucose (mg/dL)	84	±	10	83	±	19	0.951
120 min glucose (mg/dL)	81	±	13	66	±	6	0.083
Fasting insulin (μU/mL)	3.0	±	0.6	5.2	±	2.5	0.144
30 min insulin (μU/mL)	31	±	14	57	±	19	0.044
60 min insulin (μU/mL)	23	±	10	49	±	18	0.031
120 min insulin (μU/mL)	14	±	11	22	±	6	0.264
Fasting FFA (mEq/L)	0.46	±	0.13	0.51	±	0.18	0.627
30 min FFA (mEq/L)	0.37	±	0.17	0.37	±	0.10	0.962
60 min FFA (mEq/L)	0.24	±	0.08	0.27	±	0.06	0.575
120 min FFA (mEq/L)	0.21	±	0.03	0.24	±	0.06	0.364
30 min %FFA (%)	83	±	41	75	±	19	0.692
60 min %FFA (%)	55	±	22	56	±	18	0.912
120 min %FFA (%)	46	±	7	54	±	30	0.638
Δglucose30 (mg/dL)	26	±	14	18	±	18	0.509
Δinsulin30 (μU/mL)	28	±	14	52	±	17	0.045
Glucose-AUC (mg/dL/2 hour)	183	±	14	168	±	19	0.223
Insulin-AUC (μU/mL/2 hour)	40	±	20	78	±	23	0.028
FFA-AUC (mEq/L/2 hour)	0.59	±	0.17	0.63	±	0.10	0.591
Pulse rate (bpm)	48	±	6	58	±	6	0.038
CAVI	6.5	±	0.6	6.1	±	0.2	0.142

Mean±SD.

AUC, area under the concentration curve; bpm, beats per minute; CAVI, cardio-ankle vascular index; %FFA, percent changes in FFA; FFA, free fatty acid; Δglucose30, the increment in plasma glucose during the first 30 min; Δinsulin30, the increment in serum insulin during the first 30 min of the meal test; SMI, skeletal muscle mass index.

meal appeared normal in relatives. We have no explanation for higher meal-induced early phase insulin secretion in Japanese FHD+ women in the present study.

Insulin is a vascular hormone and regulates vascular tone and reactivity.^{26 27} In the postprandial state, increased circulating insulin can enhance blood flow to the microvasculature in muscle thus increasing the access of glucose and insulin to the myocytes. And then, insulin binds to receptors on the skeletal muscle cell surface to enhance glucose uptake. In healthy humans, physiological doses of insulin and mixed meals each increased microvascular perfusion volume.^{28 29} Therefore, it seems reasonable to assume that postprandial hyperinsulinemia in the presence of nearly identical postprandial glycemia may be related to the impaired vasodilatory action of insulin to deliver insulin to skeletal muscle. In the same

way, postprandial hyperinsulinemia in the presence of nearly identical postprandial FFA suppression may be related to the impaired vasodilatory action of insulin in adipose tissue. Fasting glucose averaged <90 mg/dL and TG <60 mg/dL in our FHD+ women. These observations suggest that impaired microvascular insulin sensitivity may precede glucose and lipid dysmetabolism in young FHD+ Japanese women. Experimental studies in diet-induced obesity have shown that vascular insulin resistance (impaired insulin-induced phosphorylation of endothelial nitric oxide synthase) and endothelial dysfunction (reduced nitric oxide production) preceded the onset of insulin resistance in muscle, liver, and fat, suggesting a heightened susceptibility of the vasculature to the detrimental effects of nutrient excess relative to tissues involved in glucose homeostasis and fuel

Figure 3 Mean ($\pm$ SD) plasma glucose, insulin and free fatty acids (FFA) concentration and percent changes in FFA (%FFA) in female athletes (red lines) and non-athletes (green lines) with family history of type 2 diabetes. * $p < 0.05$; ** $p < 0.01$.

storage.³⁰ Postprandial hyperinsulinemia was not found in FHD+ athletes, suggesting that endurance training improved impaired insulin sensitivity in FHD+ offspring as previously reported.^{31–33} Exercise has been reported to increase human skeletal muscle insulin sensitivity via coordinated increases in microvascular perfusion and molecular signaling.³⁴

A standardized meal test gives a more clinically relevant metabolic status whereas the primary function of glucose tolerance tests is to amplify any glucose intolerance, thus aiding diagnosis.³⁵ However, studies using a mixed meal to represent normal physiological feeding in people with FHD+ were limited.^{25 36 37} In agreement with our observations, postprandial hyperinsulinemia in the presence of comparable/slightly raised glycemia and comparable FFA have been found in non-obese relatives in mid-to-late thirties.^{25 36} However, Allerton *et al*³⁷ have shown that responses to a mixed meal, which was similar to that used in the present study, of glucose, insulin and FFA assessed by respective AUC did not differ between normal-weight offspring of mothers with type 2 diabetes aged 21 years and controls.

CAVI is a new measure of arterial distensibility that reflects both the elastic (the ascending aorta) and muscular (the leg artery) artery stiffness and has recently been validated to be a reliable screening tool for arterial distensibility.^{22 23} CAVI score was higher in FHD+ compared with FHD–, suggesting decreased arterial distensibility in FHD+ as previously reported.^{11–13} It is noted that an independent association between decreased arterial distensibility and FHD+ was observed in young normal weight Japanese FHD+ women. Furthermore, endurance training did not improve decreased arterial distensibility in our FHD+ women whereas it improved impaired microvascular insulin sensitivity as described above. Arterial distensibility was not associated with insulin sensitivity in FHD+ subjects whose age and BMI were comparable to those of our participants¹² but was associated in FHD+ subjects who tended to be obese and older.^{10 13}

This study has several strengths, including a homogeneous study population with scarce confounding factors, and accurate and reliable measures of body composition by DXA. The main limitation of our study is small sample size. The cross-sectional design of the present study complicates the drawing of causal inferences, and a single measurement of biochemical variables may be susceptible to short-term variation, which would bias the results toward the null. We used several surrogate markers in the present study, which may be less accurate. As we studied young Japanese women only, results may not be generalized to other gender, age populations, races or ethnicities. Studies with more patients are needed.

CONCLUSIONS

Young insulin sensitive Japanese women with FHD may have decreased arterial distensibility and impaired microvascular insulin sensitivity in skeletal muscle and adipose tissue, suggesting that impaired vascular insulin sensitivity may precede glucose and lipid dysmetabolism. Although female athletes with FHD did not show the metabolic changes, they had decreased arterial distensibility. These observations may suggest a genetic effect on early arterial physiology that may be relevant to later clinical disease.

Author affiliations

¹Research Institute for Nutrition Sciences, Mukogawa Women's University, Nishinomiya, Japan

²Open Research Center for Studying of Lifestyle-Related Diseases, Mukogawa Women's University, Nishinomiya, Japan

³Department of Endocrinology, First Affiliated Hospital of Kunming Medical University, Kunming, China

⁴Department of Health, Sports, and Nutrition, Faculty of Health and Welfare, Kobe Women's University, Kobe, Japan

⁵Department of Nutrition, Osaka City Juso Hospital, Osaka, Japan

⁶Department of Public Health, Shiga University of Medical Science, Otsu, Japan

⁷Graduate School of Human Science and Environment, University of Hyogo, Himeji, Japan

⁸Department of Food Sciences and Nutrition, Mukogawa Women's University, Nishinomiya, Japan

⁹Department of Medicine, Kohnan Kakogawa Hospital, Kakogawa, Japan

Acknowledgements We are indebted to all the participants for their dedicated and conscientious collaboration.

Contributors MT, BW, MH, AT, KK, SM and MK have made substantial contributions to acquisition, analysis and interpretation of data. KF has been involved in drafting the manuscript. TK has been involved in revising it critically for important intellectual content; has given final approval of the version to be published; and agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval The study was approved by the Ethics Committees of the University (No. 07-28) to be in accordance with the Helsinki declaration.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as supplementary information. The Ethics Committee

of Mukogawa Women's University does not allow us to open data except for a manuscript.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Tsutomu Kazumi <http://orcid.org/0000-0002-8855-9268>

REFERENCES

- Zhu Y, Sidell MA, Arterburn D, *et al*. Racial/Ethnic disparities in the prevalence of diabetes and prediabetes by BMI: patient outcomes research to advance learning (portal) multisite cohort of adults in the U.S. *Diabetes Care* 2019;42:2211–9.
- Gujral UP, Weber MB, Staimez LR, *et al*. Diabetes among non-overweight individuals: an emerging public health challenge. *Curr Diab Rep* 2018;18:60.
- Toyasaki M, Honda M, Nanbu S, *et al*. Weight changes in patients with type 2 diabetes since the age of 18 years. *Jap J Clin Nutr* 2007;28:312–8.
- Tanaka S, Honda M, Wu B, *et al*. Clinical features of normal weight Japanese patients with type 2 diabetes who had formerly been obese. *J Atheroscler Thromb* 2011;18:115–21.
- Doi K, Taniguchi A, Nakai Y, *et al*. Decreased glucose effectiveness but not insulin resistance in glucose-tolerant offspring of Japanese non-insulin-dependent diabetic patients: a minimal-model analysis. *Metabolism* 1997;46:880–3.
- Matsumoto K, Sakamaki H, Izumino K, *et al*. Increased insulin sensitivity and decreased insulin secretion in offspring of insulin-sensitive type 2 diabetic patients. *Metabolism* 2000;49:1219–23.
- Chen KW, Boyko EJ, Bergstrom RW, *et al*. Earlier appearance of impaired insulin secretion than of visceral adiposity in the pathogenesis of NIDDM. 5-year follow-up of initially nondiabetic Japanese-American men. *Diabetes Care* 1995;18:747–53.
- Bergman RN, Phillips LS, Cobelli C. Physiologic evaluation of factors controlling glucose tolerance in man: measurement of insulin sensitivity and beta-cell glucose sensitivity from the response to intravenous glucose. *J Clin Invest* 1981;68:1456–67.
- Kahn SE, Prigeon RL, McCulloch DK, *et al*. Quantification of the relationship between insulin sensitivity and beta-cell function in human subjects. Evidence for a hyperbolic function. *Diabetes* 1993;42:1663–72.
- Scheuner MT, Setodji CM, Pankow JS, *et al*. Relation of familial patterns of coronary heart disease, stroke, and diabetes to subclinical atherosclerosis: the multi-ethnic study of atherosclerosis. *Genet Med* 2008;10:879–87.
- Scuteri A, Tesaro M, Rizza S, *et al*. Endothelial function and arterial stiffness in normotensive normoglycemic first-degree relatives of diabetic patients are independent of the metabolic syndrome. *Nutr Metab Cardiovasc Dis* 2008;18:349–56.
- Hopkins KD, Lehmann ED, Jones RL, *et al*. A family history of NIDDM is associated with decreased aortic distensibility in normal healthy young adult subjects. *Diabetes Care* 1996;19:501–3.
- Ikonomidis I, Lambadiari V, Pavlidis G, *et al*. Insulin resistance and acute glucose changes determine arterial elastic properties and coronary flow reserve in dysglycaemic and first-degree relatives of diabetic patients. *Atherosclerosis* 2015;241:455–62.
- Kitaoka K, Takeuchi M, Tsuboi A, *et al*. Increased adipose and muscle insulin sensitivity without changes in serum adiponectin in young female collegiate athletes. *Metab Syndr Relat Disord* 2017;15:246–51.
- Tsuboi A, Takeuchi M, Kitaoka K, *et al*. Post-Prandial plasma glucose less than or equal to 70 mg/dl is not uncommon in young Japanese women. *J Clin Med Res* 2017;9:680–6.
- Takeuchi M, Tsuboi A, Kurata M, *et al*. Associations of postprandial lipemia with trunk/leg fat ratio in young normal weight women independently of fat mass and insulin resistance. *Asia Pac J Clin Nutr* 2018;27:293–9.
- Tanaka S, Wu B, Honda M, *et al*. Associations of lower-body fat mass with favorable profile of lipoproteins and adipokines in healthy, slim women in early adulthood. *J Atheroscler Thromb* 2011;18:365–72.
- Malin SK, Kashyap SR, Hammel J, *et al*. Adjusting glucose-stimulated insulin secretion for adipose insulin resistance: an index of β -cell function in obese adults. *Diabetes Care* 2014;37:2940–6.
- Matthews DR, Hosker JP, Rudenski AS, *et al*. Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 1985;28:412–9.
- Lim U, Turner SD, Franke AA, *et al*. Predicting total, abdominal, visceral and hepatic adiposity with circulating biomarkers in Caucasian and Japanese American women. *PLoS One* 2012;7:e43502.
- Kadota K, Takamura N, Aoyagi K, *et al*. Availability of cardio-ankle vascular index (CAVI) as a screening tool for atherosclerosis. *Circ J* 2008;72:304–8.
- Chirinos JA, Segers P, Hughes T, *et al*. Large-Artery Stiffness in Health and Disease: JACC State-of-the-Art Review. *J Am Coll Cardiol* 2019;74:1237–63.
- Matsushita K, Ding N, Kim ED, *et al*. Cardio-ankle vascular index and cardiovascular disease: systematic review and meta-analysis of prospective and cross-sectional studies. *J Clin Hypertens* 2019;21:16–24.
- Yabe D, Seino Y, Fukushima M, *et al*. β cell dysfunction versus insulin resistance in the pathogenesis of type 2 diabetes in East Asians. *Curr Diab Rep* 2015;15:602.
- Nyholm B, Walker M, Gravholt CH, *et al*. Twenty-Four-Hour insulin secretion rates, circulating concentrations of fuel substrates and gut incretin hormones in healthy offspring of type II (non-insulin-dependent) diabetic parents: evidence of several aberrations. *Diabetologia* 1999;42:1314–23.
- Mather KJ, Steinberg HO, Baron AD. Insulin resistance in the vasculature. *J Clin Invest* 2013;123:1003–4.
- Kusters YHAM, Barrett EJ. Muscle microvasculature's structural and functional specializations facilitate muscle metabolism. *Am J Physiol Endocrinol Metab* 2016;310:E379–87.
- Coggins M, Lindner J, Rattigan S, *et al*. Physiologic hyperinsulinemia enhances human skeletal muscle perfusion by capillary recruitment. *Diabetes* 2001;50:2682–90.
- Vincent MA, Clerk LH, Lindner JR, *et al*. Mixed meal and light exercise each recruit muscle capillaries in healthy humans. *Am J Physiol Endocrinol Metab* 2006;290:E1191–7.
- Kim F, Pham M, Maloney E, *et al*. Vascular inflammation, insulin resistance, and reduced nitric oxide production precede the onset of peripheral insulin resistance. *Arterioscler Thromb Vasc Biol* 2008;28:1982–8.
- Barwell ND, Malkova D, Moran CN, *et al*. Exercise training has greater effects on insulin sensitivity in daughters of patients with type 2 diabetes than in women with no family history of diabetes. *Diabetologia* 2008;51:1912–9.
- Sogaard D, Østergård T, Blachnio-Zabielska AU, *et al*. Training does not alter muscle ceramide and diacylglycerol in offsprings of type 2 diabetic patients despite improved insulin sensitivity. *J Diabetes Res* 2016;2016:1–12.
- Ahn CW, Song YD, Nam JH, *et al*. Insulin sensitivity in physically fit and unfit children of parents with type 2 diabetes. *Diabet Med* 2004;21:59–63.
- Sjøberg KA, Frøsig C, Kjøbsted R, *et al*. Exercise increases human skeletal muscle insulin sensitivity via coordinated increases in microvascular perfusion and molecular signaling. *Diabetes* 2017;66:1501–10.
- Owens DR, Wragg KG, Briggs PI, *et al*. Comparison of the metabolic response to a glucose tolerance test and a standardized test meal and the response to serial test meals in normal healthy subjects. *Diabetes Care* 1979;2:409–13.
- Lovegrove JA, Pinder CJ, Jones J, *et al*. Beta-Cell responses in the offspring of non-insulin dependent diabetics after a test meal. *Biochem Soc Trans* 1993;21:21.
- Allerton TD, Irving BA, Spielmann G, *et al*. Metabolic flexibility is impaired in response to acute exercise in the young offspring of mothers with type 2 diabetes. *Physiol Rep* 2019;7:e14189.