

Supplementary Table 4. Gla-300 insulin doses according to previous treatment (insulin-naïve versus previous basal insulin) and age <75 years versus ≥ 75 years.

Gla-300 doses	Total dose (U)							
	<i>Baseline</i>		<i>Hospital discharge</i>		<i>Month 3 post-discharge</i>		<i>Month 6 post-discharge</i>	
	Mean \pm SD	N	Mean \pm SD	N	Mean \pm SD	N	Mean \pm SD	N
Overall	18.8 \pm 11.8	112	38.0 \pm 19.3	105	35.2 \pm 18.5	93	36.3 \pm 18.3	93
Insulin naïve	15.4 \pm 5.6	55	31.9 \pm 14.6	51	28.6 \pm 15.6	44	30.2 \pm 16.6	45
Previous basal insulin	22.0 \pm 14.9	57	43.7 \pm 21.5	54	41.1 \pm 18.9	49	42.1 \pm 18.2	48
Aged <75 years	21.1 \pm 13.8	65	39.0 \pm 21.1	60	37.9 \pm 20.0	53	39.9 \pm 19.9	54
Aged ≥ 75 years	15.5 \pm 7.2	47	36.6 \pm 16.8	45	31.7 \pm 15.7	40	31.5 \pm 14.8	39

Gla-300, glargine 300 U/ml; SD, standard deviation.