Effect of the Affordable Care Act on diabetes care at major health centers: newly detected diabetes and diabetes medication management

Al'ona Furmanchuk,¹ Mei Liu,² Xing Song,³ Lemuel R Waitman,³ John R Meurer,⁴ Kristen Osinski,⁵ Alexander Stoddard,⁵ Elizabeth Chrischilles,⁶ James C McClay,⁷ Lindsay G Cowell,⁸ Umberto Tachinardi,⁹ Peter J Embi,⁹ Abu Saleh Mohammad Mosa,¹⁰ Vasanthi Mandhadi,¹⁰ Raj C Shah,¹¹ Diana Garcia,¹² Francisco Angulo,¹² Alejandro Patino,¹² William E Trick,¹³ Talar W Markossian,¹⁴ Laura J Rasmussen-Torvik,¹⁵ Abel N Kho,¹ Bernard S Black¹⁶

To cite: Furmanchuk Ana, Liu M, Song X, *et al.* Effect of the Affordable Care Act on diabetes care at major health centers: newly detected diabetes and diabetes medication management. *BMJ Open Diab Res Care* 2021;9:e002205. doi:10.1136/bmjdr-2021-002205

► Supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjdr-2021-002205>).

Received 10 February 2021
Accepted 13 June 2021

© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to
Dr Al'ona Furmanchuk;
alona.furmanchuk@northwestern.edu

The adoption of the Affordable Care Act (ACA)¹ in the USA expanded health insurance for low-income Americans and took two main forms: Medicaid expansion in some states and subsidized private health insurance through insurance exchanges available in all states, with deep subsidies for persons with incomes from 138% to 250% of the federal poverty limit (FPL) in Medicaid expansion states and from 100% to 250% of the FPL in non-expansion states. Prior studies found a statistically significant slightly negative² effects of the ACA on diabetes diagnoses and controversial (from insignificantly slightly positive³ to significantly positive⁴) effects on diabetes therapies at county and state levels. We examined the effect of both forms of ACA reform on the improvement of diabetes diagnostics and management in low-income patients who had access to healthcare before the ACA expansion (2011–2013).

We used electronic health records (EHR) from 11 major academic health systems in 8 states in the USA (Illinois, Iowa, Wisconsin, Kansas, Nebraska, Missouri, Texas, Indiana). The sample (see [table 1](#) for demographics) was limited to patients aged 55–74 over 2011–2018 who used care (any encounter type) at the study facilities at least once in the pre-expansion period. Due to inconsistent depiction of insurance status in EHR, patient residence in a socially deprived⁵ census tract (see online supplemental appendix for details) was used as proxy for persons who were more likely to gain insurance under the

ACA. Therefore persons aged 55–64 from the socially deprived census tracts were the treatment group. Persons aged 65–74 from socially deprived census tracts were the control group

Panel A

Panel B

Figure 1 Annual trends in healthcare utilization outcomes before and during the Affordable Care Act Medicaid expansion (dotted vertical line). States are equally weighted. Bars are 95% CI. (A) Disadvantaged patients as per cent of all patients with newly detected diabetes. (B) Disadvantaged patients (with low socioeconomic status (SES)) as per cent of individuals with prevalent diabetes who received medical management at partnering health systems.

Table 1 Sample characteristics used to measure healthcare utilization outcomes

Sample demographic characteristics for outcome measures	2011–2013		2014–2018	
	55–64	65–74	55–64	65–74
(A) Total newly detected diabetes (305 726 patients aged 55–74 years old during 2011–2018)	73 479	56 371	90 948	84 928
% from socially deprived census tracts	41.9	34.3	34.7	28.3
Sex: % female	50.2	50.2	47.6	48.0
Race: % white	65.2	71.9	69.8	75.7
Race: % black	22.0	15.7	16.9	11.4
Race: % Asian	3.2	3.3	2.6	2.8
Race: % mixed	0.0	0.0	0.0	0.0
Race: % missing	9.6	9.1	10.7	10.1
Ethnicity: % Hispanic	11.9	8.1	9.5	6.9
Ethnicity: % missing	18.1	20.3	20.8	21.7
(B) Total with prevalent diabetes and relevant medical prescriptions (67 083 patients aged 55–74 years old during 2011–2018)	34 831	32 252		
% from socially deprived census tracts	44.1	37.6		
Sex: % female	51.0	49.9		
Race: % white	62.0	59.9		
Race: % black	15.7	11		
Race: % Asian	1.7	1.6		
Race: % mixed	0.3	0.3		
Race: % missing	6.1	5.2		
Ethnicity: % Hispanic	5.4	4.1		
Ethnicity: % missing	21.5	19.7		

The pre-ACA period is 2011–2013; the ACA period is 2014–2018. For medical management of diabetes, patients with prevalent diabetes were studied (sample is the same before and during the ACA period).
ACA, Affordable Care Act.

because they had Medicare insurance. For each age group, we studied the per cent of patients of interest with newly detected diabetes⁶ and the per cent of patients with prevalent diabetes receiving diabetes-related medications before (2011–2013) and during (2014–2018) the ACA expansion. Combined age discontinuity and difference-in-difference research design was employed.

Different from individuals who had no access to healthcare² before the ACA, our sample of patients from socially deprived tracts shows no increase in rates of newly diagnosed diabetes (figure 1). An insignificant drop of -0.72 (95% CI -3.22 to 1.77) in newly diagnosed diabetes for the treated group was detected. We have to note the identification of diabetes in the sample was not limited to ambulatory settings. This makes us conclude that the study centers may have already been using all available resources to accurately diagnose diabetes before 2014, including for low-income patients. Therefore, the ACA did not lead to an improvement in diagnostics for our sample. The decline in new diabetes cases may be a positive effect of the improved access to other preventive care⁷ services and medications during the ACA.

We also assessed whether the ACA led to low-income persons with prevalent diabetes having better access to diabetes medications. We detected an insignificant increase of 0.21 (95% CI -2.10 to 2.52) in the prescription for diabetes medications in the treatment group.

The observed trend for the prescribed diabetes medications matched the 2010–2016 dispensed medication trend detected with the Medicaid State Drug Utilization Data.³ Overall, the reported increase in diabetes medication due to the ACA tended to be modest if a ‘per enrollee’-like measure was selected as opposed to an ‘all prescriptions’⁴ one.

In summary, we would like to stress that selected health outcomes are not doing the ACA justice and, as a result, underestimating the presumed improvement in the health services for low-income patients-clients of the academic centers before the ACA implementation. Such patients would face a different level of improvement in access to care comparing with ones who were completely isolated from the healthcare system before the policy took place.

Author affiliations

¹Division of General Internal Medicine and Geriatrics, Northwestern University, Chicago, Illinois, USA

²Division of Medical Informatics, Department of Internal Medicine, University of Kansas Medical Center, Kansas City, Kansas, USA

³Division of Health Management and Informatics, University of Missouri, Columbia, Missouri, USA

⁴Institute for Health & Equity, Medical College of Wisconsin, Milwaukee, Wisconsin, USA

⁵Clinical and Translational Science Institute of Southeast Wisconsin, Medical College of Wisconsin, Milwaukee, Wisconsin, USA

⁶Department of Epidemiology, The University of Iowa College of Public Health, Iowa City, Iowa, USA

⁷Department of Emergency Medicine, University of Nebraska Medical Center, Omaha, Nebraska, USA

⁸Division of Biomedical Informatics, Department of Population and Data Sciences, Department of Immunology, The University of Texas Southwestern Medical Center, Dallas, Texas, USA

⁹Department of Biostatistics, Indiana University School of Medicine, Indianapolis, Indiana, USA

¹⁰Department of Health Management and Informatics, University of Missouri School of Medicine, Columbia, Missouri, USA

¹¹Department of Family Medicine and Rush Alzheimer's Disease Center, Rush University Medical Center, Chicago, Illinois, USA

¹²Health Research and Solutions Unit, Cook County Bureau of Health Services, Chicago, Illinois, USA

¹³Department of Medicine, Cook County Bureau of Health Services, Chicago, Illinois, USA

¹⁴Department of Public Health Sciences, Loyola University Chicago, Maywood, Illinois, USA

¹⁵Department of Preventive Medicine, Northwestern University Feinberg School of Medicine, Chicago, Illinois, USA

¹⁶Pritzker School of Law, Kellogg School of Management, Northwestern University, Chicago, Illinois, USA

Acknowledgements The Rush team would like to acknowledge the contribution of Ms Ekta Kishen and Mr J Alan Simmons of the Bioinformatics/Biostatistics Core for fulfilling the data queries used in these analyses. The Loyola team would like to acknowledge the contribution of Ms Susan Zelisko of Information Technologies for conducting the data queries used in these analyses. The Northwestern team would like to thank Ms Anna Nasiri for providing valuable insight about the Northwestern Medicine EHR. The UTSW team would like to acknowledge the contribution of Mr Phillip Reeder for conducting the data queries used in this project. Data from Indiana were produced by the Regenstrief Data Services team. Daniel Hood, the Regenstrief Project Manager for GPC, led the data extraction processes for this project.

Contributors BSB, ANK and AF conceived and designed the study. All authors interpreted the data, critically reviewed the manuscript for important intellectual content and approved the final manuscript. AF and BSB developed the analytical plan. AF performed the statistical analyses. AF and BSB drafted the initial manuscript. ANK supervised the study. AF, ML, XS, LRW, JRM, KO, AS, EC, LGC, RCS, JCM, UT, VM, ASMM, DG, FA, AP, WET, TWM and ANK contributed to data acquisition. AF and BSB contributed to analysis of data. RCS, ML, XS, JRM, KO, AS, EC, JCM, UT, ASMM, WET, TWM, LR-T, PJE and ANK contributed to interpretation of data and manuscript writing.

Funding This work is supported by the Centers for Disease Control and Prevention (CDC) and the National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases (grant no 5U18DP006120-05-00) as part of the

Natural Experiments for Translation in Diabetes 2.0 (NEXT-D2) collaborative. The study funders had no role in the study design, data collection, data analysis, data interpretation or the writing of this article. The corresponding author had full access to all the data in the study and had final responsibility for the decision to submit for publication.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval The CAPRICORN and GPC institutional review boards approved this study (IRB#16030901 and IRB#STUDY00004015, respectively).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as online supplemental information.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- 1 US Gov. *The Patient Protection and Affordable Care Act. Public Law 111-148, 124 STAT 855*, 2010.
- 2 Palanki R, Chamrathy S, Palanki S. Impact of the Affordable care act on diabetes diagnoses in the United States: a county-level analysis. *Economic Affairs* 2021;41:111–22.
- 3 Cher BAY, Morden NE, Meara E. Medicaid expansion and prescription trends: opioids, addiction therapies, and other drugs. *Med Care* 2019;57:208–12.
- 4 Ghosh A, Simon K, Sommers BD. The Effect of Health Insurance on Prescription Drug Use Among Low-Income Adults: Evidence from Recent Medicaid Expansions. *J Health Econ* 2019;63:64–80.
- 5 Townsend P. Deprivation. *J Soc Policy* 1987;16:125–46.
- 6 Nichols GA, Desai J, Elston Lafata J, et al. Construction of a multisite Datalink using electronic health records for the identification, surveillance, prevention, and management of diabetes mellitus: the SUPREME-DM project. *Prev Chronic Dis* 2012;9:E110.
- 7 IRS. IRS notice 2013-57, 2013. Available: <https://www.irs.gov/pub/irs-drop/n-13-57.pdf> [Accessed Apr 2021].