

eTable 1. Baseline demographic and clinical characteristics of patients missing vs not missing metabolic data

	Missing n=1814	Not Missing n=1574	P-value
Age (yr)	58.9±12.5	58.0±12.5	0.031
Male	68.5%	70.1%	0.312
White	67.1%	71.9%	0.002
Married	52.1%	55.3%	0.067
Lives alone	25.1%	23.3%	0.221
High school education	79.8%	82.3%	0.063
Avoids care due to cost	24.8%	26.8%	0.203
Body mass index	28.8±6.1	29.0±5.9	0.303
Current smoking	41.5%	44.8%	0.063
Hypertension	61.3%	61.7%	0.817
Prior myocardial infarction	18.6%	17.9%	0.586
Prior angioplasty	16.2%	17.7%	0.260
Prior bypass surgery	9.3%	8.6%	0.493
Peripheral vascular disease	3.9%	4.1%	0.750
Prior stroke	4.1%	4.1%	0.984
Chronic heart failure	7.6%	4.7%	<0.001
Estimated GFR (mL/min/1.73m ²)	78.1±26.4	79.2±25.5	0.222
Chronic lung disease	7.2%	6.4%	0.317
Depression	14.9%	18.8%	0.003
GRACE discharge risk score	99.6±29.9	95.9±29.1	<0.001
ST-elevation MI	44.1%	50.1%	<0.001
Peak troponin (ng/dL)	29.4±68.1	32.5±79.1	0.234
Multivessel disease	43.2%	49.0%	<0.001
LV dysfunction	17.4%	17.4%	0.986
HbA1c (%)	6.2±2.0	6.1±1.4	0.032
Missing	678	0	
Fasting glucose (mg/dL)	121.8±41.1	125.2±41.0	0.036
Missing	802	0	
Total cholesterol (mg/dL)	176.2±44.6	178.4±52.8	0.212
Triglycerides (mg/dL)	139.5±120.8	149.1±152.8	0.051
HDL-C (mg/dL)	42.5±14.6	41.3±14.1	0.028
LDL-C (mg/dL)	107.1±39.1	107.7±41.9	0.705